

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Confirm* 2F

EPA Reg. No. 62719-420

For Control of Insects Infesting Almond

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Confirm* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Confirm 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Confirm 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
peach twig borer	16 - 30 (0.25 - 0.47 lb ai/acre)	<p>Spring (overwintering) generation: Make 1 to 2 applications at bloom to petal fall depending on infestation level.</p> <p>Summer generation: Apply at peak moth flight (200-300 DD, base 50°F following biofix) for each generation. Reapply at 10-to 14-day intervals under high pressure or sustained moth flight.</p> <p>Use a higher rate in the rate range for extended residual effectiveness, higher pest infestation levels, large trees, or heavy dense foliage.</p>	<ul style="list-style-type: none"> ▪ Do not apply more than 30 fl oz of Confirm 2F per application or exceed 122 fl oz of Confirm 2F per season. ▪ Pre-harvest Interval: Do not harvest within 14 days of application. ▪ Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
navel orangeworm	18 - 30 (0.28 to 0.47 lb ai/acre)	Apply at initiation of hull split and reapply 10 to 14 days later. For heavy infestation, reapply 10 to 14 days after the second application.	<div style="border: 1px solid black; padding: 5px; transform: rotate(-2deg);"> <p style="text-align: center; font-weight: bold; font-size: 1.2em;">ACCEPTED</p> <p style="text-align: center; font-weight: bold; font-size: 1.2em;">FEB 10 2004</p> <p style="font-size: 0.8em;">Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-420</p> </div>

Ground Application: Apply a minimum of 50 gallons per acre by conventional ground equipment to almond trees 4th leaf or younger. Apply a minimum of 100 gallons per acre to almond trees 5th leaf or older. Ground speed of the sprayer should not exceed 2 mph.

Aerial Application: Apply in a minimum of 20 gallons per acre. **Note:** Do not aerially apply Confirm 2F to almonds if the tree size or foliage density prohibits thorough, uniform coverage of the entire tree canopy.

2/2

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R809-001

EPA-accepted __/__/__

Initial printing.