

MAY 4 2004


Barbara J. Kaminski
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

Subject: Supplemental Label For;
Confirm 2F
EPA Registration No. 62719-420
Letter dated February 19, 2004
Letter dated April 6, 2004

Dear Ms. Kaminski:

The supplemental labels for almond and canola submitted with your letters cited above are acceptable. A stamped copy of each supplemental label is enclosed for your records. You are reminded that this supplemental labeling must be made a part of the master label for this product within 18 months.

Sincerely,



Richard Gebken
Acting Product Manager 10
Insecticide Branch
Registration Division (7505C)



2/5

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Confirm* 2F

EPA Reg. No. 62719-420

For Control of Insects Infesting Almond

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of a pplication.
- Read the label affixed to the container for Confirm* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Confirm 2F according to this supplemental labeling is subject to all u se precautions and limitations imposed by the label affixed to the container for Confirm 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
peach twig borer	16 - 30 (0.25 - 0.47 lb ai/acre)	<p>Spring (overwintering) generation: Make 1 to 2 applications at bloom to petal fall depending on infestation level.</p> <p>Summer generation: Apply at peak moth flight (200-300 DD, base 50°F following biofix) for each generation. Reapply at 10-to 14-day intervals under high pressure or sustained moth flight.</p> <p>Use a higher rate in the rate range for extended residual effectiveness, higher pest infestation levels, large trees, or heavy dense foliage.</p>	<ul style="list-style-type: none"> • Do not apply more than 30 fl oz of Confirm 2F per application or exceed 122 fl oz of Confirm 2F per season. • Pre-harvest Interval: Do not harvest within 14 days of application. • Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
navel orangeworm	18 - 30 (0.28 to 0.47 lb ai/acre)	Apply at initiation of hull split and reapply 10 to 14 days later. For heavy infestation, reapply 10 to 14 days after the second application.	

ACCEPTED with COMMENTS
In EPA Letter Dated:
MAY 4 2004
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-420

Ground Application: Apply a minimum of 50 gallons per acre by conventional ground equipment to almond trees 4th leaf or younger. Apply a minimum of 100 gallons per acre to almond trees 5th leaf or older. Ground speed of the sprayer should not exceed 2 mph.

Aerial Application: Apply in a minimum of 20 gallons per acre. **Note:** Do not aerially apply Confirm 2F to almonds if the tree size or foliage density prohibits thorough, uniform coverage of the entire tree canopy.

3/5

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R809-001

EPA-accepted 02/10/04

Initial printing.

4/5

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Confirm* 2F

EPA Reg. No. 62719-420

For Control of Insects Infesting Canola

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of a application.
- Read the label affixed to the container for Confirm* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Confirm 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Confirm 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
beet armyworm (<i>Spodoptera exigua</i>) bertha armyworm (<i>Mamestra configurata</i>)	8 (0.12 lb ai/acre)	For early season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	<ul style="list-style-type: none"> ▪ Do not apply more than 16 fl oz of Confirm 2F per application or exceed 64 fl oz of Confirm 2F per season. ▪ Pre-harvest Interval: Do not harvest within 14 days of application. ▪ In the state of New York, this product is prohibited from use in Nassau and Suffolk Counties. ▪ Refer to Rotational Crop Restrictions in the body of the product label.
	8 - 16 (0.12 to 0.25 lb ai/acre)	For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, re-apply in 10 to 14 days to protect new growth until moth flights and/or hits subside.	

ACCEPTED
with **COMMENTS**
EPA Letter Dated:

MAY 4 2004
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-420

Specific Use Directions

Equipment and spray volume should be carefully adjusted to assure thorough uniform coverage of infested parts of the crop.

Ground Application: Apply a minimum of 8 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 10 gallons per acre to densely foliated or difficult to cover crop to ensure thorough coverage.

Aerial Application: Apply in a minimum of 5 gallons per acre.

5/5

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R809-005
EPA-accepted 08/11/03
Replaces D06-809-022.

