62719 - 420

8/16/2007

174

United States Environmental Protection Agency

Ken Racke, Ph. D. Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054

Subject: Supplemental Label for; Confirm 2F EPA Registration No. 62719-420 Letter dated 8 November 2005

Dear Mr. Racke:

The supplemental label for citrus submitted with your letter cited above is acceptable. A stamped copy of the supplemental label is enclosed for your records. You are reminded that this supplemental labeling must be made a part of the master label for this product within 18 months.

Sind

ALIG 16

Richard Gebken Product Manager 10 Insecticide Branch Registration Division (7505P)

Enclosures 62719-420 RN 16August2007

CONCURRENCES										
:	H7505P									
<	MSUAREZ									
	Aug 16, 2007									

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

D3B / Confirm 2F / Prop Sec 3 Supp Citrus / 06-13-02 file: D06-809-XXX Confirm 2F Sec 3 Citrus 13June02d.doc

Confirm* 2F

EPA Reg. No. 62719-420

Proposed Section 3 supplemental labeling: Proposed supplemental labeling for use of Confirm 2F for control of insects infesting citrus.

*Trademark of Dow AgroSciences LLC

Supplemental



Dow AgroSciences LLC

Labeling

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Confirm* 2F

EPA Reg. No. 62719-420

For Control of Insects Infesting Citrus[†]

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Confirm* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Confirm 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Confirm 2F.

Directions for Use

[†]Citrus, including but not limited to: Calamondin, Citrus Citron, Citrus Hybrids (Chironja, Tangelo, Tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Pummelo, Satsuma Mandarin, Sour Orange, Sweet Orange

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

	Application Rate		
Pest	(fl oz/acre)	Application Timing	Restrictions
citrus cutworm	8 - 16 (0.125 – 0.25 lb ai/acre)	Make application at the initiation of egg lay, usually in the bloom period. Re-treat in 14-21 days to protect newly expanded fruit against heavy infestations. Additional applications may be required to protect against prolonged moth flight and egg laying.	 Do not apply more than 80 fl oz of Confirm 2F per season. Pre-harvest Interval: Do not harvest within 14 days of application.
citrus leafminer		Apply at the initiation of new flush to protect newly expanding leaves.	
omfilverous leafroller orangedog orange tortrix		Apply at first signs of feeding damage or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	A C C E P T E D 16 August 2007 Under the Federal Insecticide, Pungicide, and Rodentiside Act,
fruittree (cafrolle: western tussock moth		Apply at the initiation of egg lay or at the first sign of larval infestation. A second application may be required for control under heavy pressure or sustained moth flight	as amended, for the pesticide registered under EPA Reg. No. 62719-420

Specific Use Directions

Equipment and spray volume should be carefully adjusted to assure thorough uniform coverage of infested parts of the crop.

Ground Application: Apply a minimum of 50 gallons per acre to trees 10 feet tall or less by conventional ground equipment. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre.

Aerial Application: Apply a minimum of 20 gallons per acre in equipment that has been properly patterned and calibrated for environmental conditions at the site. This method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Spray Adjuvants: The use of Latron B-1956* spreader/sticker or similar spreader/sticker is recommended to maximize coverage and retention of the spray material. Addition of a crop oil or other adjuvant with strong penetrant properties may improve performance against citrus leafminer.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

D38'/Cenfirm 2F / Prop Sec 3 Supp Citrus / 06-13-02 D06-809-XXX EPA-accepted __/_/__ Initial printing.