



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

JUL 25 2007

Dr. Raymond S. Brinkmeyer  
Dow Agro Sciences LLC  
9330 Zionsville Road  
Indianapolis, Indiana 46268

Subject: Indar® 75WSP Fungicide  
EPA Registration Number 62719-421  
Your master label amendment and supplemental labels for  
Citrus Fruits and Sugarbeet submitted by e-mail on May  
18, 2007 (submission S811479), amending the same types  
of labels submitted on April 3, 2007 (submissions  
S808157 and S808160)

We have reviewed the subject amended labeling, submitted in  
connection with registration under the Federal Insecticide,  
Fungicide, and Rodenticide Act (FIFRA), as amended. The amended  
labeling is acceptable provided that:

1. You make the following changes to the labeling.
  - a. In the "Chemigation:" subsection on page 8 in the  
master label delete "or product bulletins" from the end  
of the sentence. If product bulletins are referenced  
in the label, they are part of the labeling and the  
Agency must review and approve them first.
  - b. In the first sentence in the "Citrus Fruits"  
specific use directions section in the master label and  
the "Citrus Fruits" supplemental label change "and  
other commodities included" to "and other crops  
included".
  - c. In the "Cranberry" specific use directions section,  
the "Use Directions" column for "fairyring complex  
(suppression)" in the table reads "Measure ring  
diameter and add 10 ft to the diameter. Begin  
applications at bud break and repeat once if


2720

necessary...". Add to these directions an explanation of what is done with or what is the significance of the number that results from the ring diameter plus 10 feet.

2. You submit one copy of your final printed master and supplemental labels to the Agency before you release product bearing this amended labeling for shipment.

If you have any questions about this letter, please contact John Bazuin at (703)305-7381 or bazuin.john@epa.gov.

Sincerely yours,



Tony Kish, Product Manager (22)  
Fungicide Branch  
Registration Division (7505C)

Attachments: Master label stamped "ACCEPTED with COMMENTS"  
Two supplemental labels stamped "ACCEPTED"

3 8 20

Base Label:

# Indar<sup>®</sup> 75WSP

Fungicide

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

JUL 25 2007

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

62,719-421

<b>Group</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-  
a-phenyl-1H-1,2,4-triazole-1-propanenitrile . 75%

Other Ingredients ..... 25%

Total ..... 100%

U.S. Patent No. 4,366,165

**Keep Out Of Reach Of Children**

## CAUTION

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**Harmful If Swallowed Or Absorbed Through The Skin**

**Avoid contact with skin, eyes or clothing.**

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### First Aid

4 8 20

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night for emergency treatment information.

**Environmental Hazards**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-421

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

**Net Weight 1 lb**  
**8 x 2 oz. Water-Soluble Pouches**

Label Booklet:

# Indar<sup>®</sup> 75WSP

## Fungicide

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-

a-phenyl-1H-1,2,4-triazole-1-propanenitrile . 75%

Other Ingredients ..... 25%

Total ..... 100%

U.S. Patent No. 4,366,165

**Keep Out Of Reach Of Children**

### CAUTION

#### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for Precautionary Statements and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-421

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC  
**Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.**

**Net Weight 1 lb.  
8 x 2 oz. Water-Soluble Pouches**

(Page 1 through end):

<b>Table of Contents</b>	<b>Page</b>
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
Engineering Controls	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Resistance Management	-
Use Rate Determination	-
Mixing	-
Handling for Water Soluble Pouches	-
Application	-
Orchard Use	-
Aerial Spray Drift Management	-
Compatibility	-
Plantback Intervals	-
Uses	-
Almond	-
Apple	-
Blueberry	-
Citrus Fruits	-
Cranberry	-
Peanut	-
Pecan	-
Stone Fruits	-
Sugarbeets	-
Wheat	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals**

**CAUTION**

**Harmful If Swallowed Or Absorbed Through The Skin**

**Avoid contact with skin, eyes or clothing.**

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

When handlers use closed systems enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

**First Aid**

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

**Environmental Hazards**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

---

**Directions for Use**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry area above freezing. The water soluble pouch may become brittle at storage temperatures below 32°F but the fungicide is not affected. Do not remove the water soluble pouches from container except for immediate use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Dispose of the empty outer foil pouch in the trash, as long as WSP is unbroken.

**Steps to be Taken in Case Material is Released or Spilled:** Keep dust to a minimum. Appropriate protective equipment must be worn when handling a spill of this material. Remove contaminated clothing promptly. Wash exposed skin areas with soap and water immediately after exposure. Contain spills immediately with inert materials (e.g., sand, earth). Transfer spilled material and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Material Safety Data Sheet.

**General Information**

Indar® 75WSP fungicide is a systemic fungicide with preventative and curative properties used for the control of specific diseases listed on this label. For optimal disease control with Indar 75WSP, apply sprays on a preventative spray program.

A non-polymer containing spray adjuvant approved for use with registered pesticide products may be added to spray solutions according to manufacturer's use instructions to achieve optimum disease control.

**Resistance Management**

Indar 75WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Indar 75WSP should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.



## Use Rate Determination

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum suggested use rates can be used, while maximum suggested label rates and shortened spray schedules are recommended for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the suggested label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

## Mixing

Always place Indar 75WSP into solution prior to adding co-applied agricultural chemicals registered for use on specific crop or crops being treated. Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending upon the water temperature and the degree of agitation, the pouches should dissolve completely within approximately ten minutes from the time they are added to the water. **Note:** Reduced product efficacy may occur if water containing suspended soil particles such as water from ponds, streams or unlined ditches is used.

## Handling for Water Soluble Pouches

Do not remove water soluble pouch from overwrap except for immediate use. Do not allow water soluble pouch to come into contact with water prior to use. Do not handle water soluble pouch with wet hands or wet gloves. Do not open water soluble pouches. Partial use of pouches is not allowed. Carefully reseal overwrap containing unopened water soluble pouches and protect package from moisture.

## Application

Do not make ground or aerial applications within 75 feet of streams, rivers, ponds, lakes or reservoirs.

**Ground:** Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. For tree fruits and nuts the same amount of Indar 75WSP should be applied per acre in either dilute or concentrate sprays.

**Aerial:** Apply in a minimum of 5 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Flagmen and loaders should wash thoroughly before eating, drinking or smoking and at the end of each day's spray operation.

## Orchard Use

Apply Indar 75WSP to orchards using a dilute or concentrate spray.

**Dilute Spray:** Dilute or thorough coverage applications are based upon the amount of spray solution necessary to thoroughly wet the trees until spray run-off. To determine the number of gallons of dilute spray required per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance.

**Concentrate Spray:** Concentrate spray applications assume all of the tree is uniformly covered with spray solution but not to the point of run-off. A lower spray volume is used to deliver the same application rate per acre as used for a dilute spray. A minimum of 50 gallons of spray solution per acre is recommended for concentrate sprays with Indar 75WSP.

## Aerial Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

**Chemigation:** Do not apply this product by chemigation to labeled crops except as specified in Dow AgroSciences supplemental labeling or product bulletins.

**Compatibility**

Indar 75WSP is compatible with most commonly used agriculture fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. For any registered adjuvant, follow the manufacturer's use instructions to achieve proper spray concentrations. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. Indar 75WSP is compatible with boron and spray oils; however, the water soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

**Plantback Intervals**

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

**Uses**

**Almond**

Disease	Indar 75WSP		Use Directions	Restrictions
	(oz/100 gal) <sup>†</sup>	(oz/acre)		
blossom blight ( <i>Monilinia</i> spp.)	0.4 – 0.67	1.25 - 2 (0.06 - 0.094 lb active)	Begin applications at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>• Do not make more than 3 applications or apply more than 6 oz of Indar 75WSP (0.28 lb active) per acre per year.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 160 days of harvest.</li> <li>• <b>Minimum Interval Between Sprays:</b> 10 days</li> </ul>

<sup>†</sup> Based on a standard of 300 gallons of dilute spray per acre or the equivalent amount of product per acre as a concentrate spray.

11 7 20

**Apple**

Disease	Indar 75WSP (oz/acre) <sup>†</sup>	Use Directions	Restrictions
flyspeck ( <i>Zygothia jamaicensis</i> ) powdery mildew ( <i>Podosphaera leucotricha</i> ) rusts ( <i>Gymnosporangium</i> spp.) scab ( <i>Venturia inaequalis</i> ) sooty blotch ( <i>Gloeodes pomigena</i> )	2 - 2.67 (0.094 - 0.125 lb active)	Begin application at green tip or when environmental conditions become favorable for disease development. Apply on a 7- to 10-day schedule through the first cover spray; thereafter, apply on a 10- to 14-day schedule. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>Do not make more than 4 applications or apply more than 10.67 oz of Indar 75WSP (0.5 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> <li>Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.</li> </ul>

<sup>†</sup> Maintain per acre use rate regardless of spray volume per acre.

**Blueberry**

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
alternaria ( <i>Alternaria</i> spp.) anthracnose ( <i>Colletotrichum gloeosporioides</i> ) leaf spot and blotch ( <i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) mummy berry disease ( <i>Monilinia vaccinii-corymbosi</i> ) phomopsis twig blight and fruit rot ( <i>Phomopsis vaccinii</i> ) powdery mildew ( <i>Microsphaera vaccinii</i> ) rusts ( <i>Pucciniastrum</i> spp.)	2 - 2.67 (0.094 - 0.125 lb active)	For mummy berry and phomopsis twig blight, begin applications at early green tip and make subsequent applications at intervals of 8 to 14 days. For control of powdery mildew and other diseases listed, begin applications prior to or at onset of disease development and repeat at 8- to 14-day intervals. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>Do not make more than 4 applications or apply more than 8 oz of Indar 75WSP (0.38 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> <li>Do not graze livestock in treated fields or offer treated material as a livestock feed item.</li> </ul>

**Citrus Fruits**

Including: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange (Sweet, Sour and Mandarin), Pummello, and Tangerine and other commodities included in the citrus fruits crop grouping

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Indar 75WSP with a nonionic surfactant or spray oil at the manufacturer's recommended rate and making thorough coverage applications.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
---------	-----------------------	----------------	--------------

greasy spot ( <i>Mycosphaerella citri</i> )	2.67 (0.125 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application or for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> <li>Do not make more than 3 applications or apply more than 8.01 oz of Indar 75WSP (0.375 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Up to day of harvest.</li> <li><b>Minimum Interval Between Sprays:</b> 21 days.</li> <li>Do not allow livestock to graze in treated areas.</li> <li>Do not apply Indar 75WSP with polymer based spray adjuvants.</li> </ul>
scab ( <i>Elsinoe fawcettii</i> )		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold ( <i>Capnodium</i> spp.)		Applications made for control of greasy spot or scab will also control sooty mold.	

**Cranberry**

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, it is recommended to use an agriculturally registered non-polymer non-ionic surfactant at the manufacturer's recommended rate.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
cottonball tip blight ( <i>Monilinia oxycocci</i> )	2 - 4 (0.094 - 0.188 lb active)	Begin applications when 50% of the shoots have begun to elongate. Refer to your local spray recommendation guides for more specific application timing information.	<ul style="list-style-type: none"> <li>Do not make more than 4 applications or apply more than 16 oz of Indar 75WSP (0.75 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> </ul>
cranberry fruit rot complex (1)		Begin applications prior to bloom at the onset of disease. Continue on a 10- to 14-day spray schedule depending upon local conditions. Refer to your local recommendation guides for more specific application information.	
fairyring complex (suppression)	4 (0.188 lb active)	Measure ring diameter and add 10 ft to the diameter. Begin applications at bud break and repeat once if necessary 14 days later. Apply in 30 to 100 gallons of water to the affected area. Irrigation for 1 to 2	

		hours following application is advisable to ensure penetration to the base of the plant.	
--	--	--	--

1. Cranberry fruit rot disease complex may contain one or more of the following fungal pathogens: *Allantophomopsis lycopodina*, *Botryosphaeria vaccinii*, *Coleophoma empetri*, *Diaporthe vaccinii*, *Glomerella cingulata-vaccinii*, *Godronia cassandrae*, *Phyllosticta vaccinii*, *Physalospora vaccinii*, *Colletotrichum* spp. , *Phomopsis vaccinii* and *Strasseria geniculata*.

**Peanut**

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
early leaf spot ( <i>Cercospora arachidicola</i> )	1.25 - 2 (0.062 - 0.094 lb active)	Begin foliar applications approximately 4 weeks after planting or at first sign of disease development and continue application on a 10- to 14-day schedule.	<ul style="list-style-type: none"> <li>Do not make more than 4 applications or apply more than 10.67 oz of Indar 75WSP (0.5 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> <li>Do not feed hay or threshings or allow livestock to graze in treated areas.</li> </ul>
late leaf spot ( <i>Cercosporidium personatum</i> )	2 - 2.67 (0.094 - 0.125 lb active)		

**Pecan**

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
downy spot ( <i>Mycosphaerella caryigena</i> ) gnomonina leafspot ( <i>Gnomonia dispersa</i> ) leaf blotch ( <i>Mycosphaerella dendroides</i> ) leaf scorch powdery mildew ( <i>Micosphaera penicillata</i> ) scab ( <i>Cladosporium caryigenum</i> ) vein spot ( <i>Gnomonia nerviseda</i> )	2 - 2.5 (0.09 - 0.12 lb active)	Begin applications at bud break and continue at 10- to 14-day intervals through pollination. Apply at 14- to 21-day intervals through the cover sprays. Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>Do not make more than 6 applications or apply more than 16 oz of Indar 75WSP (0.75 lb active) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply after shuck split or within 28 days of harvest.</li> <li>Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.</li> <li>Do not apply after shuck split.</li> </ul>

**Stone Fruits**

Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. For optimum disease control, a non-polymer spray adjuvant/wetting agent labeled for use on food crops should be added to spray solutions at the manufacturers' recommended rate.

Crop	Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
apricot cherry nectarine peach	blossom blight ( <i>Monilinia</i> spp.)	2 (0.094 lb active)	Begin applications at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and at petal-fall.	<ul style="list-style-type: none"> <li>Do not make more than 8 applications or apply more than 16 oz of Indar 75WSP (0.75 lb active) per acre per season.</li> <li><b>Preharvest Interval:</b> Up to day of harvest.</li> <li>Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.</li> </ul>
	fruit brown rot ( <i>Monilinia</i> spp.)		Begin applications 2 to 3 weeks before harvest using a 7- to 10-day spray interval.	
cherry	leaf spot ( <i>Blumeriella</i> spp.)	Follow blossom blight schedule and continue application at 10- to 14-day intervals. Additional foliar applications may be made after harvest.		
peach	scab ( <i>Cladosporium</i> spp.)	Begin applications at shuck split. Make 2 to 3 subsequent thorough coverage applications at 10- to 14-day intervals.		
plum prune	blossom blight ( <i>Monilinia</i> spp.)	2 (0.094 lb active)	Refer to apricot	
	fruit brown rot ( <i>Monilinia</i> spp.)		<ul style="list-style-type: none"> <li>Do not make more than 4 applications or apply more than 8 oz of Indar 75WSP (0.38 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Up to day of harvest.</li> <li>Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.</li> </ul>	

**Sugarbeets**

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
cercospora leaf spot ( <i>Cercospora beticola</i> ) powdery mildew ( <i>Erysiphe</i> spp.)	2.67 (0.125 lb active)	Apply Indar 75WSP at intervals of 14 to 21 days beginning when conditions are favorable for the development of disease. For resistance management, alternate applications of Indar 75WSP with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant such as crop oil concentrate that is labeled	<ul style="list-style-type: none"> <li>Do not make more than 2 applications or apply more than 5.34 oz of Indar 75WSP (0.25 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> </ul>

		<p>for crop use may be added to the spray mixture at the manufacturer's recommended rate. Add an adjuvant to the spray solution last.</p> <p>For best spray coverage of foliage, apply aerially in a recommended minimum of 5 to 7 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.</p>	
--	--	---	--

**Wheat**

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
glume blotch ( <i>Septoria nodorum</i> ) leaf blight, spot blotch, barley scald ( <i>Rhynchosporium</i> spp.) leaf blotch ( <i>Septoria tritici</i> ) powdery mildew ( <i>Erysiphe graminis</i> ) rusts ( <i>Puccinia</i> spp.) tan spot, net blotch ( <i>Pyrenophora</i> spp.)	1.25 (0.06 lb active)	Apply anytime through heading (Feekes growth stage 10.3 to 10.5) on a 7- to 21-day schedule. Apply aerially in a recommended minimum of 5 gallons of water per acre or by ground in a recommended minimum of 15 gallons of water per acre.	<ul style="list-style-type: none"> <li>Do not make more than 2 applications or apply more than 2.5 oz of Indar 75WSP (0.12 lb active) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply after completion of flowering (Feekes growth stage 10.5) or within 35 days of harvest.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest for forage and hay.</li> </ul>

**Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**Warranty Disclaimer**

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

16 7 20

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

To the extent permitted by law, the terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC  
EPA accepted \_\_\_/\_\_\_/\_\_\_



17/20

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Indar<sup>®</sup> 75WSP EPA Reg. No. 62719-421

### Disease Control in Citrus Fruits<sup>†</sup>

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Indar<sup>®</sup> 75WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Indar 75WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 75WSP.

#### Directions for Use

<sup>†</sup>Including: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange (Sweet, Sour and Mandarin), Pummello, and Tangerine and other commodities included in the citrus fruits crop grouping

Refer to product label for Indar 75WSP for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Indar 75WSP with a nonionic surfactant or spray oil at the manufacturer's recommended rate and making thorough coverage applications.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
greasy spot ( <i>Mycosphaerella citri</i> )	2.67 (0.125 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application or for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> <li>• Do not make more than 3 applications or apply more than 8.01 oz of Indar 75WSP (0.375 lb active) per acre per year.</li> <li>• <b>Preharvest Interval:</b> Up to day of harvest.</li> <li>• <b>Minimum Interval Between Sprays:</b> 21 days.</li> <li>• Do not allow livestock to graze in treated areas.</li> <li>• Do not apply Indar 75WSP with polymer based spray adjuvants.</li> </ul>
scab ( <i>Elsinoe fawcettii</i> )		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold ( <i>Capnodium</i> spp.)		Applications made for control of greasy spot or scab will also control sooty mold.	

18/20

**Resistance Management**

Indar 75WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Indar 75WSP should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

**JUL 25 2007  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

62719-421

®Trademark of Dow AgroSciences LLC

R845-077

EPA accepted:   /  /  

Replaces R845-053.

19/20

# Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

## Indar<sup>®</sup> 75WSP

EPA Reg. No. 62719-421

### Disease Control in Sugarbeets

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Indar<sup>®</sup> 75WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Indar 75WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 75WSP.

#### Directions for Use

Refer to product label for Indar 75WSP for Mixing and Application Instructions.

Disease	Indar 75WSP (oz/acre)	Use Directions	Restrictions
cercospora leaf spot ( <i>Cercospora beticola</i> ) powdery mildew ( <i>Erysiphe</i> spp.)	2.67 (0.125 lb active)	Apply Indar 75WSP at intervals of 14 to 21 days beginning when conditions are favorable for the development of disease. For resistance management, alternate applications of Indar 75WSP with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant such as crop oil concentrate that is labeled for crop use may be added to the spray mixture at the manufacturer's recommended rate. Add an adjuvant to the spray solution last. For best spray coverage of foliage, apply aerially in a recommended minimum of 5 to 7 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> <li>• Do not make more than 2 applications or apply more than 5.34 oz of Indar 75WSP (0.25 lb active) per acre per year.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> </ul>

#### Resistance Management

Indar 75WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Indar 75WSP should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

90/30

**Plantback Intervals**

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

JUL 25 2007

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

62719-426

®Trademark of Dow AgroSciences LLC

R845-078

EPA accepted: \_\_\_/\_\_\_/\_\_\_

Replaces R845-057.