

62719-424

11/02/2001

Page 1 of 67



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

NOV - 2 2001

Ms. Barbara J. Kaminski
Regulatory Manager
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268

Dear Ms. Kaminski:

Subject: GOAL® 2XL Herbicide
EPA Registration No. 62719-424
Applications and Your Letter Dated September 26 2001,
Transitive Labeling from Rohm & Haas Company to
Dow AgroSciences: Company Name and Address, Warranty
Statements, Trademark Ownership and Formatting;
Request To Amend Registration as Described in the
11 "Proposed Changes by Amendment"

The proposed subject labeling amendments have been reviewed and found acceptable under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, provided that you:

1. Revise all tank mix partners' names listed on this label to correctly give complete names of products; e.g., if the name is Dual[®] Herbicide or Gramoxone[®] Herbicide, capitalize the word "Herbicide" in these names. In many of these names you have used the lower case for the word "herbicide".
2. On page 46, revise the "Tank Mixes with Goal 2XL" section. The claims could imply that GOAL 2XL could be tank mixed with any other pesticide product. PR Notice 82-1 requires that you name the tank mix partners. Alternatively, add a section to this labeling that gives directions for determining compatibility with GOAL 2XL when tank mixing with another pesticide product or a carrier other than water.
3. Submit one (1) copy of the final printed label prior to your shipment of this pesticide product under the enclosed revised stamped label.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment under this labeling

2 7 67

-2-

constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The issue of listing trademarked name ownership on labels of pesticide products registered by this Agency is pending with the EPA, Office of General Counsel; as requested in your letter of August 2, 2001 to Ms. Michele Knorr. We are accepting at this time your removal of the list of trademarks with their owners. The list may have to be re-instated after this issue has been resolved.

Sincerely yours,

for Eugene M. Wilson
Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

Base Label:

(Logo) Dow AgroSciences

Goal* 2XL

Active Ingredient	
oxyfluorfen: 2-chloro-1-(3-ethoxy-4-nitrophenoxy)	
4-(trifluoromethyl).....	23%
Inert Ingredients.....	77%
Total.....	100%

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

NOV - 2 2001

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

62719-424

Contains 2 pounds active ingredient per gallon.
Contains petroleum distillates

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Causes Skin Irritation • Harmful If Swallowed Or Absorbed Through The Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes, on skin or on clothing. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Protective eyewear
- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (such as Nitrile, Butyl, Neoprene, and/or Barrier Laminate)
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

4 8 67

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove contaminated clothing and wash clothing before reuse.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. This product may pose an aspiration pneumonia hazard. Contains petroleum distillates

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas. Runoff may be hazardous to aquatic organisms in neighboring areas.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

Shake Well Before Using

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

5 7 67

F2B / Goal 2XL / Amend / 09-25-01

page 3

*Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents __ gal

6 7 67

Label Booklet:

(Logo) Dow AgroSciences

Goal* 2XL

Active Ingredient

oxyfluorfen: 2-chloro-1-(3-ethoxy-4-nitrophenoxy)
4-(trifluoromethyl)..... 23%

Inert Ingredients..... 77%

Total..... 100%

Contains 2 pounds active ingredient per gallon

Contains petroleum distillates

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for Precautionary Statements and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

Shake Well Before Using

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-424

EPA Est. 707-PA-01

*Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.
Made in USA

Shake Well Before Using

Herbicide

Net Contents __ gal

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Chemigation	-
Mixing	-
Crop Uses	-
Artichokes	-
Broccoli/Cabbage/Cauliflower	-
Cacao	-
Citrus(Non-Bearing)	-
Coffee	-
Conifer and Selected Deciduous Trees	-
Corn	-
Cotton	-
Cottonwood	-
Eucalyptus	-
Fallow Bed (Cotton/Soybeans)	-
Fallow land	-
Garbanzo Beans	-
Garlic	-
Guava (Hawaii Only)	-
Horseradish	-
Jojoba	-
Mint	-
Onions	-
Onions Grown for Seed	-
Papaya (Hawaii Only)	-
Soybeans	-
Taro	-
Treefruit/Nuts/Vines	-
General Use Restrictions	-
Weed Control List	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes Substantial But Temporary Eye Injury • Causes Skin Irritation • Harmful If Swallowed Or Absorbed Through The Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes, on skin or on clothing. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Protective eyewear
- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (such as Nitrile, Butyl, Neoprene, and/or Barrier Laminate)
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove contaminated clothing and wash clothing before reuse.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. This product may pose an aspiration pneumonia hazard. Contains petroleum distillates

9767

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas. Runoff may be hazardous to aquatic organisms in neighboring areas.

Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Protective eyewear
- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (such as Nitrile, Butyl, Neoprene, and/or Barrier Laminate)
- Chemical-resistant headgear for overhead exposure

Non-Agricultural Use Requirements:

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal:

Do not contaminate water, food or feed by storage or disposal

Storage: Keep from Freezing. Store above 32°F

Pesticide Disposal: Pesticide Wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Plastic Containers: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities

Steps to be Taken In Case Material Is Released or Spilled: Ventilate area. Avoid breathing vapors. Use MSHA/NIOSH self-contained breathing apparatus or airline respirator for large spills in confined areas. Dike the spill with inert material (sand, earth, etc.) and transfer the liquid or solid diking material to separate containers for recovery or disposal. Remove contaminated clothing promptly and wash exposed skin areas with soap and water. Wash clothing before reuse. Keep spill out of all sewers and bodies of water.

CHEMIGATION

Do not apply this product through any irrigation system unless the instructions for chemigation are followed. **If application by chemigation is not specifically listed under the specific crop use instructions, GOAL 2XL may not be applied to that crop through an irrigation system.**

Apply this product only through sprinkler (solid set, portable lateral, or low-volume (microsprinkler)), drip (trickle), or flood (basin) irrigation systems. Refer to use directions for specific crops for instructions as to which type of irrigation system may be used. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER CHEMIGATION (FOLIAR SPRAY USES)

To apply a pesticide using sprinkler chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

For sprinkler irrigation, sufficient water should be applied at the beginning of the irrigation period to insure

uniform wetting of the plant and/or soil surfaces. Meter GOAL 2XL at a continuous uniform rate during the middle 1/3 of the irrigation period to allow for uniform distribution to the vegetation and/or soil surface. Continue irrigation during the final 1/3 of the irrigation period to insure proper flushing of the irrigation system. During sprinkler irrigation, sufficient water should be applied to insure water penetration to a depth of two inches.

FLOOD (BASIN) CHEMIGATION (SOIL DRENCH USES)

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

GOAL 2XL should be continuously metered into the water during the entire irrigation period. Agitation in the pesticide supply tank is suggested. Best weed control results from Goal 2XL applied through flood (basin) irrigation systems are obtained when a uniform distribution and flow of irrigation water is maintained over level land.

DRIP (TRICKLE) CHEMIGATION (SOIL DRENCH USES)

To apply a pesticide using drip (trickle) chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Meter GOAL 2XL at a continuous uniform rate during the middle 1/3 of the irrigation period to allow for uniform distribution to the soil surface. For best results, Goal 2XL should be uniformly positioned across the wetted area to help reduce the "ring effect" of weed escapes, as other products begin to break down around the emitter. Continue irrigation during the final 1/3 of the irrigation period to insure proper flushing of the irrigation system.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

If the chemigation system is connected to a public water supply, the following conditions must also be met:

- Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

GENERAL USE INFORMATION

Unless otherwise directed by registered supplemental labels, follow the Directions for Use in each crop group section.

CULTURAL CONSIDERATIONS

In order for Goal 2XL to provide maximum preemergence activity:

Prior to application, the bed or soil surface should be smooth and free of crop and weed trash (decaying leaves, clippings, dead weeds, etc.). Leaves and trash may be removed by blowing the area to be treated or by thoroughly mixing the trash into the soil through cultivation prior to herbicide application.

After application, at least one-quarter inch (1/4 inch) of irrigation or rainfall should occur within 3 or 4 weeks after application. The best results from Goal 2XL are from applications to established beds or soil surfaces that are left undisturbed during the time period for which weed control is desired. Cultural practices that result in redistribution or disturbance of the soil surface after treatment will decrease the herbicidal effectiveness of GOAL 2XL. Cutting water furrows or cultivations that mix untreated soil into treated areas will also reduce the effectiveness of the treatment.

MIXING DIRECTIONS

Fill the spray tank at least one-third full of clean water. With the pump and agitator running, add the recommended amount of herbicides to the spray tank. The order of addition to the spray tank should be wettable powders first, flowables second and liquids last. Complete filling of the spray tank with water. For all applications of Goal 2XL (except onions) where postemergence weed control is desired, add 2 to 4 pints of Latron AG-98* nonionic surfactant (or comparable 80% active nonionic surfactant cleared for application to growing crops) per each 100 gallons of spray. The addition of 4 pints of Latron AG-98 per 100 gallons of spray is recommended to enhance postemergence activity when hard water (greater than 600 ppm) is used as carrier. Maintain agitation until spraying is completed.

Spray equipment should be calibrated carefully before each use.

Dosages listed on this label are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

<u>Band Width (in inches)</u>	X	Rate per	=	Amount Needed per Acre
Row Width (in inches)		Broadcast Acre		for Banded Application

ARTICHOKES (GLOBE)

POST-DIRECTED SPRAY

GENERAL INFORMATION

GOAL 2XL is an effective herbicide for postemergence and preemergence control of certain broadleaf weeds in artichokes. Goal 2XL should be directed towards the winter ditch, levees or flat rows between the artichoke rows. Artichoke fronds receiving accidental spray or drift will be injured. Over-the-top applications may exhibit severe injury to the foliage and flower bud and are not recommended.

DOSAGE

Goal 2XL is recommended as a post-directed application at 4 to 8 pints (1.0 to 2.0 lb active) per acre. Optimum control is achieved when two applications of Goal 2XL are applied at 4 pints (1.0 lb. active) per acre. The initial application should be made to susceptible weed seedlings (up to 8-leaf stage). It is recommended that a second application be made 8 to 10 weeks later. Good results may be achieved when a single application of 8 pints (2.0 lb active) of Goal 2XL is applied to susceptible weed seedlings (up to 8-leaf stage). Do not apply more than 8 pints (2.0 lb active) of Goal 2XL per treated acre per season as a result of a single application or multiple applications. Do not apply within 5 days of harvest.

WEEDS CONTROLLED POSTEMERGENCE

CHEESEWEED (MALVA)	OXALIS (BERMUDA BUTTERCUP)
GROUNDSEL, COMMON	SHEPHERDSPURSE
MUSTARD, COMMON YELLOW	SOWTHISTLE, ANNUAL
NETTLE, BURNING	

WEEDS CONTROLLED PREEMERGENCE

CHEESEWEED (MALVA)	† OXALIS (BERMUDA BUTTERCUP)
GROUNDSEL, COMMON	SHEPHERDSPURSE
LAMBSQUARTERS, COMMON	SOWTHISTLE, ANNUAL
MUSTARD, COMMON YELLOW	

† Suppression

TIMING AND METHOD OF APPLICATION

Treatments should be made after completion of the ditching operation.

Goal 2XL should be applied in a minimum of 40 gallons of water per acre depending upon density of emerged weeds. Spray volume should be increased as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use. Spray should be directed towards the winter ditch, levees or flat rows between the artichoke rows. **ARTICHOKE FRONDS RECEIVING ACCIDENTAL SPRAY OR DRIFT WILL BE INJURED.**

ARTICHOKES (GLOBE)

SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 8 pints (2.0 lb active) of Goal 2XL per treated acre per season as a result of a single application or multiple applications.
- Do not apply Goal 2XL within 5 days of harvest.
- Avoid direct spray or drift contact of Goal 2XL with artichoke flowers or buds as severe injury may result.

- Do not apply Goal 2XL to artichoke plantings within 60 days after cutting back or transplanting.

BROCCOLI/CABBAGE/CAULIFLOWER

PRE-TRANSPLANT (PREPLANT) APPLICATION FOR PREEMERGENCE BROADLEAF WEED CONTROL

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for preemergence control of certain annual broadleaf weeds. Applications must be made after completion of soil preparation but *prior* to transplanting of broccoli, cabbage or cauliflower plants. Transplanting should be completed with minimal soil disturbance. Treated soil surfaces should be left undisturbed after transplanting to obtain greatest benefit of Goal 2XL on susceptible annual broadleaf weeds during the time period for which weed control is desired. However, timely cultivations after weed emergence will assist in weed control.

Pre-transplant applications of Goal 2XL in broccoli, cabbage and cauliflower may result in a temporary initial crop response (leaf cupping or crinkling). Crop response may be enhanced if crop leaves come in direct contact with treated soil. Crops rapidly outgrow this condition and develop normally.

Severe crop response may result from the use of transplants that are under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides or storage conditions. The use of young (less than 5 weeks old), extremely succulent transplants grown in containers, less than 1 inch square, may increase the severity of crop injury. Hardening off, increasing the age of transplants or increasing the size of the rooting container will lessen the possibility and/or severity of crop injury.

DOSAGE

Goal 2XL is recommended for use at 1 to 2 pints (0.25 to 0.5 lb active) per broadcast acre. The lower rate (1-pint per acre) is recommended for preemergence weed control on coarse texture soils with less than 1% organic matter. The 2 pint per acre rate is recommended for preemergence weed control on medium to fine texture soils or soils containing greater than 1% organic matter.

Goal 2XL will assist in early season annual grass control. However, Goal 2XL must not be a basic portion of the grass herbicide program. A planned herbicide program for preemergence or postemergence grass control is recommended. Research has shown that severe crop injury can occur if Goal 2XL is applied to a field that has had an acetanilide herbicide (Dual herbicide, Lasso herbicide, Ramrod herbicide) application during the current growing season, therefore, it is not recommended.

WEEDS CONTROLLED †

CARPETWEED

PURSLANE, COMMON

PIGWEEED, REDROOT

SMARTWEED, PENNSYLVANIA

Goal 2XL at the rate of 1 to 2 pints per acre may provide partial control or suppression of galinsoga, common lambsquarters and wild mustard.

† Applications of Goal 2XL to muck soils may result in partial control or suppression of the weeds listed.

METHOD OF APPLICATION

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 20 gallons of water per acre. Use conventional ground spray equipment with flat fan nozzles at 20 to 40 psi. Do not exceed 40 psi. Accurately calibrate spray equipment prior to each use.

Thoroughly flush the spray equipment (tank, hose, pump, boom) with water before and after each use.

Residual Goal 2XL remaining in spray equipment may damage other crops.

AVOID DRIFT TO ALL OTHER CROPS AND NON-TARGET AREAS. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT. GOAL 2XL IS PHYTOTOXIC TO PLANT FOLIAGE.

**BROCCOLI/CABBAGE/CAULIFLOWER
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per treated acre per season.
- Do not apply Goal 2XL preemergence to direct-seeded broccoli, cabbage or cauliflower.
- Do not apply Goal 2XL post-transplant or postemergence (over the top) to broccoli, cabbage or cauliflower.
- For field use only. Do not apply Goal 2XL in an enclosed greenhouse structure as injury to plant foliage may result.

CACAO

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence herbicide when used alone for the control of certain annual broadleaf weeds in bearing and nonbearing cacao plantings. Preemergence control is most effective when spray is applied to clean, weed-free soil surfaces. Treated berms or soil surfaces should not be disked or disturbed in any manner as the herbicidal effectiveness of Goal 2XL may be decreased. Seedling weeds are controlled as they come in contact with soil-applied herbicide during emergence.

GOAL 2XL USED ALONE

DOSAGE

Goal 2XL is recommended for preemergence and postemergence control of susceptible weeds at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre when directed to the orchard floor beneath cacao plants, or at a dosage of up to 4 pints per acre as a pre-transplant application.

For directed spray applications, cacao transplants must be healthy and of suitable size for field transplanting. Avoid spray contact with cacao foliage as injury may result.

Dosages listed are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

WEEDS CONTROLLED POSTEMERGENCE

Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Applications to weeds beyond the four-leaf stage may result in partial control.

PURSLANE, COMMON SPURGE, GARDEN

WEEDS CONTROLLED PREEMERGENCE

Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre.

AGERATUM PURSLANE, COMMON
BUTTONWEED SPURGE, GARDEN
CROTALARIA

TIMING AND METHOD OF APPLICATION

DO NOT APPLY PREPLANT OR PREEMERGENCE TO DIRECT-SEEDED CACAO. TREATMENTS CAN BE MADE TO ESTABLISHED CACAO OR AS A PRE-TRANSPLANT OR RECENTLY TRANSPLANTED CACAO.

Treatments should only be applied to healthy cacao stock (as determined by standard commercial growing practices). Care must be taken to prevent direct spray contact with foliage. Cacao foliage receiving accidental spray or drift may be injured.

As a preemergence or postemergence treatment to weeds, apply in a minimum of 15 gallons of water per acre. Use higher volumes to assure adequate coverage in high densities of emerged weeds or heavy trash. Goal 2XL should be directed to the soil and the base of the tree. Use of a low pressure sprayer equipped with a breakaway boom and a flat fan or off-center (OC) nozzles is recommended. Spray shields are suggested for use in young trees. Spray equipment should be calibrated carefully before each use.

CACAO - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe the GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in a single application or 24 pints (6.0 lb active) per broadcast acre per year.
- Do not apply Goal 2XL within one (1) day of harvest.
- Direct spray toward the base of the trees. Avoid spray contact with foliage.
- Do not apply preplant or preemergence to direct-seeded cacao.

CITRUS (NONBEARING)

CALAMONDIN, CHIRONJA, CITRUS CITRON, GRAPEFRUIT, KUMQUAT, LEMON, LIME, MANDARIN, PUMMELO, SATSUMA MANDARIN, SOUR ORANGE, SWEET ORANGE, TANGELO, TANGERINE, TANGOR
FOR USE ONLY IN PERMANENTLY ESTABLISHED GROVES IN ARIZONA, CALIFORNIA, FLORIDA, LOUISIANA AND TEXAS

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence and/or postemergence herbicide when used alone or in recommended tank mix combinations, for the control of certain annual broadleaf weeds in nonbearing citrus plantings. Goal 2XL may be applied to newly planted trees or to young trees that will not bear fruit within one year. The most effective postemergence weed control is achieved when Goal 2XL is applied to seedling weeds at the recommended growth stage. For postemergence control of certain grassy and broadleaf weeds, a tank mix of Goal 2XL with paraquat (Gramoxone herbicide) or glyphosate (Glyphomax* herbicide) can be used. For residual grass control in citrus, a tank mixture of Goal 2XL with Devrinol herbicide, simazine, Solicam herbicide or Surflan* herbicide can be used. Contact herbicides such as paraquat (Gramoxone) or glyphosate (Glyphomax) may also be added to the tank mixture. Check individual product labels to determine suitability and use rates for various crops.

GOAL 2XL USED ALONE

GEOGRAPHIC USE DIRECTIONS ARIZONA AND CALIFORNIA

DOSAGE

Goal 2XL is recommended for postemergence control at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre. For preemergence control of susceptible weeds, use 8 pints (2.0 lb active) per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE (weeds up to 4 inches high) - Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Applications to weeds beyond this 4-inch stage may result in partial control.

CHEESEWEED (MALVA)	MINERSLETTUCE
FIDDLENECK, COAST	NETTLE, BURNING
† FILAREE, BROADLEAF	PIGWEEED, REDROOT
† FILAREE, REDSTEM	REDMAIDS
† FILAREE, WHITESTEM	SHEPHERDSPURSE
GROUNDSEL, COMMON	SOWTHISTLE, ANNUAL
HENBIT	

† Goal 2XL at the 8-pint rate (2.0 lb active) will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

WEEDS CONTROLLED PREEMERGENCE - Apply 5 to 8 pints (1.25 to 2.0 lb active) of Goal 2XL per broadcast acre.

BURCLOVER	LETTUCE, PRICKLY
CHEESEWEED (MALVA)	PIGWEEED, REDROOT
FIDDLENECK, COAST	PURSLANE, COMMON
FILAREE, BROADLEAF	REDMAIDS
FILAREE, REDSTEM	ROCKET, LONDON
FILAREE, WHITESTEM	SHEPHERDSPURSE
GROUNDSEL, COMMON	SOWTHISTLE, ANNUAL
HENBIT	SPURGE, PROSTRATE
KNOTWEED, PROSTRATE	SPURGE, SPOTTED
LAMBSQUARTERS, COMMON	

FLORIDA, LOUISIANA AND TEXAS

DOSAGE

Goal 2XL is recommended for postemergence control at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre. For preemergence control of susceptible weeds, Goal 2XL is recommended at 8 pints (2.0 lb active) per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE - Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. The lower rate is recommended for the control of susceptible seedling weeds in the early postemergence stage, up to the 4-leaf stage. The higher rate (2.0 lb active) should be used for weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in partial control.

BALSAMAPPLE	PEPPERWEED, VIRGINIA
† CUDWEED, NARROWLEAF	PIGWEEED, REDROOT
†† EVENINGPRIMROSE, CUTLEAF	POINSETTIA, WILD
GROUNDCHERRY, CUTLEAF	PURSLANE, COMMON
GROUNDCHERRY, WRIGHT	PUSLEY, FLORIDA
LAMBSQUARTERS, COMMON	SIDA, PRICKLY (TEAWEED)
MORNINGGLORY, ANNUAL	SMARTWEED, PENNSYLVANIA
NIGHTSHADE, AMERICAN BLACK	SOWTHISTLE, ANNUAL
NIGHTSHADE, BLACK	

WEEDS CONTROLLED PREEMERGENCE - Apply 8 pints (2.0 lb active) of Goal 2XL per broadcast acre.

CUDWEED, NARROWLEAF	POINSETTIA, WILD
†† EVENINGPRIMROSE, CUTLEAF	PUSLEY, FLORIDA
GROUNDCHERRY, CUTLEAF	SIDA, PRICKLY (TEAWEED)

LAMBSQUARTERS, COMMON	SMARTWEED, PENNSYLVANIA
NIGHTSHADE, AMERICAN BLACK	SOWTHISTLE, ANNUAL
NIGHTSHADE, BLACK	SPURGE, PROSTRATE
PEPPERWEED, VIRGINIA	SPURGE, SPOTTED
PIGWEEED, REDROOT	

† Maximum 0.5-inch diameter.

†† Highest rate and/or multiple applications may be required for acceptable control. Do not apply more than 16 pints (4.0 lb active) per broadcast acre during any 12-month period as a result of multiple applications.

ALL STATES - ARIZONA, CALIFORNIA, FLORIDA, LOUISIANA AND TEXAS

TIMING AND METHOD OF APPLICATION

Goal 2XL should be directed to the soil and the base of trees. Avoid direct spray contact on the citrus foliage. Use a low-pressure sprayer equipped with a breakaway boom and flat fan nozzles. An off-center (OC) nozzle positioned at the end of the boom may be desired.

SPRAY VOLUME:

Weed Stage	Gallons of Water Per Acre
Preemergence	40 or more
Postemergence up to 4-inch or 4-leaf stage	40 or more
Exceeding 4-inch or 4-leaf stage	100 or more

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

DOSAGE

For preemergence control of susceptible grassy and broadleaf weeds in citrus plantings, a tank mixture of Goal 2XL with Devrinol, simazine, Solicam or Surflan can be applied. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

For postemergence control of susceptible grassy and broadleaf weeds, a tank mixture of paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL or combinations of Goal 2XL plus Devrinol, simazine, Solicam or Surflan can be used. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

WEEDS CONTROLLED

In addition to the weeds controlled by Goal 2XL used alone, control of susceptible weeds listed on the respective labels for the following products is also obtained:

Devrinol	simazine †
paraquat (Gramoxone)	Solicam
glyphosate (Glyphomax)	Surflan

† In addition, provides preemergence control of horseweed (marestalk).

**CITRUS (NONBEARING)
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe the GENERAL USE RESTRICTIONS listed at the end of this label.

- Apply Goal 2XL only to nonbearing citrus trees.
- Do not apply more than 8 pints of Goal 2XL (2.0 lb active) per broadcast acre in a single application or more than 16 pints (4.0 lb active) per broadcast acre during any 12-month period as a result of multiple applications.
- Goal 2XL or any of the combinations recommended on this label should only be applied to healthy growing trees.
- Do not apply during periods of new foliage growth. Applications should be made after foliage has fully expanded and hardened off.
- Direct spray toward the base of trees. Avoid direct sprays contact on the citrus foliage.

COFFEE

BEARING AND NONBEARING COFFEE IN HAWAII

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence herbicide when used alone for the control of certain annual broadleaf weeds in bearing and nonbearing coffee plantings.

For postemergence control of certain grassy and broadleaf weeds, a tank mixture of either paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL can be applied to seedling weeds.

Check individual product labels to determine suitability and use rates for crop.

GOAL 2XL USED ALONE

DOSAGE

For preemergence control of susceptible weeds, Goal 2XL is recommended at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre as a preemergence application directed to the orchard floor beneath coffee plants, or at a dosage of up to 4 pints per broadcast acre as a pre-transplant application.

For directed spray applications, coffee transplants must be healthy and of suitable size for field transplanting. Avoid spray contact with coffee foliage as injury may result.

Goal 2XL may be applied postemergence (over the top) to dormant coffee transplants. Applications must only be made prior to bud break to avoid possible phytotoxicity to the coffee foliage. Over-the-top applications made after buds start to swell may result in injury to the coffee plant and are not recommended.

Dosages listed on this label are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

WEEDS CONTROLLED POSTEMERGENCE

Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Applications to weeds beyond the four leaf stage may result in partial control.

PURSLANE, COMMON

SPURGE, GARDEN

WEEDS CONTROLLED PREEMERGENCE

Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre.

AGERATUM

PURSLANE, COMMON

BUTTONWEED

SPURGE, GARDEN

CROTALARIA**TIMING AND METHOD OF APPLICATION**

DO NOT APPLY PREPLANT OR PREEMERGENCE TO DIRECT SEEDED COFFEE.

Treatments should only be applied to healthy coffee stock (as determined by standard commercial growing practices). Care must be taken to prevent direct spray contact with foliage. Coffee foliage receiving accidental spray or drift may be injured.

As a preemergence or postemergence treatment to weeds, apply in a minimum of 30 gallons of water per acre. Use higher volumes to ensure adequate coverage in high densities of emerged weeds or heavy trash. Goal 2XL should be directed to the soil and the base of the tree. Use of a low-pressure sprayer equipped with a breakaway boom and flat fan or off-center (OC) nozzles is recommended. Spray equipment should be calibrated carefully before each use.

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

For postemergence control of susceptible grassy and broadleaf weeds in coffee plantings, a tank mixture of Goal 2XL with either glyphosate (Glyphomax) or paraquat (Gramoxone) may be applied as a directed spray. Apply at recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

WEEDS CONTROLLED POSTEMERGENCE

In addition to the weeds controlled by Goal 2XL used alone, control of susceptible weeds listed on the respective labels for the following products is also obtained.

paraquat (Gramoxone)

glyphosate (Glyphomax)

COFFEE**SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply preplant or preemergence to direct-seeded coffee.
- Direct spray toward the base of the trees. Avoid spray contact with foliage.
- Goal 2XL may be applied as a postemergence (over the top) application to dormant transplants. Do not apply over the top to coffee transplants after buds start to swell.
- Goal 2XL or any of the combinations recommended on this label should be applied to only healthy growing trees/transplants under standard commercial growing practices.
- Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in a single application or 24 pints (6.0 lb active) per broadcast acre per year.
- Do not apply Goal 2XL within one (1) day of harvesting.
- Applications of Goal 2XL during periods of rapid new foliage growth may cause injury.

**CONIFER SEEDBEDS, TRANSPLANTS, CONTAINER STOCK and
SELECTED FIELD GROWN DECIDUOUS TREES**

GOAL 2XL is effective as a preemergence and/or postemergence herbicide for the control of certain annual grassy and broadleaf weeds in conifer seedbeds, transplant and container stock, and in selected field-grown deciduous trees. Preemergence weed control is most effective when spray is applied to clean, weed-free soil surfaces. Treated soil surfaces should not be disturbed as the herbicidal effectiveness of GOAL 2XL may be decreased. Seedling weeds are controlled during emergence as they come in contact

with the soil-applied herbicide. The most effective postemergence weed control is achieved when Goal 2XL is applied to seedling weeds less than 4 inches in height.

Occasionally after the use of Goal 2XL, a spotting, crinkling, or flecking may appear on leaves of conifer and deciduous species. Leaves that receive direct or indirect (drift) spray contact may be injured. The conifer and deciduous species typically outgrow this condition rapidly and develop normally.

IMPORTANT. When applied as directed, the conifer and selected deciduous species listed on this label have shown tolerance to GOAL 2XL. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of listed species on this label or under all possible growing conditions. The user should exercise reasonable judgment and caution with this product. Until familiar with results under user growing conditions, limit application of this product to a few plants in a small treated area to determine plant tolerance and extent of injury if such occurs, prior to initiating large-scale applications.

WEEDS CONTROLLED

When Goal 2XL is applied preemergence or postemergence at recommended dosages and weed stages, the following grasses and broadleaf weeds are controlled.

- | | |
|-------------------------|-------------------------|
| † BARNYARDGRASS | MUSTARD, BLUE |
| BEDSTRAW, CATCHWEED | MUSTARD, TUMBLE |
| BITTERCRESS, LESSER | MUSTARD, WILD |
| † BLUEGRASS, ANNUAL | NETTLE, BURNING |
| BUCKWHEAT, WILD | NIGHTSHADE, BLACK |
| BURCLOVER | NIGHTSHADE, HAIRY |
| CARPETWEED | OATS, WILD |
| † CLOVER, RED | ORACH, RED |
| † CLOVER, WHITE | PEPPERWEED, |
| | YELLOWFLOWER |
| COCKLEBUR, COMMON | PIGWEEED, PROSTRATE |
| † CRABGRASS, LARGE | PIGWEEED, REDROOT |
| † FIDDLENECK, COAST | PIMPERNEL, SCARLET |
| FILAREE, BROADLEAF | PURSLANE, COMMON |
| FILAREE, REDSTEM | REDMAIDS |
| FIREWEED (FROM SEED) | ROCKET, LONDON |
| FLIXWEED | SANDSPURRY, RED |
| † FOXTAIL, GIANT | † SHEPHERDSPURSE |
| † GOOSEGRASS | SIDA, PRICKLY |
| GROUNDCHERRY, CUTLEAF | SMARTWEED, PENNSYLVANIA |
| GROUNDCHERRY, WRIGHT | SORREL, RED (FROM SEED) |
| GROUNDSEL, COMMON | SOWTHISTLE, ANNUAL |
| HENBIT | SPEEDWELL, BIRDSEYE |
| JIMSONWEED | †† SPURGE, PROSTRATE |
| KNOTWEED, PROSTRATE | †† SPURGE, SPOTTED |
| LADYSTHUMB | SPURRY, CORN |
| LAMBSQUARTERS, COMMON | TANSYMUSTARD |
| LETTUCE, PRICKLY | †† THISTLE, BULL |
| MALLOW, LITTLE | THISTLE, RUSSIAN |
| MAYWEED | VELVETLEAF |
| MINERSLETTUCE | WITCHGRASS |
| † MORNINGGLORY, IVYLEAF | †† WOODSORREL, YELLOW |
| † MORNINGGLORY, TALL | |

† Highest rate and/or multiple applications may be required for acceptable control.

†† Preemergence control only.

Goal 2XL is most effective when applied preemergence to annual grasses. Postemergence applications should be made to seedling grasses not exceeding the 2-leaf stage. The addition of 0.25% (2 pints/100 gallons of spray solution) of Latron AG-98 or comparable 80% active nonionic surfactant, cleared for application on growing crops, enhances the Goal 2XL activity on emerged weeds.

When determining an appropriate use rate where a range of rates are provided, use higher rates where heavy weed pressure is anticipated, or where medium and fine soil textures exist and high organic matter soils are present.

CONIFER SEEDBEDS

To assist in the establishment of conifer seedbeds, Goal 2XL can be applied as a preemergence application following seeding. Postemergence applications should be delayed until a minimum of 5 weeks after emergence of the conifer seedlings. During periods of cool, cloudy weather, make certain that seedlings have hardened off prior to spraying.

Conifers are tolerant to preemergence and postemergence applications of Goal 2XL. Goal 2XL will provide both postemergence and residual preemergence control of many broadleaf weeds and annual grass species.

CONIFER SPECIES

Goal 2XL may be applied to conifer seedbeds of species:

DOUGLAS FIR	<i>Pseudotsuga menziesii</i>
FIR	
FRASER	<i>Abies fraseri</i>
GRAND	<i>Abies grandis</i>
NOBLE	<i>Abies procera</i>
HEMLOCK	
EASTERN HEMLOCK	<i>Tsuga canadensis</i>
PINE	
AUSTRIAN	<i>Pinus nigra</i>
EASTERN WHITE	<i>Pinus strobus</i>
HIMALAYAN	<i>Pinus wallichiana</i>
JACK	<i>Pinus banksiana</i>
LOBLOLLY	<i>Pinus taeda</i>
LOGEPOLE	<i>Pinus contorta</i>
LONGLEAF	<i>Pinus palustris</i>
MONTEREY	<i>Pinus radiata</i>
MUGHO	<i>Pinus mugo</i>
PONDEROSA	<i>Pinus ponderosa</i>
SCOTCH	<i>Pinus sylvestris</i>
SHORTLEAF	<i>Pinus echinata</i>
SLASH	<i>Pinus elliottii</i>
VIRGINIA	<i>Pinus virginiana</i>

SPRUCE

BLUE
DWARF ALBERTA
NORWAY
SITKA

Picea pungens
Picea glauca Conica
Picea abies
Picea sitchensis

PREEMERGENCE DOSAGE

Apply 1 to 4 pints (0.25 lb. to 1.0 lb. active) of Goal 2XL per broadcast acre as a preemergence application prior to conifer emergence. Where grassy weeds are present, a rate of 2 to 4 pints (0.5 to 1.0 lb. active) of Goal 2XL per broadcast acre is recommended. In known areas of high weed competition, 4 pints (1.0 lb. active) of Goal 2XL per broadcast acre are recommended.

TIMING AND METHOD OF APPLICATION

Goal 2XL should be thoroughly mixed with clean water at recommended concentration and applied at 20 to 40 psi in a minimum of 20 gallons of water per treated acre. Broadcast to beds and irrigate prior to weed emergence with 1/2 to 3/4 inch of sprinkler irrigation.

POSTEMERGENCE DOSAGE

Apply 1 to 2 pints (0.25 lb. to 0.5 lb. active) of Goal 2XL per broadcast acre with each postemergence application. Depending of subsequent weed flushes, multiple applications may be necessary to achieve season-long weed control.

TIMING AND METHOD OF APPLICATION

Postemergence applications should be delayed until a minimum of 5 weeks after emergence of conifer seedlings. During periods of cool, cloudy weather, make certain that seedlings have hardened off prior to spraying. Application should be made to seedling weeds (less than 4 inches in height). Goal 2XL should be thoroughly mixed with clean water at recommended concentration and applied as a broadcast application at 20 to 40 psi in a minimum of 20 gallons of water per treated acre.

**CONIFER TRANSPLANTS AND CONTAINER STOCK
(INCLUDES 2-0 SEEDLING AND CHRISTMAS TREE PLANTINGS)**

Many container-grown conifers and conifer transplants are tolerant to preemergence and postemergence applications of Goal 2XL. Applied postemergence, Goal 2XL will provide both postemergence and preemergence control of many broadleaf weeds and grasses listed in the "Weeds Controlled" Section above. Postemergence applications should be applied before bud break or after foliage has had an opportunity to harden off. Conifers may be transplanted from seedbeds and sprayed directly providing bud break has not occurred.

The following conifer species in addition to species listed under the CONIFER SEEDBED section have been shown to be tolerant to Goal 2XL.

ARBORVITAE

Thuja occidentalis
Thuja orientalis

RED CEDAR

Juniperus virginiana

JUNIPER

Juniperus chinensis
Juniperus horizontalis
Juniperus procumbens
Juniperus sabina
Juniperus scopulorum

WESTERN HEMLOCK

Tsuga heterophylla

YEW

Taxus species

DOSAGE

For preemergence or postemergence weed control apply 4 to 8 pints (1.0 lb. to 2.0 lb active) of Goal 2XL per broadcast acre.

TIMING AND METHOD OF APPLICATION

For optimum weed control, preemergence applications should be made immediately after transplanting seedlings or to weed-free container stock. Postemergence applications should be made to weeds less than 4 inches in height. Two applications may be necessary, in fall-transplanted conifer fields, for season-long weed control. The addition of 0.25% (2 pints/100 gals. of spray solution) of Latron AG-98 or comparable 80% active nonionic surfactant, cleared for application to growing crops, enhances Goal 2XL activity on emerged weeds. Goal 2XL must be applied only to conifer transplants prior to bud break or after foliage has had an opportunity to harden off. Thoroughly mix with clean water at recommended concentration and apply at 20 to 40 psi in a minimum of 20 gallons of water per treated acre. Spray over the top of transplants. Heavy rainfall immediately following application to emerged weeds may reduce effectiveness.

TANK MIXTURES FOR SELECTED FIELD GROWN CONIFERS

In addition to the weeds controlled by Goal 2XL used alone, tank mixes with other preemergence or postemergence herbicides registered for this use, may provide a broader spectrum of weed control. Goal 2XL may be tank mixed with products containing the following active ingredients registered for use in conifer plantings:

glyphosate
 napropamide
 oryzalin
 pendimethalin
 prodiamine
 pronamide
 sethoxydim

Determine the additional weed species to be controlled and based on label claims, select the product(s), which would give effective control of the targeted weed(s). When using tank mixes of two or more products, use conditions must be in accordance with the more (most) restrictive of the label limitations and precautions of the mixing partners.

IMPORTANT: Read and follow container labels of tank-mix partners and use as directed by labeling. Follow the most restrictive labeling.

CONIFER - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 8 pints (2.0 lb active) of this product per broadcast acre per year.
- **NOT FOR CONIFER RELEASE IN FOREST MANAGEMENT PROGRAMS OR FOR FOREST REGENERATION APPLICATIONS.**
- Do not apply Goal 2XL in an enclosed greenhouse structure as injury to plant foliage may result.
- Do not store or transport treated container stock in an enclosed structure until completion of 4 irrigations (minimum 21 days) as injury to non-labeled plants may occur.
- Always apply Goal 2XL only to healthy conifer stock. Do not apply Goal 2XL to conifers that are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result.
- Do not graze or feed livestock forage cut from areas treated with Goal 2XL.

SELECTED FIELD-GROWN DECIDUOUS TREES

Many field-grown deciduous trees are tolerant to applications of Goal 2XL directed to the soil and base of the plant. Goal 2XL will provide both preemergence and postemergence control of many broadleaf weeds and grasses.

DECIDUOUS TREE SPECIES	
†† Almond	<i>Prunus spp.</i>
†† Apple	<i>Malus X domestica</i>
†† Apricot	<i>Prunus spp.</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>
Ash, White	<i>Fraxinus americana</i>
Birch, River	<i>Betula nigra</i>
†† Cherry	<i>Prunus spp.</i>
†† Chestnut	<i>Castanea spp.</i>
†† Crabapple	<i>Malus spp.</i>
Dogwood	<i>Cornus florida</i>
Eucalyptus	<i>Eucalyptus viminalis</i> <i>Eucalyptus pulverulenta</i> <i>Eucalyptus camaldulensis</i>
†† Filbert	<i>Corylus spp.</i>
Lilac	<i>Syringa vulgaris</i>
Locust, Black	<i>Robinia pseudoacacia</i>
† Maple, Black	<i>Acer nigrum</i>
† Maple, Red	<i>Acer rubrum</i>
† Maple, Sugar	<i>Acer saccharum</i>
Myrtle, Crepe	<i>Lagerstroemia indica</i>
†† Nectarine	<i>Prunus spp.</i>
†† Nut, Hickory	<i>Carya sp.</i>
†† Nut, Macadamia	<i>Macadamia ternifolia</i>
Oak, Chestnut	<i>Quercus prinus.</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus. rubra</i>
Oak, Water	<i>Quercus nigra</i>
Oak, Willow	<i>Quercus phellos</i>
Olive, Russian	<i>Elaeagnus angustifolia</i>
Poplar	<i>Populus spp.</i>
Poplar, Tulip	<i>Liriodendron tulipifera</i>
†† Peach	<i>Prunus persica</i>
†† Pear	<i>Pyrus spp.</i>
†† Pecan	<i>Carya spp.</i>
†† Pistachio	<i>Pistacia vera</i>
†† Plum	<i>Prunus spp.</i>
†† Prune	<i>Prunus spp.</i>
Redbud	<i>Cercis canadensis</i>
Sweetgum	<i>Liquidamber styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>
†† Walnut, Black	<i>Juglans nigra</i>

† Do not apply to maple trees used for production of maple sap or maple syrup.

†† Apply as directed to nonbearing trees. For bearing treefruit, nut and vine crops, refer to Treefruit/Nut/Vine section of this label for use directions.

DOSAGE

Apply 2 to 8 pints (0.5 lb. to 2.0 lb active) of Goal 2XL per acre as a spray onto to the soil area surrounding deciduous plants for preemergence or early postemergence weed control. This product may be applied as a single or split application.

DO NOT apply more than 8 pints of product per season.

For spot treatments, refer to the following table for dosage recommendations. Sprays must be uniform and applied to the soil on a spray-to-wet basis. When spraying to control weeds on a preemergence or postemergence basis, 1 gallon of spray mixture should cover 400 square feet (this is equivalent to applying Goal 2XL at a use rate of approximately one gallon per acre in a spray volume of 110 gallons per acre). It is recommended that an 80% active nonionic surfactant be added to the spray mixture at a rate of 1 tablespoon (0.5 fluid ounces) per gallon of spray when making postemergence applications.

Pounds Active/Acre	Pints GOAL 2XL (L.O.)/Acre	Fluid Ounces (milliliters) of GOAL 2XL in one gallon of spray mix to treat 400 Sq. Ft	Fluid Ounces (milliliters) of GOAL 2XL in one quart of spray mix to treat 100 Sq. Ft
2	8	1.2 (35)	0.3 (9)

TIMING

Goal 2XL can be applied after transplanting or to established deciduous trees. For optimum weed control, applications should be made prior to weed germination.

For maximum safety to deciduous species mentioned on this label, post-directed applications of GOAL 2XL should be made to the soil prior to bud swell in the spring or after trees have initiated dormancy in the fall. Care must be taken to avoid contact of spray drift or mist with foliage of or green bark of deciduous trees.

Goal 2XL may be phytotoxic to the foliage of non-target plants. Avoid making applications of this product under conditions that favor drift to non-target areas.

Note: Applications made after bud swell may result in injury to deciduous trees and are not recommended. If a non-dormant application is required due to weed competition, do not apply during periods of new foliage growth. Applications should be made after foliage has fully expanded and hardened off. Direct spray toward the soil at the base of the trees and use greater than 50 gallons of water per acre.. Splashing soil can carry Goal 2XL, which may injure the leaves of some deciduous trees.

METHOD OF APPLICATION

Goal 2XL should be directed to the soil. Avoid direct spray or drift onto foliage, flowers or green bark. Apply in 20 or more gallons of water per acre to provide uniform spray distribution and coverage to the soil surface. Use higher volumes to ensure adequate soil coverage in high densities of emerged weeds or heavy trash. Thorough spray coverage is essential to maximize the postemergence activity of GOAL 2XL. Use a low-pressure (20 to 40 psi.) sprayer. The use of spray shields that reduce exposure of foliage and bark to GOAL 2XL spray is suggested. Spray equipment should be calibrated carefully before each use.

TANK MIXTURES FOR SELECTED FIELD GROWN DECIDUOUS TREES

In addition to the weeds controlled by Goal 2XL used alone, tank mixes with other preemergence or postemergence herbicides registered for this use, may provide a broader spectrum of weed control. Goal 2XL may be tank mixed with products containing the following active ingredients registered for use in conifer plantings:

- glyphosate
- napropamide
- oryzalin
- pendimethalin
- prodiamine
- pronamide
- sethoxydim

Determine the additional weed species to be controlled and based on label claims, select the product(s) which would give effective control of the targeted weed(s). When using tank mixes of two or more products, use conditions must be in accordance with the more (most) restrictive of the label limitations and precautions of the mixing partners.

IMPORTANT: Read and follow container labels of tank-mix partners and use as directed by labeling. Follow the most restrictive labeling.

FIELD-GROWN DECIDUOUS TREES SPECIFIC USE RESTRICTIONS

- DO NOT apply more than 8 pints (2.0 lb active) of this product per broadcast acre per year.
- The use directions described here for field-grown deciduous trees do not apply for bearing treefruit, nut and vine crops. For selected bearing treefruit, nut and vine crops, refer to Treefruit/Nut/Vine section of this label use directions.
- Apply this product to the soil surface surrounding trees prior to bud swell or after trees have initiated dormancy in the fall. Although not recommended, if a non-dormant application is required, apply as a directed spray when foliage has fully expanded and hardened off. Do not apply during periods of new foliage growth.
- Avoid direct or indirect spray contact to foliage flowers and green bark.
- DO NOT apply this product when weather conditions favor drift. Avoid drift to non-target areas. Goal 2XL is phytotoxic to plant foliage.
- DO NOT apply Goal 2XL to trees that have been weakened or are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, wind injury, drought, flooding, previously applied pesticides, insects, or winter injury as severe injury may result.
- DO NOT graze or feed livestock forage cut from areas treated with Goal 2XL.

CORN

FOR USE ONLY AS DIRECTED SPRAY ON FIELD CORN IN CONJUNCTION WITH THE USDA WITCHWEED ERADICATION PROGRAM IN NORTH CAROLINA AND SOUTH CAROLINA

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for the control of witchweed (*Striga asiatica*), and works both preemergence and postemergence against witchweed.

DOSAGE

Use 2 to 3 pints of Goal 2XL (0.5 to 0.75 lb active) per acre for the first application. The 2 pint rate (0.5 lb active) per acre should be the standard use rate, with the 3 pint rate (0.75 lb active) per acre for isolated infestations. Repeat treatments should be made at rates of 1 to 2 pints (0.25 to 0.5 lb active) per acre.

Applications should be made in 10 to 30 gallons of water per acre. Use Latron AG-98 spreader in the spray mixture at the rate of 0.25% by water volume or 1 quart in 100 gallons of spray mix.

TIMING AND METHOD OF APPLICATION

Fields in the witchweed infested area selected for treatment with Goal 2XL should be examined during the early part of the growing season to determine uniformity of corn stand and grassy weed pressure. Weedy fields should be cultivated prior to the initial application so as to obtain the best possible soil coverage in the first spray application. Apply during May-August to emerged witchweed before bloom or as soon as possible after bloom appears, to avoid seed set. Corn should have a minimum height of 24 inches at the first application. After this application has been made, the fields should be inspected regularly for any breakthrough of the witchweed. If breakthrough occurs, then a second spray should be applied like the first. This application will be made postemergence to the witchweed, preferably before bloom or as soon as possible past the first appearance of witchweed bloom, to avoid seed set.

In all applications direct the Goal 2XL spray at the base of the corn plant and uniformly over the entire row surface. Do not spray over the top of the corn, as this may result in severe corn injury. Spray droplets contacting the lower leaves will cause necrotic spotting or streaking of sprayed tissue. Spray should contact only the lower 3 to 8 inches of the corn stalk and any leaves in this zone.

CORN - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 5 pints of Goal 2XL (1.25 lb active) per acre to a corn crop during the growing season.
- Do not apply any application within 60 days of harvest.
- Do not use corn plants from a treated field for green chop, ensilage, forage, or fodder.

COTTON

POST-DIRECTED SPRAY

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for use as a post-directed application for broadleaf weed control in cotton. Cotton leaves that are accidentally sprayed will exhibit necrotic spotting and may drop from the plant, therefore, care must be exercised to avoid spray contact with the cotton leaves. Crop response may be enhanced if applications are made when excessive soil moisture is present or if rainfall occurs following application. Cotton will outgrow this condition and continue to develop normally.

DOSAGE

Goal 2XL is recommended as a post-directed application at 1 to 2 pints (0.25 to 0.5 lb active) per acre.[†] Optimum control is achieved when 2 pints of Goal 2XL (0.5 lb active) per acre [†] are applied to weed seedlings not exceeding 4 true leaves. Effective control of succulent weed seedlings in the 2 to 3 leaf stage can usually be obtained when 1 pint of Goal 2XL (0.25 lb active) per acre [†] are applied. See MIXING DIRECTIONS for surfactant recommendation. Weeds should be in the seedling stage, young and actively growing. Do not count cotyledon leaves.

[†] Dosages listed are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

<u>Band Width</u> (in inches)	X	Rate per	=	Amount Needed per Acre
Row Width (in inches)		Broadcast Acre		for Banded Application

WEEDS CONTROLLED POSTEMERGENCE

When Goal 2XL is applied as a post-directed application at the recommended weed stage and dosage in cotton, the following weeds are controlled:

COCKLEBUR, COMMON		NIGHTSHADE, HAIRY
CROTON, TROPIC		PIGWEEED, REDROOT
GROUNDCHERRY, CUTLEAF	†	POINSETTIA, WILD
GROUNDCHERRY, WRIGHT		PURSLANE, COMMON
JIMSONWEED		SESBANIA, HEMP
LAMBSQUARTERS, COMMON	††	SICKLEPOD
MORNINGGLORY, ANNUAL (UP TO 6 LEAF)	†	SIDA, PRICKLY (TEAWEED)
NIGHTSHADE, AMERICAN BLACK		SMARTWEED, PENNSYLVANIA
NIGHTSHADE, BLACK		VELVETLEAF

† Multiple applications may be required for acceptable control.

†† Post-direct applications of Goal 2XL will kill or suppress seedlings not exceeding the one true leaf stage.

TIMING**SOUTHERN COTTON****ALABAMA, ARKANSAS, GEORGIA, LOUISIANA, MISSISSIPPI, MISSOURI, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS and VIRGINIA**

Cotton plant height must be a minimum 6 inches or greater. Application to cotton plants less than 6 inches tall may result in severe crop injury and is not recommended. In cotton 6 to 8 inches tall, Goal 2XL must be applied using rigid precision ground sprayer equipment. The use of spray shields is recommended to avoid spray contact with cotton foliage. Use branch lifters or shields if excessive spray contact on larger cotton plants (8 inches or greater) cannot be avoided by the directed spray.

WESTERN COTTON**ARIZONA AND CALIFORNIA**

Cotton plant height must be a minimum 6 inches or greater. Application to cotton plants less than 6 inches tall may result in severe crop injury and is not recommended. In cotton 6 to 8 inches tall, Goal 2XL must be applied using rigid precision ground sprayer equipment. The use of spray shields is recommended to avoid spray contact with cotton foliage. Use branch lifters or shields if excessive spray contact on larger cotton plants (8 inches or greater) cannot be avoided by the directed spray.

To obtain the maximum benefit of postemergence activity, encourage weed emergence by irrigating prior to spraying. Irrigate immediately following herbicide application to obtain greatest benefit of preemergence activity from Goal 2XL on nightshade and groundcherry species.

METHOD OF APPLICATION**SOUTHERN AND WESTERN COTTON**

Accurate, uniform placement of Goal 2XL spray is essential for effective weed control and to minimize cotton injury. As a directed postemergence application, Goal 2XL should be applied at 20 to 25 psi using 20 to 40 gallons of spray on a broadcast acre basis. Do not exceed 25 psi. Spray should be directed towards the base of the cotton plant. Cotton foliage receiving accidental spray or drift may be injured. Weeds should be in the seedling stage, young and actively growing.

Goal 2XL can be applied using a post-direct spray rig with only 2 flat fan nozzles per row, 1 nozzle on each side of the row. Additional care should be taken when adjusting sprayer prior to application. For best coverage, it is suggested to use 4 flat fan nozzles per row, 2 nozzles on each side of the row. The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and

downward. With either sprayer system, nozzles should be adjusted to cover the weed foliage with minimum contact to the cotton plant. **Do not use cone nozzles.**

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for tank mixtures, the most restrictive situations must apply.

DOSAGE

For postemergence control of susceptible grassy and broadleaf weeds in cotton, a tank mixture of Goal 2XL with either Bladex herbicide, Karmex herbicide (diuron) or MSMA can be applied as a post-directed application. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

COTTON - SOUTHERN AND WESTERN

SPECIFIC USE RESTRICTIONS

In addition to the following, also observe **GENERAL USE RESTRICTIONS** listed at the end of this label.

- **SOUTHERN COTTON:** Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL per season as a result of a single application or multiple applications. Do not apply within 90 days of harvest.
- **WESTERN COTTON:** Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per broadcast acre in a single application, or more than a total of 4 pints (1.0 lb. active) of Goal 2XL per broadcast acre per season as a result of multiple applications. Do not apply within 75 days of harvest.

COTTONWOOD

GENERAL INFORMATION

GOAL 2XL is an effective herbicide for postemergence and preemergence control of certain broadleaf weeds in cottonwood plantings. Goal 2XL may be applied postemergence or be post-directed to the base of the cottonwood tree. Applications must only be made prior to bud break to avoid possible phytotoxicity to the cottonwood foliage. Applications made after bud break may result in injury to the cottonwood plant and are not recommended.

DOSAGE

Apply 4 to 8 pints (1.0 to 2.0 lb active) of Goal 2XL per broadcast acre for preemergence and postemergence weed control. The addition of 1 quart of Latron AG-98 or a comparable 80% active nonionic surfactant per 100 gallons of spray mix will assist in spray coverage and wetting of weeds for postemergence control.

WEEDS CONTROLLED

When Goal 2XL is applied preemergence or postemergence to weed seedlings (not exceeding 6-leaf stage) at recommended dosages, the following broadleaf weeds are controlled:

GROUNDSEL, COMMON	MUSTARD, HEDGE
KNOTWEED, PROSTRATE	SHEPHERDSPURSE
LAMBSQUARTERS, COMMON	SMARTWEED, PENNSYLVANIA

TIMING AND METHOD OF APPLICATION

For optimum weed control, Goal 2XL should be applied prior to weed emergence. Preemergence applications should be made prior to or immediately after transplanting dormant cottonwood seedlings. Applications must be made prior to bud break of the cottonwood trees.

Goal 2XL should be applied in a minimum of 20 gallons of water per acre depending upon density of emerged weeds. Spray volume should be increased as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

COTTONWOOD

SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Goal 2XL should only be applied to dormant healthy cottonwood stock.
- Do not apply more than 8 pints (2.0 lb active) per treated acre per growing season as a result of single or multiple applications.

EUCALYPTUS

GENERAL INFORMATION

GOAL 2XL is an effective herbicide for postemergence and preemergence control of certain broadleaf weeds in permanently established eucalyptus (*E. viminalis*, *E. pulverulenta*, *E. camaldulensis*) plantings.

In new plantings, Goal 2XL should be applied immediately prior to or immediately following transplanting of dormant eucalyptus seedlings. In established plantings, Goal 2XL may be applied postemergence (over the top) or be post-directed to the base of the eucalyptus tree. Applications must only be made prior to bud break to avoid possible phytotoxicity to the eucalyptus foliage. Applications made after bud break may result in injury to the eucalyptus plant and are not recommended.

DOSAGE

Apply 4 to 8 pints (1.0 to 2.0 lb active) of Goal 2XL per broadcast acre for preemergence and postemergence weed control. The addition of 1 quart of Latron AG-98 or a comparable 80% active nonionic surfactant per 100 gallons of spray mix will assist in spray coverage and wetting of weeds for postemergence control.

WEEDS CONTROLLED

When Goal 2XL is applied preemergence or postemergence to weed seedlings (not exceeding 6-leaf stage) at recommended dosages, the following broadleaf weeds are controlled:

WEEDS CONTROLLED POSTEMERGENCE

- | | |
|----------------------|--------------------|
| CHEESEWEED (MALVA) | MINERSLETTUCE |
| FIDDLENECK, COAST | NETTLE, BURNING |
| † FILAREE, BROADLEAF | PIGWEEED, REDROOT |
| † FILAREE, REDSTEM | REDMAIDS |
| † FILAREE, WHITESTEM | SHEPHERDSPURSE |
| GROUNDSEL, COMMON | SOWTHISTLE, ANNUAL |
| HENBIT | |

WEEDS CONTROLLED PREEMERGENCE

- | | |
|--------------------|-------------------|
| BURCLOVER | LETTUCE, PRICKLY |
| CHEESEWEED (MALVA) | PIGWEEED, REDROOT |
| FIDDLENECK, COAST | PURSLANE, COMMON |
| FILAREE, BROADLEAF | REDMAIDS |
| FILAREE, REDSTEM | ROCKET, LONDON |

FILAREE, WHITESTEM	SHEPHERDSPURSE
GROUNDSEL, COMMON	SOWTHISTLE, ANNUAL
HENBIT	SPURGE, PROSTRATE
KNOTWEED, PROSTRATE	SPURGE, SPOTTED
LAMBSQUARTERS, COMMON	

† Goal 2XL at the 8-pint rate (2.0 lb active) will provide control of filaree up to the 6-leaf stage.

TIMING AND METHOD OF APPLICATION

For optimum weed control, Goal 2XL should be applied prior to weed emergence. Postemergence applications should be applied to seedling weeds (up to the 6-leaf stage). Applications must be made prior to bud break of either transplants or established eucalyptus trees.

Goal 2XL should be applied at 20 to 40 psi in a minimum of 20 gallons of water per acre depending upon density of emerged weeds. Spray volume should be increased as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

**EUCALYPTUS
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Goal 2XL should only be applied to dormant healthy eucalyptus stock.
- Do not apply more than 8 pints (2.0 lb active) per treated acre per growing season as a result of single or multiple applications.

**FALLOW BED
GROUND OR AERIAL APPLICATION OF GOAL 2XL ON FALLOW BEDS**

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphomax) for the control of winter annual broadleaf weeds to be planted to the crops listed below.

MINIMUM TREATMENT-PLANTING INTERVAL

DIRECT SEEDED CROPS	Goal 2XL Use Rate	
	up to 1 pint/A	up to 2 pints/A
CARROT	90 DAYS	90 DAYS
POTATO	60 DAYS	60 DAYS
SUGAR BEET	60 DAYS	90 DAYS
OTHER ROOT/TUBER CROPS	90 DAYS	90 DAYS
ONIONS	180 DAYS	180 DAYS
OTHER BULB VEGETABLES	180 DAYS	180 DAYS
CABBAGE	90 DAYS	90 DAYS
CAULIFLOWER	90 DAYS	90 DAYS
OTHER BRASSICA CROPS	120 DAY	120 DAYS
LETTUCE	90 DAYS	120 DAYS
OTHER LEAFY VEGETABLES (EXCEPT BRASSICA CROPS)	120 DAYS	120 DAYS
PEPPER	90 DAYS	120 DAYS
TOMATO	60 DAYS	120 DAYS

OTHER FRUITING VEGETABLES	120 DAYS	120 DAYS
CANTALOUPE	60 DAYS	90 DAYS
SQUASH	90 DAYS	120 DAYS
WATERMELON	60 DAYS	60 DAYS
OTHER CUCURBITS	90 DAYS	120 DAYS
DRY BEANS	60 DAYS	60 DAYS
PEANUT	60 DAYS	60 DAYS
OTHER LEGUME VEGETABLES	60 DAYS	60 DAYS
SAFFLOWER	60 DAYS	60 DAYS
CEREAL GRAINS	10 MONTHS	10 MONTHS
(includes barley, buckwheat, corn, proso millet, pearl millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, wild rice)		

COTTON AND SOYBEAN (See
specific labeling for fallow beds to
be planted to cotton or soybeans)

TRANSPLANTED CROPS	Goal 2XL Use Rate	
	up to 1 pint/A	up to 2 pints/A
BROCCOLI	0 DAYS	30 DAYS
CABBAGE	0 DAYS	30 DAYS
CAULIFLOWER	0 DAYS	30 DAYS
CELERY	30 DAYS	30 DAYS
CONIFER	0 DAYS	0 DAYS
GARLIC	0 DAYS	30 DAYS
GRAPE/KIWI	0 DAYS	0 DAYS
ONION	0 DAYS	30 DAYS
PEPPER	30 DAYS	30 DAYS
STRAWBERRIES	30 DAYS	30 DAYS
TOMATO	30 DAYS	30 DAYS
TREEFRUIT/NUT/CITRUS	0 DAYS	0 DAYS

IMPORTANT:

The fallow beds should be worked thoroughly to a depth of at least 2.5 inches prior to planting; weed control should not be expected following breaking of the soil surface. FAILURE TO ACHIEVE THOROUGH AND COMPLETE INCORPORATION, OR TO FOLLOW THE RECOMMENDED TREATMENT-PLANTING INTERVAL, MAY RESULT IN STAND REDUCTION AND/OR VIGOR REDUCTION OF THE PLANTED CROP.

Crop injury may be enhanced if newly seeded crops or transplants are under stress due to drought, flooding, excessive fertilizer or soil salts, low soil temperatures, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.

EXERCISE EXTREME CARE TO AVOID HERBICIDE CONTACT WITH ANY DESIRABLE DORMANT OR NON-DORMANT CROP, PLANT, TREE OR VEGETATION AS SEVERE INJURY MAY RESULT.

GOAL 2XL USED ALONE

DOSAGE

Goal 2XL may be applied at 1 to 2 pints (0.25 to 0.5 lb active) per broadcast acre. The lower rate (1 pint per acre) should provide up to 4 weeks of preemergence control of susceptible weeds and provide postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre)

should provide preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible weeds (up to 6-leaf stage). Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

WEEDS CONTROLLED

Goal 2XL should provide preemergence and postemergence † control of the following weeds when used at recommended dosages and weed stage.

CHEESEWEED (MALVA)	MUSTARD, SPECIES
FIDDLENECK, COAST	NETTLE, BURNING
FILAREE, BROADLEAF	REDMAIDS
FILAREE, REDSTEM	ROCKET, LONDON
GROUNDSEL, COMMON	SHEPHERDSPURSE
HENBIT	SOWTHISTLE, ANNUAL
MINERSLETTUCE	

† Thorough spray coverage is essential to maximize the postemergence activity of Goal 2XL. For postemergence control when applied by air, a tank mixture of Goal 2XL with glyphosate (Glyphomax) is recommended.

GOAL 2XL is a contact herbicide, therefore, coverage is essential for acceptable postemergence control. If dense weed populations, oversized weed seedlings, volunteer grains, annual grasses or unfavorable environmental conditions exist, a tank mixture of Goal 2XL with glyphosate (Glyphomax) for postemergence control is recommended.

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

DOSAGE

Goal 2XL can be tank mixed with glyphosate (Glyphomax) to obtain postemergence control of annual grassy weeds, volunteer grains and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lb active) of Goal 2XL with labeled rates of glyphosate (Glyphomax). Apply at the recommended rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

METHOD OF APPLICATION

GROUND APPLICATION

Goal 2XL should be applied in a minimum of 20 gallons of water per acre. The volume of water used should be increased as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

AERIAL APPLICATION

Goal 2XL should be applied using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 10 gallons per acre (minimum 5 GPA for GOAL 2XL/glyphosate (Glyphomax) tank mix). Applications should be made at a height of 6 to 10 feet above the soil surface. It is suggested that the nozzles on the spray booms should not be placed any closer to the wing or rotor tips than 3/4 of the span; this will minimize the formation of spray or wing tip vortice roll. Nozzles should be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100 microns or less in diameter.

AVOID DRIFT

WHEN APPLYING TO FALLOW BEDS, EXTREME CARE MUST BE EXERCISED TO PREVENT SPRAY DRIFT WHICH COULD RESULT IN DAMAGE TO OTHER CROPS OR DESIRABLE VEGETATION. USE THE FOLLOWING GUIDELINES WHEN AERIAL APPLICATIONS ARE TO BE MADE.

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least 1/2 mile from all crops and desirable vegetation, except the following:
Maintain a minimum downwind buffer zone of:
 - 150 feet from dormant treefruit/nut/vine crops and overwintering sugar beets.
 - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets, and non-targeted vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any non-targeted vegetable fallow bed, crop, or desirable vegetation.

The use of a drift control agent may be required by local regulations. However, the drift control agent may decrease the weed control activity.

IMPORTANT

Aerial applicators must be familiar with the EPA-registered label and follow the use precautions. Spraying Goal 2XL in a manner other than as recommended is done at the user's risk. Users are responsible for all loss or damage that results from such spraying. In addition, aerial applicators should follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, the most restrictive situations should apply to avoid drift hazards.

FALLOW BED - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per acre per fallow season.

FALLOW BED (COTTON/SOYBEANS)**GROUND OR AERIAL APPLICATION OF GOAL 2XL ON FALLOW BEDS (TO BE PLANTED TO COTTON OR SOYBEANS)****NOT FOR USE ON FALLOW BEDS TO BE PLANTED TO SOYBEANS IN CALIFORNIA****GENERAL INFORMATION**

Goal 2XL is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphomax) or paraquat (Gramoxone) for the control of winter annual broadleaf weeds in fallow beds to be planted to either cotton or soybeans. Do not apply Goal 2XL within 7 days prior to planting. The fallow beds should be worked thoroughly to a depth of at least 2 inches prior to planting. It is important to thoroughly break the soil surface prior to planting. Weed control should not be expected following breaking of the soil surface.

EXERCISE EXTREME CARE TO AVOID HERBICIDE CONTACT WITH ANY DESIRABLE DORMANT OR NON-DORMANT CROP, PLANT, TREE OR VEGETATION AS SEVERE INJURY MAY RESULT.

GOAL 2XL USED ALONE**DOSAGE**

Goal 2XL may be applied at 1 to 2 pints (0.25 to 0.5 lb active) per broadcast acre. The lower rate (1 pint per acre) should provide up to 4 weeks of preemergence control of susceptible weeds and provide postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre) should provide preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible weeds (up to 6-leaf stage). Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

WEEDS CONTROLLED

Goal 2XL should provide preemergence and postemergence[†] control of the following weeds when used at recommended dosages and weed stage.

BUTTERCUP, SMALLFLOWER	MUSTARD SPECIES
CHEESEWEED (MALVA)	NETTLE, BURNING
†† EVENINGPRIMROSE, CUTLEAF	OXALIS
FIDDLENECK, COAST	PIGWEEED, REDROOT
FILAREE, BROADLEAF	PURSLANE, COMMON
FILAREE, REDSTEM	REDMAIDS
GERANIUM, CAROLINA	ROCKET, LONDON
GROUNDCHERRY, CUTLEAF	SHEPHERDSPURSE
GROUNDSEL, COMMON	SIDA, PRICKLY
HENBIT	SOWTHISTLE, ANNUAL
LADYSTHUMB	VELVETLEAF (WILD COTTON)
MINERSLETTUCE	

[†] Thorough spray coverage is essential to maximize the postemergence activity of Goal 2XL. For postemergence control when applied by air, a tank mixture of Goal 2XL with either glyphosate (Glyphomax) or paraquat (Gramoxone) is recommended.

^{††} Requires maximum rate and/or multiple applications for effective control.

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

DOSAGE

Goal 2XL can be tank mixed with either glyphosate (Glyphomax) or paraquat (Gramoxone) to obtain postemergence control of annual grassy weeds, volunteer grains and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lb active) of Goal 2XL with labeled rates of either glyphosate (Glyphomax) or paraquat (Gramoxone). Apply at the recommended rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

OUTSIDE OF CALIFORNIA: For enhanced contact activity (burndown/suppression) to either glyphosate (Glyphomax) or paraquat (Gramoxone), add Goal 2XL at a rate of 3.5 to 6.5 ounces (0.05 to 0.1 lb. active) per acre to labeled rates of either glyphosate (Glyphomax) or paraquat (Gramoxone). Apply at the recommended rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

If a fallow bed treatment is applied thirty days or more prior to planting and at least three significant rainfalls (0.25 inch or greater) have occurred following application, cotton or soybeans can be planted directly into the stale seedbed. If these conditions cannot be met, soil incorporation is required as directed above.

METHOD OF APPLICATION**GROUND APPLICATION**

Goal 2XL should be applied in a minimum of 20 gallons of water per acre. The volume of water used should be increased as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

AERIAL APPLICATION

Goal 2XL should be applied using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 5 gallons per acre (in California, minimum 10 GPA when applied alone or tank mixed with paraquat [Gramoxone]). Applications should be made at a height of 6 to 10 feet above the soil surface. It is suggested that the nozzles on the spray booms should not be placed any closer to the wing or rotor tips than 3/4 of the span; this will minimize the formation of spray or wing tip vortices. Nozzles should be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100 microns or less in diameter.

AVOID DRIFT

WHEN APPLYING TO FALLOW BEDS, EXTREME CARE MUST BE EXERCISED TO PREVENT SPRAY DRIFT WHICH COULD RESULT IN DAMAGE TO OTHER CROPS OR DESIRABLE VEGETATION. USE THE FOLLOWING GUIDELINES WHEN AERIAL APPLICATIONS ARE TO BE MADE.

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least 1/2 mile from all crops and desirable vegetation, except the following:
Maintain a minimum downwind buffer zone of:
 - 150 feet from dormant treefruit/nut/vine crops and overwintering sugar beets.
 - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets, and non-targeted vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any non-targeted vegetable fallow bed, crop, or desirable vegetation.

The use of a drift control agent may be required by local regulations. However, the drift control agent may decrease the weed control activity.

IMPORTANT

Aerial applicators must be familiar with the EPA-registered label and follow the use precautions. Spraying Goal 2XL in a manner other than as recommended is done at the user's risk. Users are responsible for all loss or damage that results from such spraying. In addition, aerial applicators should follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, the most restrictive situations should apply to avoid drift hazards.

FALLOW BED (COTTON/SOYBEANS)**SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per acre per fallow season.
- Do not apply Goal 2XL within 7 days prior to planting of cotton or soybeans.

FALLOW LAND

FOR USE ONLY IN IDAHO, OREGON AND WASHINGTON

GENERAL INFORMATION

Goal 2XL is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphomax) for the control of certain annual broadleaf weeds in a fallow land system. Goal 2XL can be used as an effective tool to reduce weed growth prior to the establishment of a dry soil mulch. Use of this product is restricted to summer fallow land that will be planted back the following year to winter wheat, barley or oats.

GOAL 2XL USED ALONE

DOSAGE

Goal 2XL should be used at 0.5 to 2 pints (0.12 to 0.5 lb active) per broadcast acre.

WEEDS CONTROLLED

Goal 2XL will provide postemergence control and preemergence activity of the following broadleaf weeds when used at recommended dosages.

- | | |
|-------------------|--------------------|
| FIDDLENECK, COAST | MUSTARD, TUMBLE |
| HENBIT | (JIM HILL MUSTARD) |
| LETTUCE, PRICKLY | PIGWEEED, REDROOT |
| (CHINA LETTUCE) | PURSLANE, COMMON |
| MUSTARD, BLUE | SHEPHERDSPURSE |
| (PURPLE MUSTARD) | SOWTHISTLE, ANNUAL |

TIMING AND METHOD OF APPLICATION

The most effective postemergence weed control is achieved when Goal 2XL is applied to seedling weeds (less than 4 inches in height). Seedling weeds are controlled as they come in contact with the soil-applied herbicide during emergence.

Goal 2XL should be applied in 20 to 40 gallons of water per acre depending upon density of emerged weeds. Use a low pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

TANK MIXES WITH GOAL 2XL

DOSAGE

For postemergence control of annual grassy weeds, Goal 2XL can be tank mixed with glyphosate (Glyphomax). Tank mix 0.5 to 2 pints (0.12 to 0.5 lb active) of Goal 2XL with 0.75 to 1 pint (0.38 to 0.5 lb active) of glyphosate (Glyphomax) for each acre treated. Refer to the FALLOW AND REDUCED TILLAGE SYSTEM section on the glyphosate (Glyphomax) label for specific use directions and restrictions.

Fill the spray tank at least one-third full of clean water and add the recommended amounts of Goal 2XL and glyphosate (Glyphomax) while the pump and agitator are running. Complete filling of the spray tank with water. Add 1 quart of Latron AG-98 or comparable 80% active nonionic surfactant cleared for use on growing crops, per 100 gallons of spray. Maintain agitation until spraying is complete.

FALLOW LAND - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

GARBANZO BEANS **FOR USE ONLY IN ARIZONA AND CALIFORNIA**

GENERAL INFORMATION

Goal 2XL is effective as a preemergence herbicide when used alone for the control of certain annual broadleaf weeds in garbanzo beans. Preemergence control is most effective when spray is applied to clean, weed-free soil surfaces. Seedling weeds are controlled as they come in contact with soil-applied herbicide during emergence. Timely cultivations will usually assist in weed control.

Garbanzo beans are tolerant to preemergence applications of Goal 2XL, however, under certain conditions, Goal 2XL can cause severe but temporary crop injury. Heavy splashing rain shortly after crop emergence or wet soil conditions during early growth stages can produce leaf cupping, crinkling, stunting or defoliation of the garbanzo seedlings. When injury occurs, it is often limited to the first few leaves that develop shortly after crop plants emerge from the soil. Delays in crop development and/or maturity may result. Garbanzo beans do recover from this injury with little to no impact on yield.

GOAL 2XL USED ALONE

DOSAGE

Goal 2XL is recommended for preemergence control of susceptible winter annual broadleaf weeds at 1 pint (0.25 lb active) per broadcast acre.

WEEDS CONTROLLED PREEMERGENCE

Goal 2XL used alone at recommended dosages provides preemergence control of the following broadleaf weeds:

GROUNDSEL, COMMON
MALLOW, LITTLE

ROCKET, LONDON
SHEPHERDSPURSE

TIMING AND METHOD OF APPLICATION

As a preemergence application, apply in a minimum of 25 gallons of water per acre. Use conventional ground spray equipment to make a single broadcast application, after planting but prior to weed and crop emergence, with flat fan or hollow cone nozzles. Spray equipment should be calibrated carefully before each use.

GARBANZO BEANS - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 1 pint (0.25 lb active) per broadcast acre of Goal 2XL in a single application.
- For application only in Arizona and California.
- Do not feed bean, vines or hay.

GARLIC

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for postemergence application to direct-seeded and transplanted garlic for early postemergence control of certain annual broadleaf and grass weeds. Initial spray application should be made only when the garlic have reached the development stage specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label. On garlic transplants spray as soon

after transplanting as practical. Goal 2XL can cause necrotic lesions, twisting, pigtailling or stunting of the garlic plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the development stage of the garlic plants as specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label.

DOSAGE

SEEDED GARLIC

NORTHEASTERN STATES

CONNECTICUT, MAINE, MASSACHUSETTS, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, RHODE ISLAND AND VERMONT

Goal 2XL is recommended for postemergence control at 2 to 4 fluid ounces (0.03 to 0.06 lb active) per acre when applied postemergence to seeded garlic that has at least three (3) true leaves. Multiple treatments at the aforementioned rate may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

WESTERN STATES

ARIZONA, COLORADO, IDAHO, NEVADA, NEW MEXICO, OREGON, TEXAS, UTAH AND WASHINGTON

Goal 2XL is recommended for postemergence control at 0.5 to 1 pint (0.12 to 0.25 lb active) per acre when applied postemergence to garlic that has at least two (2) true leaves. Multiple treatments at the aforementioned rates may be applied. Do not apply more than 2.5 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

CALIFORNIA ONLY

GENERAL INFORMATION

GOAL 2XL is a selective for preemergence use (by air, ground, or sprinkler application), post-direct use when applied by ground equipment, or postemergence (over the top) application when applied via sprinkler irrigation for control of certain broadleaf and grass weeds in garlic in California.

Chemigation: For application using only solid set or portable lateral sprinkler irrigation systems, apply Goal 2XL at the recommended broadcast application rate per acre as described below. Follow the application directions for "Sprinkler Chemigation" given in the CHEMIGATION section of this label.

Preemergence Garlic Applications in California

Apply Goal 2XL at a rate of 1 pint (0.25 lb active) per broadcast acre as a preemergence application to garlic. Methods of application may be ground, sprinkler, or aerial.

Ground Application: If applied using ground application equipment, Goal 2XL should be applied in a minimum of 20 gallons per acre. Use conventional ground spray equipment with flat nozzles at 20 to 40 psi.

Sprinkler Chemigation: Apply Goal 2XL at the recommended broadcast application rate. Sufficient sprinkler irrigation water should be applied to insure water penetration to a depth of two inches.

Aerial Application: If applied using aerial application, Goal 2XL should be applied using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 10 gallons per acre. Applications should be made at a height of 6 to 10 feet above the soil surface. It is suggested that the nozzles on the spray booms should not be placed any closer to the wing or rotor tips than $\frac{1}{4}$ of the span; this will minimize the formation of spray or wing tip vortice roll. Nozzles should be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100

microns or less in diameter.

Garlic Response to Preemergence Applications With Goal 2XL: A chlorotic band around some of the leaves may be observed after the first irrigation (or rainfall) following garlic emergence. Symptoms may be more severe if garlic emerges under cool, wet, overcast, or foggy weather. This condition is temporary and should not affect the vigor or development of the garlic plant.

Postemergence (and Directed) Garlic Applications in California

Apply Goal 2XL at rates up to 1 pint (0.25 lb active) per broadcast acre as a postemergence (or directed) application in garlic. The garlic must be at least 12 inches in height at application. Weeds should be in the seedling stage, young, and actively growing. Methods of application may be post-directed or by sprinkler chemigation.

Post Direct Application: For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

Band Width (in inches)	X	Rate per	=	Amount Needed per Acre
Row Width (in inches)		Broadcast Acre		for Banded Application

Accurate, uniform placement of Goal 2XL spray is essential for effective weed control and to minimize garlic injury. As a directed, postemergence application, Goal 2XL should be applied using a low-pressure sprayer using a minimum of 20 gallons of spray on a broadcast acre basis. Apply Goal 2XL as a directed treatment to the soil area at the base of the plants and to the adjacent bed top and furrow areas. Nozzles should be adjusted to cover the weed foliage with minimum contact to the garlic plant. Reduce tractor speed and smooth furrows to minimize excessive bouncing of the spray boom.

Sprinkler Chemigation: Apply Goal 2XL at the recommended broadcast application rate. Sufficient sprinkler irrigation water should be applied to insure water penetration to a depth of two inches.

Garlic Response to Postemergence Applications With Goal 2XL: Goal 2XL may cause chlorotic leaf banding, necrotic lesions, or stunting of the garlic plants. Symptoms will be more severe if applications are made during cool, wet, overcast, or foggy weather. Garlic will outgrow these conditions and continue to develop normally.

AVOID DRIFT: WHEN APPLYING GOAL 2XL TO GARLIC IN CALIFORNIA, EXTREME CARE MUST BE EXERCISED TO PREVENT SPRAY DRIFT WHICH COULD RESULT IN DAMAGE TO OTHER CROPS OR DESIRABLE VEGETATION. WHEN APPLYING BY AIR OR THROUGH SPRINKLER CHEMIGATION SYSTEMS, USE THE FOLLOWING GUIDELINES:

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least ½ mile from all crops and desirable vegetation, except the following:
Maintain a minimum downwind buffer zone of:
 - 150 feet from dormant treefruit, dormant vines, and overwintering sugar beets.
 - 650 feet from jojoba, legumes, small grains, seedling sugar beets, pastures, and vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any vegetable fallow bed, crop, or desirable vegetation.

Cultural Considerations for use in California

On mineral soils, in order to provide maximum preemergence activity, the soil surface should be smooth

and free of excessive trash (clippings, dead weeds, etc.)

Cultural practices that result in redistribution or disturbance of the soil surface after spraying or that mix untreated soil in treated areas will reduce the effectiveness of the treatment. The best results from Goal 2XL are from applications on established beds that are left undisturbed during the time period for which weed control is desired.

ALL OTHER STATES

Goal 2XL is recommended for postemergence control at 0.5 pints (0.12 lb active) per acre when applied postemergence to garlic that has at least two (2) true leaves. Multiple treatments at the aforementioned rates may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

TRANSPLANTED GARLIC

Transplanted garlic is most tolerant of a postemergence application immediately after transplanting. For all states except the northeastern states listed under the DOSAGE - SEEDED GARLIC section, an application of up to 2 pints (0.5 lb active) per acre within two days after transplanting may be made. If less than 2.5 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

For transplanted garlic in the northeastern states, apply the same rates listed in the DOSAGE - SEEDED section within two days after transplanting.

Dosages listed are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

<u>Band Width (in inches)</u>	X	Rate per	=	Amount Needed per Acre
Row Width (in inches)		Broadcast Acre		for Banded Application

WEEDS CONTROLLED

Goal 2XL will provide postemergence control of the following weeds when applied at the recommended dosage and leaf stage (2 to 4 leaves).

- | | |
|--------------------------|--------------------|
| CANARYGRASS (ANNUAL) | PUNCTUREVINE |
| EVENINGPRIMROSE, CUTLEAF | †PURSLANE, COMMON |
| GROUNDSEL, COMMON | ROCKET, LONDON |
| MALLOW, LITTLE (MALVA) | SAGE, LANCELEAF |
| NIGHTSHADE, BLACK | †SHEPHERDSPURSE |
| †PIGWEEED, PROSTRATE | SOWTHISTLE, ANNUAL |
| †PIGWEEED, REDROOT | |

† Specific weeds controlled at rates recommended for use in Northeastern States (see DOSAGE section).

TIMING AND METHOD OF APPLICATION

For best postemergence control of susceptible weeds, apply when the weeds are in the 2 to 4-leaf stage. Application of Goal 2XL after the weeds exceed the maximum leaf stage may result in reduced weed control. More than one postemergence application may be necessary to control subsequent weed flushes.

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 40 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles at 20 to 40 psi. Accurately calibrate spray equipment prior to each use. Avoid drift to all other crops and non-target areas. Thoroughly flush the spray equipment (tank, hose, pump, boom) with water before and after each use. Residual Goal 2XL remaining in spray equipment may damage other crops.

GARLIC - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- In all states except Northeastern states, do not start spraying until the garlic (direct seeded) have two (2) fully developed true leaves. In the Northeastern states (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, Vermont), do not start spraying until the garlic (direct seeded) have three (3) fully developed true leaves. Applications made prior to the recommended garlic development stage may result in serious injury and is not recommended.
- Do not apply more than a total of 2 pints (0.5 lb active) per acre of Goal 2XL during one use season.
- Do not apply within 60 days of harvest.
- Use only on dry bulb garlic.
- Do not apply to garlic grown for seed.
- Do not mix Goal 2XL with oils, surfactants, liquid fertilizers or pesticides except as specified on other approved Rohm and Haas Supplemental Labeling.
- Do not apply Goal 2XL preemergence to direct-seeded garlic.
- Do not apply to garlic plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases.

GUAVA

FOR USE ONLY IN HAWAII

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence herbicide when used alone for the control of certain annual broadleaf weeds in bearing and nonbearing guava plantings. For postemergence control of certain grassy and broadleaf weeds, a tank mixture of either paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL can be applied to seedling weeds. Check individual labels to determine suitability and use rates for crop.

GOAL 2XL USED ALONE

DOSAGE

Goal 2XL is recommended for postemergence control of susceptible weeds at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre.

For preemergence control of susceptible weeds, use 5 to 8 pints (1.25 to 2.0 lb active) of Goal 2XL per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE

Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Applications to weeds beyond the 4-leaf stage may result in partial control.

PURSLANE, COMMON

SPURGE, GARDEN

WEEDS CONTROLLED PREEMERGENCE

Apply 5 to 8 pints (1.25 to 2.0 lb active) of Goal 2XL per broadcast acre.

AGERATUM

PURSLANE, COMMON

BUTTONWEED

SPURGE, GARDEN

CROTALARIA

TIMING AND METHOD OF APPLICATION

Treatments should be applied only to healthy guava trees. Care must be taken to prevent direct spray or drift from contacting green stems, fruit or foliage, as injury may result. Applications should be made only after new foliage has hardened off, or injury may result.

As a preemergence or postemergence treatment to weeds, apply in a minimum of 15 gallons of water per acre. Use higher volumes to assure adequate coverage in high densities of emerged weeds or heavy trash. Goal 2XL should be directed to the soil and the base of the tree. Use of a low pressure sprayer equipped with a breakaway boom and flat fan or off-center (OC) nozzles is recommended. An off-center nozzle positioned at the end of the boom may be desired. Spray shields are suggested for use in young trees.

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

DOSAGE

For postemergence control of susceptible grassy and broadleaf weeds in guava plantings, a tank mixture of Goal 2XL with either paraquat (Gramoxone) or glyphosate (Glyphomax) can be used. Apply at recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

WEEDS CONTROLLED POSTEMERGENCE

In addition to the weeds controlled by Goal 2XL used alone, control of susceptible weeds listed on the respective labels for the following products is also obtained:

paraquat (Gramoxone)
glyphosate (Glyphomax)

GUAVA - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in a single application or more than 16 pints (4.0 lbs active) per season.
- Do not apply Goal 2XL within 1 day of harvest.
- Direct spray toward the base of the trees. Avoid direct plant contact.
- Goal 2XL or any of the combinations recommended on this label should be applied to only healthy growing trees.
- GOAL 2XL applications should be made only after new foliage has hardened off.

HORSERADISH**GENERAL INFORMATION**

GOAL 2XL is a selective herbicide recommended for preemergence control of certain broadleaf weeds. Applications must be made after the horseradish roots have been planted and prior to plant emergence. (Emerged plants that receive direct or indirect (drift) spray contact will be injured.) It may be desirable to cultivate immediately prior to application to remove germinated weeds.

Do not use Goal 2XL on horseradish plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.

DOSAGE

Apply Goal 2XL at a rate of 2 pints (0.5 lb active) per broadcast acre as a preemergence application to horseradish.

WEEDS CONTROLLED

Goal 2XL will provide preemergence control of the following weeds when used at the recommended dosage:

LAMBSQUARTERS, COMMON	SHEPHERDSPURSE
PIGWEEED, REDROOT	SMARTWEED, PENNSYLVANIA
PURSLANE, COMMON	

TIMING AND METHOD OF APPLICATION

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 20 gallons of water per acre. Use conventional ground spray equipment with flat fan nozzles at 20 to 40 psi. Accurately calibrate spray equipment prior to each use.

HORSERADISH - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per broadcast acre as a single application.

JOJOBA**GENERAL INFORMATION**

GOAL 2XL is a selective herbicide for postemergence and preemergence control of certain broadleaf weeds in jojoba. Goal 2XL should be post-directed to the base of the jojoba plant to avoid possible phytotoxicity to the jojoba foliage. Over-the-top applications may exhibit burning, crinkling or bronzing of jojoba foliage, particularly to the youngest leaves, flowers, or buds present at the time of application.

DOSAGE

Goal 2XL is recommended for postemergence and preemergence control of susceptible seedling weeds (up to 12 inches in height) at 8 pints (2.0 lb active) per broadcast acre. For optimum residual control, apply during the fall or winter.

For early postemergence control of susceptible seedling weeds (less than 8 inches in height) apply Goal 2XL at a rate of 4 pints (1.0 lb. active) per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE

FIDDLENECK, COAST	MINERSLETTUCE
†† FILAREE, BROADLEAF	NETTLE, BURNING
†† FILAREE, REDSTEM	† PIGWEED, REDROOT
†† FILAREE, WHITESTEM	REDMAIDS
† GROUNDSEL, COMMON	SHEPHERDSPURSE
HENBIT	SOWTHISTLE, ANNUAL
MALLOW, LITTLE (MALVA, CHEESEWEED)	

WEEDS CONTROLLED PREEMERGENCE

BURCLOVER	LETTUCE, PRICKLY
FIDDLENECK, COAST	MALLOW, LITTLE (MALVA, CHEESEWEED)
FILAREE, BROADLEAF	PIGWEEED, REDROOT
FILAREE, REDSTEM	PURSLANE, COMMON
FILAREE, WHITESTEM	REDMAIDS
GROUNDSEL, COMMON	ROCKET, LONDON
HENBIT	SHEPHERDSPURSE
KNOTWEED, PROSTRATE	SOWTHISTLE, ANNUAL

LAMBSQUARTERS, COMMON

† Highest rate may be required for acceptable postemergence control.

†† Goal 2XL at the 8-pint rate (2.0 lbs active) will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

TIMING AND METHOD OF APPLICATION

Apply the first application of Goal 2XL after jojoba plants have grown to a minimum 6-inch height or greater. Additional applications should be applied as needed for post and preemergence weed control. Weed height should not exceed 12 inches or unsatisfactory weed control may result.

Goal 2XL should be applied in a minimum spray volume of 40 gallons of water per acre depending upon density of emerged weeds. Spray volume should be increased as weed height and density increase. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

JOJOBA - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Avoid direct spray or drift contact of Goal 2XL with jojoba flowers or buds as severe injury may result.
- Do not apply more than 8 pints (2.0 lb active) per broadcast care in a single application.

MINT**SPEARMINT/PEPPERMINT**

FOR USE ONLY IN CALIFORNIA, IDAHO, MONTANA, NEVADA, OREGON, SOUTH DAKOTA, UTAH AND WASHINGTON

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for the control of certain annual grasses and broadleaf weeds in spearmint and peppermint grown in California, Idaho, Montana, Nevada, Oregon, South Dakota, Utah and Washington. Applications should only be made to spearmint and peppermint that are dormant.

METHOD OF APPLICATION

Application must be made prior to new spring growth or severe crop injury may result. Goal 2XL should be thoroughly mixed with clean water at recommended concentration and applied at 20 to 40 psi in 20 to 40 gallons of water per acre.

WEEDS CONTROLLED

When Goal 2XL is applied as a dormant application at recommended dosages in spearmint and peppermint, the following annual weeds are controlled:

† BEDSTRAW, CATCHWEED	NIGHTSHADE, HAIRY
† BLUEGRASS, ANNUAL	† OATS, WILD
FLIXWEED	ORACH, RED
GROUNDSEL, COMMON	PEPPERWEED,
	YELLOWFLOWER
LAMBSQUARTERS, COMMON	PIGWEEED, REDROOT
LETTUCE, PRICKLY	† RYEGRASS, ITALIAN
(CHINA LETTUCE)	SHEPHERDSPURSE
MUSTARD, BLUE	SOWTHISTLE, ANNUAL
(PURPLE MUSTARD)	TANSYMUSTARD
MUSTARD, TUMBLE	THISTLE, RUSSIAN

(JIM HILL MUSTARD)

† Control of annual grasses is best obtained when Goal 2XL is applied prior to emergence. Postemergence control of winter annual grasses is generally unsatisfactory if applications are made after the 1 to 2-leaf stage.

WESTERN OREGON

PEPPERMINT (WILLAMETTE VALLEY)

Apply 2 to 3 pints (0.5 to 0.75 lb active) of Goal 2XL from November through February to dormant peppermint only. Treatments in January or February generally provide better residual preemergence control of annual broadleaf weeds. Full season weed control should not be expected from this treatment.

• DO NOT APPLY GOAL 2XL IN THE WILLAMETTE VALLEY
• TO MINT THAT HAS BEEN PLOWED.

**OREGON AND WASHINGTON (EAST OF CASCADES)
CALIFORNIA, MONTANA , IDAHO, NEVADA, SOUTH DAKOTA AND UTAH.**

SPEARMINT AND PEPPERMINT

Apply 4 to 8 pints (1 to 2 lb active) of Goal 2XL from December through March to dormant mint only. Later winter applications will provide maximum activity on summer weeds. Summer grass control may be inconsistent. For best results, fall-plowed fields should be harrowed to provide a smooth surface prior to application. Plowed fields should not be harrowed after Goal 2XL has been applied, as soil disturbance will decrease the herbicidal effectiveness. In furrow-irrigated fields, corrugating must be done prior to application. Corrugating after application can cover treated rows with untreated soil, resulting in poor weed control.

**MINT (SPEARMINT AND PEPPERMINT)
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than one application of Goal 2XL per season.
- Apply Goal 2XL only to healthy spearmint and peppermint. Do not apply to spearmint or peppermint that has been weakened by disease, drought, flooding, excessive fertilizer, soil salts, previously applied pesticides, nematodes, insects, or winter injury, as severe injury may result.

NON-CROP

RIGHTS-OF-WAY SITES, FENCE ROWS, STORAGE AREAS, ROADSIDES, ETC.

GENERAL INFORMATION

Goal 2XL is recommended for postemergence and preemergence control of certain broadleaf weeds in non-crop areas.

WEEDS CONTROLLED POSTEMERGENCE (weeds up to 4 inches high): Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. The lower rate in the rate range is recommended for control of susceptible weeds in the early postemergence stage, less than 4 inches in height. The higher rate (2.0 lb active) should be used for weeds up to 12 inches in height. Applications to weeds beyond the 4-inch stage may result in partial control.

**WEEDS CONTROLLED POSTEMERGENCE
CHEESEWEED (MALVA)**

FIDDLENECK, COAST
 FILAREE, BROADLEAF
 FILAREE, REDSTEM
 GROUNDSEL, COMMON
 HENBIT
 MINERSLETTUCE
 NETTLE, BURNING
 PIGWEED, REDROOT
 PURSLANE, COMMON
 REDMAIDS
 SHEPHERDSPURSE
 SOWTHISTLE, ANNUAL

WEEDS CONTROLLED PREEMERGENCE: Apply 5 to 8 pints (1.25 to 2.0 lb active) per broadcast acre.

WEEDS CONTROLLED PREEMERGENCE

BURCLOVER
 CHEESEWEED (MALVA)
 FIDDLENECK, COAST
 FILAREE, BROADLEAF
 FILAREE, REDSTEM
 GROUNDSEL, COMMON
 HENBIT
 KNOTWEED, PROSTRATE
 LAMBSQUARTERS, COMMON
 LETTUCE, PRICKLY
 PIGWEED, REDROOT
 PURSLANE, COMMON
 REDMAIDS
 ROCKET, LONDON
 SHEPHERDSPURSE
 SOWTHISTLE, ANNUAL

TIMING AND METHOD OF APPLICATION

GOAL 2XL should be applied in a minimum of 40 gallons of water per acre. Best preemergence results are achieved when spray is applied to a relatively weed free soil surface. The volume of water used should be increased as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

DOSAGE

For preemergence control of susceptible grassy and broadleaf weeds, a tank mixture of Goal 2XL with diuron (Karmex) or simazine can be applied. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

For postemergence control of susceptible grass and broadleaf weeds, a tank mixture with paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL can be used. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not feed or allow animals to graze on any areas treated with Goal 2XL.

ONIONS**GENERAL INFORMATION**

GOAL 2XL is a selective herbicide for postemergence application to direct-seeded and transplanted onions for early postemergence control of certain annual broadleaf and grass weeds. Initial spray application should be made only when the onions have reached the development stage specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label. On onion transplants spray as soon before or after transplanting as practical. Goal 2XL can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the development stage of the onion plants as specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label.

DOSAGE**SEEDED ONIONS****NORTHEASTERN STATES: CONNECTICUT, MAINE, MASSACHUSETTS, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, RHODE ISLAND AND VERMONT**

Goal 2XL is recommended for postemergence control at 2 to 4 fluid ounces (0.03 to 0.06 lb active) per acre when applied postemergence to seeded onions that have *at least three (3) true leaves*. Multiple treatments at the aforementioned rate may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

WESTERN STATES: ARIZONA, CALIFORNIA, COLORADO, IDAHO, NEVADA, NEW MEXICO, OREGON, TEXAS, UTAH AND WASHINGTON

Goal 2XL is recommended for postemergence control at 0.5 to 1 pint (0.12 to 0.25 lb active) per acre when applied postemergence to onions that have *at least two (2) true leaves*. Multiple treatments at the aforementioned rates may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

CHEMIGATION: For application using sprinkler irrigation (solid set or portable lateral) systems, apply specified dosage of GOAL 2XL per acre as described in DOSAGE- SEEDED ONIONS-WESTERN STATES section above. Follow all directions given in the CHEMIGATION section of this label when making applications using irrigation systems.

AVOID DRIFT

WHEN APPLYING TO ONIONS, EXTREME CARE MUST BE EXERCISED TO PREVENT SPRAY DRIFT WHICH COULD RESULT IN DAMAGE TO OTHER CROPS OR DESIRABLE VEGETATION. USE THE FOLLOWING GUIDELINES WHEN APPLICATIONS OF GOAL 2XL ARE TO BE MADE THROUGH THE SPRINKLER IRRIGATION SYSTEM:

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least ½ mile from all crops and desirable vegetation, except for the following:
Maintain a minimum downwind buffer zone of:
 - 150 feet from dormant treefruit, dormant vines and overwintering sugar beets.

- 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets and vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
 4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any vegetable fallow bed, crop, or desirable vegetation.

ALL OTHER STATES

Goal 2XL is recommended for postemergence control at 0.5 pints (0.12 lb active) per acre when applied postemergence to onions that have *at least two (2) true leaves*. Multiple treatments at the aforementioned rates may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

TRANSPLANTED ONIONS

POST TRANSPLANT: Transplanted onions are most tolerant of a postemergence application immediately after transplanting. For all states except the northeastern states listed under the DOSAGE - SEEDED ONIONS section, an application of up to 2 pints (0.5 lb active) per acre within two days after transplanting may be made. If less than 2 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season. For transplanted onions in the northeastern states, apply the same rates listed in the DOSAGE - SEEDED ONIONS section within two days after transplanting.

PRE-TRANSPLANT: (Not for use in Northeastern or Western states, except as specifically directed on other approved supplemental labeling.) Goal 2XL is recommended for use as a pre-transplant application at 1 to 2 pints (0.25 to 0.5 lb active) per broadcast acre. Applications must be made after completion of soil preparation, but prior to transplanting of onion plants. Transplanting should be completed with minimal soil disturbance. Treated soil surfaces should be left undisturbed after transplanting to obtain greatest benefit of GOAL herbicide on susceptible annual broadleaf weeds during the time period for which weed control is desired. However, timely cultivations after weed emergence will assist in weed control. If less than 2 pints per acre is applied as a pre-transplant treatment, postemergence applications can be made as instructed in the DOSAGE - SEEDED ONIONS section of this label. Do not exceed the maximum use rate of 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

Dosages listed are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

WEEDS CONTROLLED

Goal 2XL will provide postemergence control of the following weeds when applied at the recommended dosage and leaf stage (2 to 4 leaves):

- | | | | |
|----|--------------------------|----|--------------------|
| a | CANARYGRASS (ANNUAL) | ab | PUNCTUREVINE |
| | EVENINGPRIMROSE, CUTLEAF | | PURSLANE, COMMON |
| | GROUNDSEL, COMMON | | ROCKET, LONDON |
| | MALLOW, LITTLE (MALVA) | b | SAGE, LANCELEAF |
| | NIGHTSHADE, BLACK | | SHEPHERDSPURSE |
| b | PIGWEEED, PROSTRATE | | SOWTHISTLE, ANNUAL |
| ab | PIGWEEED, REDROOT | | |

- a Weeds controlled when applied as a pre-transplant application. In addition, Goal 2XL at the rate of 1 to 2 pints per acre will provide control/suppression of carpetweed, Pennsylvania smartweed, galinsoga, common lambsquarters, and wild mustard. Applications of Goal 2XL to muck soils may result in partial control or suppression of the weeds listed.
- b Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

TIMING AND METHOD OF APPLICATION

For best postemergence control of susceptible weeds, apply when the weeds are in the 2-to 4-leaf stage. Application of Goal 2XL after the weeds exceed the maximum leaf stage may result in reduced weed control. More than one postemergence application may be necessary to control subsequent weed flushes.

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 40 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles at 20 to 40 psi. Accurately calibrate spray equipment prior to each use. Avoid drift to all other crops and non-target areas. Thoroughly flush the spray equipment (tank, hose, pump, boom) with water before and after each use. Residual Goal 2XL remaining in spray equipment may damage other crops.

ONIONS - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- In all states, except Northeastern states, do not start spraying until the onions (direct-seeded) *have two (2) fully developed true leaves*. In the Northeastern states (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, Vermont), do not start spraying until the onions (direct-seeded) *have three (3) fully developed true leaves*. Applications made prior to the recommended onion development stage may result in serious injury and is not recommended.
- Do not apply more than a total of 2 pints (0.5 lb active) per acre of Goal 2XL during one use season.
- Do not apply within 45 days of harvest.
- Use only on dry bulb onions.
- Do not apply to onions grown for seed, except as specified on other approved supplemental labeling.
- Tank mixtures of Goal 2XL with oils, surfactants, liquid fertilizers or other pesticides may result in enhanced crop response/injury and are the responsibility of the user.
- Do not apply Goal 2XL preemergence to direct-seeded onions.
- Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases.

ONIONS GROWN FOR SEED

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for postemergence application to onions grown for seed, for early postemergence control of certain annual broadleaf and grassy weeds. Initial spray application should be made only when the onions have reached the development stage specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label. Goal 2XL can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the development stage of the onion plants as specified in the DOSAGE section and the SPECIFIC USE RESTRICTIONS section of this label.

NOTICE: Some varieties or inbred lines of onions may be more susceptible to Goal 2XL. Care should be taken to insure that the particular onion variety or line being grown is tolerant to Goal 2XL. It is suggested that all onion varieties or lines be tested in limited areas to ensure an adequate level of crop tolerance prior to an application for postemergence weed control.

WEEDS CONTROLLED

Goal 2XL will provide postemergence control of the following weeds when applied at the recommended dosage and leaf stage (2 to 4 leaves):

CANARYGRASS (ANNUAL)	PUNCTUREVINE
EVENINGPRIMROSE, CUTLEAF †	PURSLANE, COMMON
GROUNDSEL, COMMON	ROCKET, LONDON
MALLOW, LITTLE (MALVA)	SAGE, LANCELEAF
NIGHTSHADE, BLACK	SHEPHERDSPURSE
† PIGWEED, PROSTRATE	SOWTHISTLE, ANNUAL
† PIGWEED, REDROOT	

† Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

DOSAGE**NORTHEASTERN STATES**

CONNECTICUT, MAINE, MASSACHUSETTS, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, RHODE ISLAND AND VERMONT

Goal 2XL is recommended for postemergence control at a maximum use rate of 2 fluid ounces (0.03 lb active) per acre when applied postemergence to seeded onions that have *at least four (4) true leaves*. Multiple treatments at the aforementioned rate may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL as a result of multiple applications in one season.

ALL OTHER STATES

Goal 2XL is recommended for postemergence control at a maximum use rate of 0.5 pints (0.125 lb active) per acre when applied postemergence to onions that have *at least three (3) true leaves*. Multiple treatments at the aforementioned rates may be applied. Do not apply more than 2 pints (0.5 lb active) per broadcast acre of Goal 2XL in one season.

TIMING AND METHOD OF APPLICATION

For best postemergence control of susceptible weeds, apply when the weeds are in the 2 to 4-leaf stage. Application of Goal 2XL after the weeds exceed the maximum leaf stage may result in reduced weed control. More than one postemergence application may be necessary to control subsequent weed flushes.

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 40 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles, at 20 to 40 psi. Do not exceed 40 psi. Accurately calibrate spray equipment prior to each use. Thoroughly flush the spray equipment (tank, hose, pump, boom) with water before and after each use. Residual Goal 2XL remaining in spray equipment may damage other crops.

DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT. AVOID DRIFT TO ALL NON-TARGET AREAS. GOAL 2XL IS PHYTOTOXIC TO PLANT FOLIAGE.

CHEMIGATION: For application using sprinkler irrigation (solid set or portable lateral) systems, apply specified dosage of GOAL 2XL per acre as described above. Follow all directions given in the CHEMIGATION section of this label when making applications using irrigation systems.

AVOID DRIFT

WHEN APPLYING TO ONIONS, EXTREME CARE MUST BE EXERCISED TO PREVENT SPRAY DRIFT WHICH COULD RESULT IN DAMAGE TO OTHER CROPS OR DESIRABLE VEGETATION. USE THE FOLLOWING GUIDELINES WHEN APPLICATIONS OF GOAL 2XL ARE TO BE MADE THROUGH THE SPRINKLER IRRIGATION SYSTEM:

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least ½ mile from all crops and desirable vegetation, except for the following:
Maintain a minimum downwind buffer zone of:
 - 150 feet from dormant treefruit, dormant vines and overwintering sugar beets.
 - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets and vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any vegetable fallow bed, crop, or desirable vegetation.

**ONIONS GROWN FOR SEED
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- In all states, do not start spraying until the onions have reached the minimum leaf stage specified in the DOSAGE section of this label. Applications made prior to recommended onion development stage may result in serious injury and is not recommended.
- Do not apply more than a total of 2 pints (0.5 lb active) per acre of Goal 2XL during one use season.
- Do not apply within 60 days of harvest.
- Do not mix Goal 2XL with oils, surfactants, liquid fertilizers or other pesticides except as specified on other approved Rohm and Haas Company Supplemental Labeling.
- Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.

**PAPAYA
FOR USE ONLY IN HAWAII**

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for use as a post-directed application for broadleaf weed control in papaya. Occasionally, after the use of Goal 2XL, a spotting, crinkling or flecking may appear on the leaves of the papaya. Leaves or green stalks that receive direct or indirect (drift) spray contact will be injured.

Do not use Goal 2XL on papaya plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.

DOSAGE AND TIMING

Apply Goal 2XL at a rate of 4 pints (1.0 lb. active) per broadcast acre as a directed spray to the orchard floor. The initial application should occur no earlier than 4 months after transplanting or 6 months after direct seeding, and after the papaya has reached a minimum height of 4 feet. Applications may be repeated at approximate 4-month intervals.

Goal 2XL provides effective control of susceptible weed seedlings in the 4-leaf stage. Do not apply more than 4.0 pints (1.0 lb. active) of GOAL 2XL per broadcast acre in a single application, or more than 12.0 pints (3.0 lb active) per broadcast acre per year as a result of multiple applications.

WEEDS CONTROLLED

Goal 2XL will provide preemergence and postemergence control of the following weeds when used at the recommended dosage. Application to weeds beyond the 4-leaf stage may result in partial control:

AMARANTH, SPINY
PURSLANE, COMMON

SPURGE, GARDEN

METHOD OF APPLICATION

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 15 gallons of water per broadcast acre. Accurately calibrate spray equipment prior to each use.

Accurate, uniform placement of Goal 2XL is essential for effective weed control and to minimize crop injury. GOAL 2XL must be applied as a directed spray to the orchard floor beneath the papaya plants. Do not allow the herbicide solution, spray, drift or mist to contact green bark, stems, fruit or foliage as injury may result. Goal 2XL must be applied using rigid precision ground sprayer equipment.

PAPAYA - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not allow herbicide solution, spray, drift or mist to contact green bark, stems, fruit or foliage as injury may result.
- Do not apply more than 4.0 pints (1.0 lb. active) of Goal 2XL per broadcast acre in a single directed spray or more than 12 pints (3.0 lb active) per broadcast acre per year as a result of multiple applications.
- Do not apply Goal 2XL within 1 day of harvest.
- For use only in papaya grown in Hawaii.

SOYBEANS

NOT FOR USE IN CALIFORNIA

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence and postemergence (post-directed) herbicide for the control of broadleaf weeds in soybeans. Applications can be made early preplant in conservation tillage soybeans, preemergence in no-till (double-crop) and conventional soybeans, or post-directed in conventional till soybeans. Seedling weeds are controlled as they come in contact with the herbicide either during emergence or through a post-directed application. Follow specific use directions and restrictions for recommended use and timing of applications.

Soybeans are tolerant to preemergence and post-directed applications of recommended dosages of Goal 2XL; however, under certain conditions, Goal 2XL can cause temporary injury. Heavy splashing rain shortly after crop emergence or cold, wet soil conditions during early growth stages can produce leaf cupping and crinkling. When injury occurs, it is generally limited to the first few leaves that develop shortly after crop plants emerge from the soil. Soybeans recover from this injury and yields are not adversely affected. Soybean leaves that are accidentally sprayed during a post-directed application will exhibit necrotic spotting and injury to the soybean plant. Therefore, care must be exercised to avoid spray contact with the soybean leaves.

DOSAGE AND TIMING

CONSERVATION TILLAGE

SOYBEANS EARLY PREPLANT

Goal 2XL is effective for preemergence and postemergence control of susceptible broadleaf weeds when surface applied at 1.5 to 3 pints (0.38 to 0.75 lb active) per broadcast acre to the stale seedbed prior to the planting of conservation tillage soybeans. It is suggested that applications be made approximately 14 days prior to planting. The higher rate of 2 to 3 pints (0.5 to 0.75 lbs active) will assist in early season

annual grass control. However, Goal 2XL must not be a basic portion of the grass herbicide program. A planned program utilizing herbicides registered for early preplant, preemergence or postemergence grass control in soybeans is recommended.

The use of ridge or slot planters or other planting equipment that results in minimal soil disturbance is recommended. Soil surfaces should not be disturbed as the herbicidal effectiveness of GOAL 2XL may be decreased. Seedling weeds are controlled as they come in contact with the soil-applied herbicide during emergence. Timely cultivations will usually assist in weed control.

NO-TILL (DOUBLE-CROP) SOYBEANS

PREEMERGENCE

Goal 2XL is effective for preemergence and postemergence control of susceptible broadleaf weeds when applied at 0.5 to 2 pints (0.12 to 0.5 lb active) per broadcast acre. For postemergence control of certain grassy and broadleaf weeds a tank mix of either paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL can be used. For residual grass control in no-till soybeans, a tank mixture of Bronco herbicide, Dual, Lasso, or Surflan with Goal 2XL or combinations of Goal 2XL plus paraquat (Gramoxone) or glyphosate (Glyphomax) can be used. Follow specific use directions and restrictions for these combination tank mixes. Application should be made within one day after planting. Later applications may result in severe crop injury and are not recommended.

WEEDS CONTROLLED PREEMERGENCE

Goal 2XL used alone, at recommended dosages, provides preemergence control of the following broadleaf weeds:

† GROUNDCHERRY, CUTLEAF	POINSETTIA, WILD
JIMSONWEED	SHEPHERDSPURSE
LAMBSQUARTERS, COMMON	SIDA, PRICKLY (TEAWEED)
† NIGHTSHADE, AMERICAN BLACK	SMARTWEED, PENNSYLVANIA
† NIGHTSHADE, BLACK	† SOWTHISTLE, COMMON
PIGWEED, REDROOT	VELVETLEAF

† Suppression of this weed occurs when Goal 2XL is applied at the reduced rate recommended for the GOAL 2XL/metribuzin tank mix combination.

WEEDS CONTROLLED POSTEMERGENCE (POST-DIRECTED APPLICATION)

When Goal 2XL is applied as a post-direct application at the recommended weed stage and dosage in soybeans, the following weeds are controlled:

COCKLEBUR, COMMON	NIGHTSHADE, HAIRY
CROTON, TROPIC	PIGWEED, REDROOT
GROUNDCHERRY, CUTLEAF	† POINSETTIA, WILD
GROUNDCHERRY, WRIGHT	PURSLANE, COMMON
JIMSONWEED	SESBANIA, HEMP
LAMBSQUARTERS, COMMON	SHEPHERDSPURSE
MORNINGGLORY, ANNUAL	†† SICKLEPOD
(UP TO 6 LEAF)	† SIDA, PRICKLY (TEAWEED)
MUSTARD, WILD	smartweed, PENNSYLVANIA
NIGHTSHADE, AMERICAN BLACK	VELVETLEAF
NIGHTSHADE, BLACK	

† Multiple applications may be required for acceptable control.

†† Post-direct applications of Goal 2XL will kill or suppress seedlings not exceeding the one true leaf stage.

Two pints of Latron AG-98, or comparable 80% active nonionic surfactant cleared for application to growing crops, per each 100 gallons of spray solution are suggested in all tank mixtures containing Goal 2XL when postemergence weed control is desired.

TANK MIXES WITH GOAL 2XL

Goal 2XL when applied at 0.6 to 0.8 pint (0.16 to 0.2 lb active) per acre as a tank mix combination with metribuzin (Sencor DF herbicide or Lexone DF herbicide) at 0.33 lb. product (0.25 lb active) per acre, is effective for preemergence control of susceptible broadleaf weeds. Do not apply this tank mix to sandy soils or coarse soils (sandy loam or loamy sand) containing less than 2% organic matter. Do not use on soils with less than 1/2% organic matter, or on alkaline soils with a pH above 7.4 as crop injury may occur. Application should be made within one day following planting. Later applications may result in severe crop injury and are not recommended. The GOAL 2XL/metribuzin herbicide tank mix may be applied as a preemergence application following a preplant incorporated grass herbicide treatment or as a three-way tank mix in a preemergence application with either Dual, Lasso or Surflan.

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

DOSAGE

Refer to the following tables for labeled use rates.

NO-TILL (DOUBLE-CROP) SOYBEANS PREEMERGENCE

Soil Texture	RATE OF PRODUCT PER BROADCAST ACRE (PINTS PER ACRE)						
	GOAL 2XL	Dual 8E †	Lasso 4E †	Surflan AS ††	paraquat (Gramoxone)	glyphosate (Glyphomax)	Bronco †
Coarse	0.5 to 1.5	1.5	4.0 to 5.0	1.5	1.0 to 2.0	2.0 to 3.0	6.5 to 10.0
Medium	0.5 to 2.0	2.0	5.0 to 6.0	2.0	1.0 to 2.0	2.0 to 3.0	8.0 to 10.0
Fine	0.5 to 2.0	2.0 to 2.5	5.0 to 6.0	3.0	1.0 to 2.0	2.0 to 3.0	8.0 to 10.0
Muck or Peat	†††	†††	†††	†††	†††	†††	†††

† Use the higher rate of Bronco, Dual or Lasso on soils containing more than 3% organic matter.

†† When using Surflan 75 WP, multiply pints by 0.67 to obtain the amount of Surflan 75WP product required. Do not use Surflan on soils containing more than 5% organic matter.

††† Do not use.

CONVENTIONAL TILLED SOYBEANS PREEMERGENCE

Goal 2XL is effective for preemergence control of susceptible broadleaf weeds when applied at 1 to 5 pints (0.25 to 0.38 lb active) per broadcast acre. Application should be made within one day of planting. Later applications may result in severe crop injury and are not recommended. The higher rate (0.38 lb active) will assist in early season annual grass control. However, Goal 2XL must not be a basic portion of the grass herbicide program. Goal 2XL may be applied alone as a preemergence application following a preplant incorporated grass herbicide treatment or as a tank mix in a preemergence application with Dual, Lasso or Surflan.

**CONVENTIONAL TILLED SOYBEANS
PREEMERGENCE**

RATE OF PRODUCT PER BROADCAST ACRE (PINTS PER ACRE)

Soil Texture	RATE OF PRODUCT PER BROADCAST ACRE (PINTS PER ACRE)					(lb PER ACRE)
	GOAL 2XL	Dual 8E†	Lasso 4E†	Surflan A.S. ††	Metribuzin DF ††††	
Coarse	0.6 to 1.5	1.25 to 1.5	3.0 to 4.0	1.0 to 1.5		0.33
Medium	0.6 to 1.5	1.5 to 2.0	4.0 to 6.0	1.5 to 2.0		0.33
Fine	0.6 to 1.5	2.0 to 2.5	4.0 to 6.0	2.0 to 2.5		0.33
Muck or Peat	†††	†††	†††	†††		†††

- † Use the higher rate of Bronco, Dual or Lasso on soils containing more than 3% organic matter.
- †† When using Surflan 75 WP, multiply pints by 0.67 to obtain the amount of Surflan 75WP product required. Do not use Surflan on soils containing more than 5% organic matter.
- ††† Do not use.
- †††† Sencor DF or Lexone DF.

WEEDS CONTROLLED PREEMERGENCE

When Goal 2XL is tank mixed with Bronco, Dual, Lasso or Surflan and applied preemergence, in addition to the weeds controlled preemergence by Goal 2XL alone, control of the following weeds is also obtained:

- | | |
|------------------|------------------------|
| BARNYARDGRASS | JOHNSONGRASS, SEEDLING |
| CRABGRASS, LARGE | PANICUM, FALL |
| FOXTAIL, GIANT | RAGWEED, COMMON |
| FOXTAIL, YELLOW | SIGNALGRASS, BROADLEAF |

WEEDS CONTROLLED POSTEMERGENCE

When Goal 2XL is tank mixed with Bronco, paraquat (Gramoxone) or glyphosate (Roundup-Glyphomax) and applied postemergence, in addition to the weeds controlled postemergence by Goal 2XL alone, control of the following weeds is also obtained:

- | | |
|-------------------|-----------------------|
| BLUEGRASS, ANNUAL | FOXTAIL, YELLOW |
| CRABGRASS, LARGE | LAMBSQUARTERS, COMMON |
| FOXTAIL, GIANT | RAGWEED, COMMON |
| FOXTAIL, GREEN | SANDBUR, FIELD |

TIMING AND METHOD OF APPLICATION

As a preemergence treatment, apply in 20 to 60 gallons of water per acre. If Bronco or glyphosate (Glyphomax) are included in the tank mix, apply in 20 to 40 gallons of water per acre. To insure complete coverage, spray volume should be increased as the density of emerged weeds, crop residue or stubble increases. Use conventional spray equipment with flat fan or flood jet nozzles. Spray equipment should be calibrated carefully before each use.

POST-DIRECTED SPRAY

**GOAL 2XL USED ALONE
DOSAGE**

Goal 2XL is recommended as a post-directed application at 1 pint (0.25 lb active) per acre. Optimum control is achieved when Goal 2XL is applied to seedling weeds not exceeding 4 true leaves. See MIXING DIRECTIONS for surfactant recommendations. Weeds should be in the seedling stage, young and actively growing. Do not count cotyledon leaves.

TANK MIXES WITH GOAL 2XL

For improved broadleaf weed control, a tank mixture of Goal 2XL plus Butoxone herbicide or Butyrac 200 herbicide is suggested. Use 1 pint of Goal 2XL (0.25 lb active) with 1 pint of Butoxone (0.22 lb active) or

0.7 to 0.9 pint of Butyrac 200 (0.175 to 0.22 lb active) per broadcast acre. See MIXING DIRECTIONS for surfactant recommendations. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

TIMING

Soybean plant height must be a minimum 8 inches or greater. Use branch lifters or shields if excessive spray contact to the soybean plant cannot be avoided.

METHOD OF APPLICATION

Accurate, uniform placement of Goal 2XL spray is essential for effective weed control and to minimize soybean injury. As a directed postemergence application, Goal 2XL should be applied at 20 to 25 psi using 20 to 40 gallons of spray on a broadcast acre basis. Do not exceed 25 psi. Spray should be directed towards the base of the soybean plant. Soybean foliage receiving accidental spray or drift may be injured. Weeds should be in the seedling stage, young and actively growing.

Goal 2XL can be applied using a post-direct spray rig with only 2 flat fan nozzles per row, 1 nozzle on each side of the row. Additional care should be taken when adjusting the sprayer prior to application. For best coverage, it is suggested to use 4 flat fan nozzles per row, 2 nozzles on each side of the row. The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and downward. With either sprayer system, nozzles should be adjusted to cover the weed foliage with minimum contact to the soybean plant. **Do not use cone nozzles.**

TANK MIXTURE OF GOAL 2XL WITH COMMAND 6EC HERBICIDE SOYBEANS (NOT FOR USE IN CALIFORNIA)

Goal 2XL when applied preemergence at 0.6 to 0.8 pint (0.16 to 0.2 lb active) per acre in a tank mix combination with Command 6EC (EPA Reg. No. 279-3054) at 1 to 1 $\frac{2}{3}$ pints (0.75 to 1.25 lb active) is effective for the control of susceptible annual grass and broadleaf weeds in soybeans. Application should be made within one day following planting. Later applications may result in severe crop injury and are not recommended.

WEEDS CONTROLLED PREEMERGENCE

A tank mix of Goal 2XL with Command 6EC at recommended dosages, provides preemergence control of the following weeds:

GRASS WEEDS

BARNYARDGRASS
CRABGRASS
 (CRABGRASS, LARGE)
 (CRABGRASS, SMOOTH)
CUPGRASS, SOUTHWEST
CUPGRASS, WOOLLY
FOXTAIL
 (FOXTAIL, GIANT)
 (FOXTAIL, GREEN)
 (FOXTAIL, ROBUST PURPLE)
 (FOXTAIL, YELLOW)
GOOSEGRASS
JOHNSONGRASS (SEEDLING)
PANICUM
 (PANICUM, FALL)
 (PANICUM, TEXAS)
SANDBUR, FIELD
SIGNALGRASS, BROADLEAF
 (BRACHIARIA)

BROADLEAF WEEDS

BEGGARWEED, FLORIDA
CROTON, TROPIC
† GROUNDCHERRY, CUTLEAF
JIMSONWEED
LAMBSQUARTERS
MALLOW, VENICE
† NIGHTSHADE, BLACK
PIGWEEED, REDROOT
PURSLANE, COMMON
PUSLEY, FLORIDA
SHEPHERDSPURSE
SIDA, PRICKLY
SMARTWEED, PENNSYLVANIA
† SOWTHISTLE, COMMON
VELVETLEAF

59 7 67

† Suppression.

SOYBEANS - SPECIFIC ENVIRONMENTAL HAZARDS

This product is highly toxic to freshwater clams, oysters, aquatic invertebrates and aquatic plants. Do not apply Goal 2XL where visible erosion to aquatic habitats and/or wetlands occurs. (See container for further information on Environmental Hazards).

SOYBEANS - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not make more than two applications of Goal 2XL per growing season.
- Do not apply more than 2 pints (0.5 lbs active) of Goal 2XL per acre during one growing season as a result of preemergence application in no-till (double-crop) or conventional till soybeans, or post-directed in conventional till soybeans. If early preplant application is made, do not apply more than 3 pints (0.75 lb active) of Goal 2XL per acre during one growing season.
- Do not apply a post-directed application of Goal 2XL to soybeans after the initial appearance of blooms.

TARO

FOR USE IN HAWAII ONLY

GENERAL INFORMATION

GOAL 2XL is a selective herbicide for preemergence and post-direct application to dryland taro for the control of certain broadleaf weeds.

NOTE: Dryland taro is defined as a taro grown without irrigation, or by using irrigation practices that do not result in runoff, irrigation return flow, or other loss of irrigation water from the production area. If irrigation is used, the water applied shall not exceed the field capacity of the soil.

Occasionally, after the use of Goal 2XL, a spotting, crinkling or flecking may appear on the leaves of the taro. Leaves that receive direct or indirect (drift) spray contact will be injured.

Do not use Goal 2XL on taro plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.

DOSAGE

Apply Goal 2XL at a rate of 2 pints (0.5 lb active) per broadcast acre as a single preemergence application within one week after transplanting (and prior to emergence) of the taro.

GOAL 2XL is also recommended as a post-direct application of 1 pint (0.25 lb active) per acre. Effective control of succulent weed seedlings in the 2-to 3-leaf stage can usually be obtained. Do not apply more than 1 pint (0.25 lb active) of GOAL 2XL per acre in a single post-direct application, or more than 2 pints (0.5 lb active) per broadcast acre per season as a result of multiple post-direct applications.

Dosages listed are for broadcast application. For banded application, the amount of Goal 2XL used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

WEEDS CONTROLLED

Goal 2XL will provide preemergence and postemergence control of the following weeds when used at the recommended dosage. Applications to weeds beyond the 3-leaf stage may result in partial control:

AMARANTH, SPINY
PURSLANE, COMMON

SPURGE, GARDEN

TIMING AND METHOD OF APPLICATION

Goal 2XL should be thoroughly mixed with clean water at recommended concentrations, and applied in a minimum of 15 gallons of water per acre.

When applied preemergence, use conventional ground spray equipment with flat fan nozzles at 20 to 40 psi. Accurately calibrate spray equipment prior to each use.

When applied as a post-direct spray, sprays must be directed to the base of the taro plant. Accurate, uniform placement of Goal 2XL is essential for effective weed control and to minimize crop injury. Taro foliage receiving accidental spray or drift will be injured. Goal 2XL must be applied using rigid precision ground sprayer equipment. As a directed postemergence application, Goal 2XL should be applied at 20 to 25 psi using 20 to 40 gallons of spray on a broadcast acre basis. Do not exceed 25 psi.

TARO - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 2 pints (0.5 lb active) of Goal 2XL per broadcast acre as a single preemergence application.
- Do not apply more than 1 pint (0.25 lb active) of Goal 2XL per broadcast acre in a single post-direct spray or more than 2 pints (0.5 lb active) per broadcast acre per season as a result of multiple post-direct applications.
- Do not apply more than 4 pints (1.0 lb. active) of Goal 2XL per broadcast acre per season as a result of preemergence and post-direct applications.
- Do not apply Goal 2XL within 6 months of harvest of taro (corms, leaves).
- For use only to dryland taro grown in Hawaii. (Dryland taro is defined as taro grown without irrigation, or by using irrigation practices that do not result in run-off, irrigation return flow, or other loss of irrigation water from the production area. If irrigation is used, the water applied shall not exceed the field capacity of the soil.)

TREEFRUIT/NUT/VINE**DORMANT APPLICATION**

ALMOND, APPLE, APRICOT, AVOCADO, BEECHNUT, BRAZIL NUT, BUTTERNUT, CASHEW, CHERRY, CHESTNUT, CHINQUAPIN, CRAB APPLE, DATE, FEIJOA, FIG, FILBERT, GRAPES, HICKORY NUT, KIWI, LOQUAT, MACADAMIA NUT, MAYHAW, NECTARINE, OLIVES, PEACH, PEAR, PECAN, PERSIMMON, PISTACHIO, PLUM, POMEGRANATES, PRUNE, QUINCE, WALNUT

GENERAL INFORMATION

GOAL 2XL is effective as a preemergence and/or postemergence herbicide when used alone or in recommended combinations, for the control of certain annual broadleaf weeds, in certain bearing and nonbearing treefruit, nut or vine plantings.

The most effective postemergence weed control is achieved when Goal 2XL is applied to seedling weeds. For postemergence control of certain grassy and broadleaf weeds, a tank mixture of Goal 2XL with either paraquat (Gramoxone) or glyphosate (Glyphomax) can be used.

For preemergence control of susceptible grassy and broadleaf weeds in certain treefruit, nut or vine plantings, a tank mixture of Goal 2XL with napropamide (Devrinol), diuron (Karmex), pronamide (Kerb* herbicide), simazine, norflurazon (Solicam) or oryzalin (Surflan) can be applied. Contact herbicides such as paraquat (Gramoxone) or glyphosate (Glyphomax) may also be added to the tank mixture. Check individual product labels to determine suitability and use rates for various crops.

**GOAL 2XL USED ALONE
GEOGRAPHIC USE DIRECTIONS
ARIZONA AND CALIFORNIA
DOSAGE**

Goal 2XL is recommended for postemergence control at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre. For preemergence control of susceptible weeds, use 5 to 8 pints (1.25 to 2.0 lb active) per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE (weeds up to 4 inches high)—Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. Applications to weeds beyond this 4-inch stage may result in partial control.

CHEESEWEED (MALVA)	MINERSLETTUCE
FIDDLENECK, COAST	NETTLE, BURNING
† FILAREE, BROADLEAF	PIGWEEED, REDROOT
† FILAREE, REDSTEM	REDMAIDS
† FILAREE, WHITESTEM	SHEPHERDSPURSE
GROUNDSEL, COMMON	SOWTHISTLE, ANNUAL
HENBIT	

† Goal 2XL at the 8-pint rate (2.0 lb active) will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

WEEDS CONTROLLED PREEMERGENCE—Apply 5 to 8 pints (1.25 to 2.0 lb active) of Goal 2XL per broadcast acre.

BURCLOVER	LAMBSQUARTERS, COMMON
CHEESEWEED (MALVA)	LETTUCE, PRICKLY
FIDDLENECK, COAST	PIGWEEED, REDROOT
FILAREE, BROADLEAF	PURSLANE, COMMON
FILAREE, REDSTEM	REDMAIDS
FILAREE, WHITESTEM	ROCKET, LONDON
GROUNDSEL, COMMON	SHEPHERDSPURSE
HENBIT	SOWTHISTLE, ANNUAL
KNOTWEED, PROSTRATE	

ALL OTHER STATES (EXCEPT CALIFORNIA AND ARIZONA)

DOSAGE

Goal 2XL is recommended for postemergence control at 2 to 8 pints (0.5 to 2.0 lb active) per broadcast acre. For preemergence control of susceptible weeds, use 5.0 to 8 pints (1.25 to 2.0 lb active) per broadcast acre.

WEEDS CONTROLLED POSTEMERGENCE—Apply 2 to 8 pints (0.5 to 2.0 lb active) of Goal 2XL per broadcast acre. The lower rate is recommended for the control of susceptible seedling weeds in the early postemergence stage up to the 4-leaf stage. The higher rate (2.0 lb active) should be used for weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in partial control.

- | | |
|-----------------------------|-------------------------|
| BALSAMAPPLE | PEPPERWEED, VIRGINIA |
| COCKLEBUR, COMMON | PIGWEEED, REDROOT |
| † CUDWEED, NARROWLEAF | POINSETTIA, WILD |
| †† EVENINGPRIMROSE, CUTLEAF | PURSLANE, COMMON |
| GROUNDCHERRY, CUTLEAF | SESBANIA, HEMP |
| GROUNDCHERRY, WRIGHT | SHEPHERDSPURSE |
| JIMSONWEED | SIDA, PRICKLY (TEAWEED) |
| LAMBSQUARTERS, COMMON | SMARTWEED, PENNSYLVANIA |
| MORNINGGLORY, ANNUAL | SOWTHISTLE, ANNUAL |
| NIGHTSHADE, AMERICAN BLACK | VELVETLEAF |
| NIGHTSHADE, BLACK | |

† Maximum 0.5-inch diameter

†† Highest rate and/or multiple applications may be required for acceptable control. Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in one season.

WEEDS CONTROLLED PREEMERGENCE—Apply 5 to 8 pints (1.25 to 2.0 lb active) of Goal 2XL per broadcast acre.

- | | |
|----------------------------|-------------------------|
| CAMPHORWEED | PIGWEEED, REDROOT |
| CUDWEED, NARROWLEAF | POINSETTIA, WILD |
| † EVENINGPRIMROSE, CUTLEAF | SIDA, PRICKLY |
| GROUNDCHERRY, CUTLEAF | SMARTWEED, PENNSYLVANIA |
| JIMSONWEED | SOWTHISTLE, ANNUAL |
| LAMBSQUARTERS, COMMON | SPURGE, PROSTRATE |
| NIGHTSHADE, AMERICAN BLACK | SPURGE, SPOTTED |
| NIGHTSHADE, BLACK | VELVETLEAF |
| PEPPERWEED, VIRGINIA | |

† Highest rate and/or multiple applications may be required for acceptable control. Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in one season.

ALL STATES

TIMING AND METHOD OF APPLICATION

In Arizona and California, GOAL 2XL can be applied during the period following completion of final harvest up to February 15 (February 1st in the Coachella Valley, California). Applications made after the calendar dates above, but prior to bud swell, may result in significant crop injury and are the responsibility of the user.

In all states, do not apply Goal 2XL after buds start to swell until completion of final harvest. Do not apply when fruit/nuts are present. Goal 2XL can be applied upon completion of final harvest.

As a preemergence treatment, apply a minimum of 40 gallons of water per acre. Use higher volumes to ensure adequate coverage in high densities of emerged weeds or heavy trash. Best preemergence results are achieved when spray is applied to a relatively weed-free established berm or soil surface. Goal 2XL should be directed to the soil and the base of dormant trees or vines. Use a low-pressure sprayer equipped with a breakaway boom and flat fan nozzles. An off-center (OC) nozzle positioned at the end of the boom may be desired. See SPECIFIC USE RESTRICTIONS for Goal 2XL application on dormant tree or vine plantings.

SPRAY VOLUME:

Weed Stage	Gallons of Water per Acre
Preemergence	40 or more
Postemergence (up to 4-inch or 4-leaf stage)	40 or more
Exceeding 4-inch or 4-leaf stage	100 or more

TANK MIXES WITH GOAL 2XL

IMPORTANT: Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

DOSAGE

For preemergence control of susceptible grassy and broadleaf weeds in certain bearing and nonbearing treefruit, nut or vine plantings, a tank mixture of Goal 2XL with napropamide (Devrinol), diuron (Karmex), pronamide (Kerb), simazine, norflurazon (Solicam) or oryzalin (Surflan) can be applied. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

For postemergence control of susceptible grassy and broadleaf weeds in certain treefruit, nut or vine plantings, a tank mixture of paraquat (Gramoxone) or glyphosate (Glyphomax) with Goal 2XL or combinations of Goal 2XL plus napropamide (Devrinol), diuron (Karmex), pronamide (Kerb), simazine, norflurazon (Solicam) or oryzalin (Surflan) with either paraquat (Gramoxone) or glyphosate (Glyphomax) can be used. Apply at the recommended rates and growth stages to susceptible weed species in a manner described on the respective labels.

WEEDS CONTROLLED

In addition to the weeds controlled by Goal 2XL used alone, control of susceptible weeds listed on the respective labels for the following products is also obtained:

diuron (Karmex)	oryzalin (Surflan)
glyphosate (Glyphomax)	paraquat (Gramoxone)
napropamide (Devrinol)	pronamide (Kerb)
norflurazon (Solicam)	† simazine

† In addition, simazine provides preemergence control of horseweed (marestail).

CHEMIGATION (ALL STATES): For application using sprinkler (low-volume (microsprinkler)), drip (trickle), and flood (basin) irrigation systems, apply specified dosage of GOAL 2XL per acre as described in DOSAGE section above. Follow all directions given in the CHEMIGATION section of this label when making applications using irrigation systems. Do not allow treated irrigation water to contact the fruit or foliage.

**TREEFRUIT/NUT/VINE
SPECIFIC USE RESTRICTIONS**

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply Goal 2XL during the period between bud swell and completion of final harvest or when fruit/nuts are present. Goal 2XL can be applied upon completion of final harvest.
- **IN ARIZONA AND CALIFORNIA, GOAL 2XL can be applied during the period following completion of final harvest up to February 15 (February 1st in the Coachella Valley, California).**

64 / 67

Applications made after the calendar dates above, but prior to bud swell, may result in significant crop injury and are the responsibility of the user.

- Do not apply more than 8 pints (2.0 lb active) per broadcast acre of Goal 2XL in one season.
- Do not apply to grapes or kiwi established less than 3 years unless vines are on a trellis wire a minimum of 3 feet above the soil surface.
- Do not apply to grapes or kiwi that are not staked or trellised unless vines are free-standing.
- Goal 2XL or any of the combinations recommended on this label should be applied to only healthy growing trees or vines.
- Direct spray toward the base of tree or vines. Avoid direct plant contact.

SPECIFIC USE RESTRICTIONS FOR INDIVIDUAL CROPS ARE FOUND UNDER DIRECTIONS FOR USE IN EACH CROP GROUP SECTION.

GENERAL USE RESTRICTIONS: USE RESTRICTIONS THAT APPLY TO ALL REGISTERED APPLICATIONS ARE AS FOLLOWS:

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not contaminate irrigation water or water used for domestic purposes.
- Do not use any plants treated with Goal 2XL for feed or forage.
- Do not feed or allow animals to graze on any areas treated with Goal 2XL.
- Goal 2XL should be applied only by ground application equipment except as specifically directed on this label or on other approved Rohm and Haas Company Supplemental Labeling.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. Goal 2XL is phytotoxic to plant foliage.
- Thoroughly flush spray equipment (tank, pump, hoses and boom) with clean water before and after each use. Residual Goal 2XL remaining in spray equipment may damage other crops. To assist removal of Goal 2XL residues in spray equipment, Latron AG-98 or Latron CS-7 may be added at the rate of 1 quart per 100 gallons of water during flushing.
- Use Goal 2XL only for recommended purposes and at recommended rates.
- Do not treat ditch banks or waterways with Goal 2XL.

•ROTATION CROP RESTRICTIONS

Do not rotate to small-grain crops (includes barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, wild rice) within 10 months following a GOAL treatment.

- Do not direct seed any crops, other than GOAL-labeled crops, within 60 days following a GOAL treatment.
- Do not transplant seedling crops, other than GOAL-labeled crops, within 30 days following a GOAL treatment. **IMPORTANT: TREATED SOIL MUST BE THOROUGHLY INCORPORATED TO A DEPTH OF 4 INCHES AFTER HARVEST (OR ABANDONING) OF THE TREATED CROP BUT PRIOR TO PLANTING OF THE ROTATIONAL CROP. FAILURE TO ACHIEVE THIS THOROUGH AND COMPLETE INCORPORATION OR TO FOLLOW THE REQUIRED MINIMUM PLANT-BACK INTERVAL MAY RESULT IN CROP INJURY, STAND REDUCTION AND/OR VIGOR REDUCTION OF THE PLANT-BACK CROP. See specific fallow bed labeling regarding crop planting information for applications of GOAL 2XL made to a fallow bed or fallow field.**

WEEDS LISTED

AGERATUM	<i>Ageratum conyzoides</i>
AMARANTH, SPINY	<i>Amaranthus spinosus</i>
BALSAMAPPLE	<i>Momordica charantia</i>
BARNYARDGRASS (WATERGRASS)	<i>Echinochloa crus-galli</i>
BEDSTRAW, CATCHWEED	<i>Galium aparine</i>
BITTERCRESS, LESSER	<i>Cardamine oligosperma</i>
BLUEGRASS, ANNUAL	<i>Poa annua</i>
BUCKWHEAT, WILD	<i>Polygonum convolvulus</i>
BURCLOVER	<i>Medicago hispida</i>
BUTTERCUP, SMALLFLOWER	<i>Ranunculus abortivus</i>
BUTTONWEED	<i>Borreria laevis</i>
CAMPHORWEED	<i>Heterotheca subaxillaris</i>
CANARYGRASS (ANNUAL)	<i>Phalaris canariensis</i>
CARPETWEED	<i>Mollugo verticillata</i>
CHEESEWEED (MALVA)	<i>Malva parviflora</i>
CLOVER, RED	<i>Trifolium pratense</i>
CLOVER, WHITE	<i>Trifolium repens</i>
COCKLEBUR, COMMON	<i>Xanthium pensylvanicum</i>
CRABGRASS, LARGE (HAIRY)	<i>Digitaria sanguinalis</i>
CROTALARIA	<i>Crotalaria species</i>
CROTON, TROPIC	<i>Croton glandulosus</i>
CUDWEED, NARROWLEAF	<i>Gnaphalium falcatum</i>
EVENINGPRIMROSE, CUTLEAF	<i>Oenothera laciniata</i>
FIDDLENECK, COAST	<i>Amsinckia intermedia</i>
FILAREE, BROADLEAF	<i>Erodium botrys</i>
FILAREE, REDSTEM	<i>Erodium cicutarium</i>
FILAREE, WHITESTEM	<i>Erodium moschatum</i>
FIREWEED (FROM SEED)	<i>Epilobium angustifolium</i>
FLIXWEED	<i>Descurainia sophia</i>
FOXTAIL, GIANT	<i>Setaria faberi</i>
FOXTAIL, GREEN	<i>Setaria viridis</i>
FOXTAIL, YELLOW	<i>Setaria lutescens</i>
GERANIUM, CAROLINA	<i>Geranium carolinianum</i>
GOOSEGRASS	<i>Eleusine indica</i>
GROUNDCHERRY, CUTLEAF	<i>Physalis angulata</i>
GROUNDCHERRY, WRIGHT	<i>Physalis wrightii</i>
GROUNDSEL, COMMON	<i>Senecio vulgaris</i>
HENBIT	<i>Lamium amplexicaule</i>
HORSEWEED (MARESTAIL)	<i>Conyza canadensis</i>
JIMSONWEED	<i>Datura stramonium</i>
JOHNSONGRASS, SEEDLING	<i>Sorghum halepense</i>
KNOTWEED, PROSTRATE	<i>Polygonum aviculare</i>
LADYSTHUMB (SMARTWEED)	<i>Polygonum persicaria</i>
LAMBSQUARTERS, COMMON	<i>Chenopodium album</i>
LETTUCE, PRICKLY (CHINA LETTUCE)	<i>Lactuca serriola</i>
MALLOW, LITTLE (MALVA)	<i>Malva parviflora</i>
MAYWEED (DOG FENNEL)	<i>Anthemis cotula</i>
MINERSLETTUCE	<i>Montia perfoliata</i>
MORNINGGLORY SPECIES, ANNUAL	<i>Ipomoea species</i>
MORNINGGLORY, IVYLEAF	<i>Ipomoea hederacea</i>

MORNINGGLORY, TALL	<i>Ipomoea purpurea</i>
MUSTARD, BLACK	<i>Brassica nigra</i>
MUSTARD, BLUE (PURPLE MUSTARD)	<i>Chorispora tenella</i>
MUSTARD, COMMON YELLOW	<i>Brassica campestris</i>
MUSTARD, HEDGE	<i>Sisymbrium officinale</i>
MUSTARD, TUMBLE (JIM HILL MUSTARD)	<i>Sisymbrium altissimum</i>
MUSTARD, WILD	<i>Brassica kaber</i>
NETTLE, BURNING	<i>Urtica urens</i>
NIGHTSHADE, AMERICAN BLACK	<i>Solanum americanum</i>
NIGHTSHADE, BLACK	<i>Solanum nigrum</i>
NIGHTSHADE, HAIRY	<i>Solanum sarrachoides</i>
OATS, WILD	<i>Avena fatua</i>
ORACH, RED	<i>Atriplex rosea</i>
OXALIS (BERMUDA BUTTERCUP)	<i>Oxalis pes-caprae</i>
PANICUM, FALL	<i>Panicum dichotomiflorum</i>
PEPPERWEED, VIRGINIA	<i>Lepidium virginicum</i>
PEPPERWEED, YELLOWFLOWER	<i>Lepidium perfoliatum</i>
PIGWEEED, PROSTRATE	<i>Amaranthus blitoides</i>
PIGWEEED, REDROOT	<i>Amaranthus retroflexus</i>
PIMPERNEL, SCARLET	<i>Anagallis arvensis</i>
POINSETTIA, WILD	<i>Euphorbia heterophylla</i>
PUNCTUREVINE	<i>Tribulus terrestris</i>
PURSLANE, COMMON	<i>Portulaca oleracea</i>
PUSLEY, FLORIDA	<i>Richardia scabra</i>
RAGWEED, COMMON	<i>Ambrosia artemisiifolia</i>
REDMAIDS	<i>Calandrinia caulescens</i>
ROCKET, LONDON	<i>Sisymbrium irio</i>
RYEGRASS, ITALIAN	<i>Lolium multiflorum</i>
SAGE, LANCELEAF	<i>Salvia reflexa</i>
SANDBUR, FIELD	<i>Cenchrus incertus</i>
SANDSPURRY, RED	<i>Spergularia rubra</i>
SESBANIA, HEMP	<i>Sesbania exaltata</i>
SHEPHERDSPURSE	<i>Capsella bursa-pastoris</i>
SICKLEPOD	<i>Cassia obtusifolia</i>
SIDA, PRICKLY (TEAWEED)	<i>Sida spinosa</i>
SIGNALGRASS, BROADLEAF	<i>Brachiaria platyphylla</i>
SMARTWEED, PENNSYLVANIA	<i>Polygonum pensylvanicum</i>
SORREL, RED (FROM SEED)	<i>Rumex acetosella</i>
SOWTHISTLE, ANNUAL	<i>Sonchus oleraceus</i>
SPEEDWELL, BIRDSEYE	<i>Veronica persica</i>
SPURGE, GARDEN	<i>Euphorbia hirta</i>
SPURGE, PROSTRATE	<i>Euphorbia supina</i>
SPURGE, SPOTTED	<i>Euphorbia maculata</i>
SPURRY, CORN	<i>Spergula arvensis</i>
TANSYMUSTARD	<i>Descurainia pinnata</i>
THISTLE, BULL	<i>Cirsium vulgare</i>
THISTLE, RUSSIAN	<i>Salsola kali</i>
VELVETLEAF	<i>Abutilon theophrasti</i>
WITCHGRASS	<i>Panicum capillare</i>
WITCHWEED	<i>Striga asiatica</i>
WOODSORREL, COMMON YELLOW	<i>Oxalis stricta</i>

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC
EPA-accepted date: __/__/__