8/1/2003

1814

Page

.

# Dimension\* 0.25G

62719-429

G2B / Dimension 0.25G / Amend / 06-23-03

**Turf and Ornamental Herbicide** 

For season-long control or suppression of listed annual grasses and broadle af weeds in ornamental landscape areas, established lawns and ornamental turit, including golf course fairways, roughs and tee boxes.

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

In the state of New York, this product may be applied only by commercial applicators at no more than 185 lb (0.5 lb active ingredient) per acre (4.25 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Active Ingredient:

This product contains 0.135 pounds of the active ingredient, dithiopyr, per 50 pound bag. Product protected by U.S. Patent No. 4,692,184. Other patents pending

# Keep Out of Reach of Children

See back or container/side panel for additional Precautionary Statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.


Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-429

EPA Est. No. \_\_\_\_\_

Net Weight \_\_ Ib

\*Trademark of Dow AgroSciences LLC Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA



F

# 

# CAUTION: Causes Eye Irritation

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling:

# **First Aid**

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency medical treatment information.

# **Environmental Hazards**

This product is toxic to fish. Drift and runoff may adversely affect aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters

# **Directions for Use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

# **Storage and Disposal**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Storage:** Store this product only in its original container only. In case of spill, sweep up material and dispose of material according to label directions.

**Pesticide Disposal:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

**Container Disposal:** Do not reuse emptied container. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or, if allowed by state and local authorities. by burning. If burned, stay out of smoke.

# Use Directions for Turf

Dimension\* 0.25% turf and ornamental herbicide provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

# **Use Precautions for Turf**

- Use of this product on golf greens is not recommended.
- Do not apply this product to Colonial bentgrass.
- For best results, cultural practices that disturb the soil, such as core, spike, hydro-aerification, or verticutting, and topdressing should be done before application of this product. This product should not be applied until turfgrasses have recovered from stresses related to these or other mechanical

operations. Irrigation or rainfall within 7 days after application will improve the performance of this product.

- Reseeding or Sprigging: Reseeding, overseeding, or sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 12 weeks from the time of application. If overseeding with perennial ryegrass, delay overseeding at least 8 weeks after application. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact.
- This product may be used on seeded, sodded, or sprigged turfgrass that is well established.\* Allow the turf to develop a good root system and uniform stand before application. Use of this product on turfgrass not well established, weakened by lack of moisture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to recover before application of this product.
- This product may be applied as a single application or as a split application in the spring, summer or fall.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow domestic animals to feed or forage on the treated turf or clippings from treated turf.

When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

# **Cool-Season Grasses**

bentgrass, creeping <sup>T</sup>	Agrostis stolonifera
bluegrass, Kentucky	Poa pratensis
fescue, fine tt	Festuca rubra
fescue, tall	Festuca arundinacea
ryegrass, perennial	Lolium perenne

#### Warm-Season Grasses

bahiagrass	Paspalum notatum
bermudagrass	Cynodon dactylon
buffalograss	Buchloe dactyloides
carpetgrass	Axonopus affinis
centipedegrass	Eremochloa ophiuroides
kikuyugrass	Pennisetum clandestínum
St. Augustinegrass	Stenotaphrum secundatum
zoysiagrass	Zoysia japonica

<sup>†</sup> Do not apply this product to Colonial bentgrass. Use of this product on certain varieties of creeping bentgrass, such as 'Carmen', 'Cohansey', or 'Seaside' will result in undesirable turf injury. Not all varieties of creeping bentgrass have been tested.

<sup>††</sup> The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'.

# **Application Equipment and Instructions**

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage, and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction of application. Avoid the use of spreaders that apply granular materials in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

#### Preemergence Control of Annual Grasses and Annual Broadleaf Weeds

# G2B / Dimension 0.25G / Amend / 06-23-03

Used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for "..." crabgrass in early stages of development, but only if application is made to crabgrass prior to the 3-leaf stage of growth or first tiller. Use the higher recommended use rates where weed infectations are dense and/or applications are made early and longer residual control is desirable.

#### Weeds Controlled and Use Rate Recommendations

Weeds Controlled		Suggested Use I	Rate Per Application.
		Pounds/Acre	Pounds/1000 sq ft
crabgrass, large crabgrass, smooth crabgrass, southern	foxtail, green foxtail, yellow	100 to 150 <sup>tr</sup>	2.3 to 3.4 <sup>+</sup>
barnyardgrass chickweed clover, hop crabgrass, tropical crowfootgrass dichondra, Carolina lespedeza, common kikuyugrass oxalis, buttercup oxalis (creeping woodsorrel)	oxalis (yellow woodsorrel) pineappleweed ryegrass, annual ryegrass, perennial smutgrass speedwell, corn spurge, prostrate spurge, spotted	150 to 185 T	3.4 to 4.25
bluegrass, annual bittercress chickweed geranium, Carolina goosegrass	henbit parsley-piert pineappleweed purslane, common shepard's-purse	185 **	4.25 **

<sup>1</sup> Higher use rates are recommended where weed infestations are heavy and/or dense.

<sup>11</sup> Use of this product at this rate will provide only partial control and/or suppression of listed weeds.

# **Restrictions:**

- Do not apply more than 185 pounds of this product per acre per application and no more than 3 applications per year (555 pounds) of this product per acre per year.
- In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 lb/1000 sq ft), (0.5 lb active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

# Use Directions for Ornamental Plantings

Dimension 0.25 Turf and Ornamental Herbicide provides preemergence control of listed annual grass and broadleaf weeds in ornamental plantings. Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

4 7 14

Page 4

# Use Rates

Refer to the following table for treatment dosage for treating small areas.

				•		:	۰.
	Ounces	Ounces (Grams) :		:	•_	•	• . •
Pounds Product/Acre	Product/1000 Square Feet	Product/100 Square Feet	ľ	•	•		•
185	68	6.8 (193 grams)					

# **Application Timing and Use Recommendations**

This product is a preemergence herbicide that will not control established weeds. Make applications prior • to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil/media surface as the herbicide barrier will be broken.

#### Weeds Controlled

Used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for crabgrass in early stages of development This product will not control established weeds, except for crabgrass in early stages of development, but only if application is made to crabgrass prior to the 3-leaf stage of growth or first tiller. Therefore, the area to be treated should be free of weeds prior to application. Repeat applications may be made during the growing season at 3 to 4 month intervals to maintain weed control.

Annual Grasses	
Common Name	Scientific Name
barley	Hordeum spp.
barnyardgrass	Echinochloa crus-galli
bluegrass, annual	Poa annua
brome	Bromus spp.
crabgrass, large	Digitaria sanguinalis
crabgrass, smooth	Digitaria ischaemum
crabgrass, Southern	Digitaria ciliaris
crowfootgrass	Dactyloctenium aegyptium
dallisgrass (seedling)	Paspalum dilatatum
goosegrass	Eleusíne Indica
foxtail, green	Setaria verdi
foxtail, yellow	Setaria faberi
kikuyugrass	Pennisetum clandestinum
oats, wild	Avena fatua
ryegrass (annual & perennial)	Lolium spp.
smutgrass	Sporobolus indicus
Broadleaf Weeds	
Common Name	Scientific Name
bittercress	Cardamine spp.
carpetweed	Mollugo verticillata
chickweed	<i>Stellaria</i> spp.
geranium, Carolina	Geranium carolinianum
henbit	Lamium spp.
knotweed, prostrate	Polygonum saviculare
lespedeza, common	Lespedeza striata
lambsquarters	Chenopodium spp.
marestail	Hippuris vulgaris
medic, black	Medicago lupulina

#### Weeds controlled or suppressed by Dimension 0.25G when applied at 185 pounds per acre:

Page 5

#### G2B / Dimension 0.25G / Amend / 06-23-03

mustard	Brassica spp.
oxalis, buttercup	Oxalis pes-caprae
pineappleweed	Matricaria matricarioides
pigweed, redroot	Amaranthus retroflexus
parsley-piert	Alchemilla arvensis
purslane, common	Portulaca oleracea
rocket, London	Sisymbrium irio
shepherdspurse	Capsella bursa-pastoris
speedwell, corn	Veronica arvensis
spurge, garden	Euphorbia hirta
spurge, prostrate	Euphorbia humistrata
spurge, spotted	Euphorbia maculata
woodsorrel, creeping	Oxalis corniculata
woodsorrel, yellow	Oxalis stricta

Page 6

14

# **Application Equipment and Instructions**

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction of application.

Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

# **Use Precautions for Ornamental Plantings:**

- Only apply this product to established ornamentals.
- Do not apply this product to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or diseases).
- Do not make applications when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not apply this product on grasses grown for seed.
- · Do not graze livestock or feed forage cut from areas treated with this product.
- Do not apply this product in enclosed structures and greenhouses.
- Do not apply another herbicide within 4 weeks of application of this product.

#### **Restrictions:**

- Do not apply more than 185 pounds of this product per acre per application and no more than 3 applications per year (555 lb) of this product per acre per year.
- In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 ounces/1000 sq ft), (0.5 lb active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

# **Tolerant Ornamentals**

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Under your growing conditions, a limited number of plants should be treated for tolerance evaluations, prior to initiating large-scale applications. Follow directions given above to determine plant tolerance under your growing conditions prior to large-scale use

.

ĉ

Page 7

7714

Common Name	Botanical Name	Tolerant Cultivars	]
Abelia, Dwarf	Abelia X grandiflora	Nana	
Ajuga	Ajuga reptans	Bronze	
	Ajuga genevensis	Bronze Beauty	ه د ا
Almond, Flowering	Prunus gladulosa		
Apple †	Malus pumila	3 * *	•
Arborvitae	Thuja occidentalis	Nigra	
		Pyramidalis	
		Smaragh	
		Techny	ſ
		Woodwardii	
Arborvitae, Golden	Thuja orientalis		
Aster, Chinese	Callistephus chinensis	Dwarf Queen	
Ash, Green	Fraxinus pennsylvanica		
Ash, Mountain	Sorbus aucuparia	l	
Ash, Purple	Fraxinus americana		
Azalea	Rhododendron spp.	Brilliant	
		Buccaneer	
		Carror	
		Chimes (Belgian)	
		Elsie Lee	
		Exbury	
		Fashion	
		Hardijzer Beauty	
		Hershey Red	
		Higasa	
		Hinocrimson	
		Holland (Hybrid)	
		Marion Lee	
		Northern Lights	
		Orange Cup	
		Orchid Lights Snow	
		Southern Charm	
Azalea, Flame	Rhododendron calendulaceum		
Azalea, Kirishima			
Bamboo, Heavenly			
Barberry	Berberís thunbergii	Aurea	
Barberry		Dwarf Pigmy	
		Green	
		Kobold	
		Pygmy Red	
		Rose Glow	
Barberry, Purple		Atropurpurea	
Basket flower	Gaillardia grandiflora		
Beach grass	Ammophila breviligulata		
Bearberry (common)	Arctostaphylos uva-ursi	Massachusetts	
Bee Balm	Monarda didyma		
Begonia	Begonia spp.		
Birch, River	Betula nigra		
Blackeyed Susan	Rudbeckia hirta	Goldstrum	
Blanket Flower	Gaillardia spp.		

.

*.* 

4

8 8 14

Blueberry <del>†</del>	Vaccinium spp.	Bluecrop Blue Jay Jersey North Blue Northland	
Bottlebrush	Callistemon citrinus		
Boxwood, Japanese		Japonica	• • • •
Boxwood, Weller	Buxus sempervirens		
Broom	Cytisus spp.	Moonlight	
	Genista pilosa	Vancouver Gold	
Bugle Carpet			
Camellia	Camellia japonica	Debutante Mathotiana Supreme Chansonette	
	Camellia sasanqua		
Candy Tuft	Iberis spp.	Snow White	
Carex. Variegated	Carex		
Cedar, Red	Juniperus virginiana		
Celosia	Celosia spp.		
Centaura	Centaurea montana	Scarlet Plumosa	
Cockscomb, Plumosa	Celosia cristata	Red Kewpie	
Coleus	Coleus blumei		
Columbine	Aguilegia spp.		
Copper leaf	Acalypha wilkesiana		
Coreopsis	Coreopsis spp.	Moonbeam,	
Corn Flower	Centaurea spp.		
Cotoneaster	Cotoneaster apiculatus		
Coyotebrush	Baccharis pilularis		
Cycads	Cycas revoluta	······································	
Cypress, Bald	Taxodium distichum		
Cypress, Italian	Cupressus sempervirens	Glauca	
Cypress, Japanese False	Chamaecyparis obtusa	Gracilis	
Cypress, Leyland	Cupressocyparis leylandii		
Daffodil	Narcissus spp.	King Alfred	
Daylilly	Hemerocallis spp.	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilsonís Yellow	
Dianthus (Sweet William)	Dianthus spp.		
Delphinium	Delphinium spp.	Magic fountain	
Dogwood	Cornus florida		
Dogwood, American	Cornus sericea	Flavarimaea	
Douglas Fir	Pseudotsuga menziesii	<u> </u>	
Dusty Miller	Senecio cineraria		
Elm, Drake	Ulmus parvifolia		

Page 9

9 7 14

Euonymus	Euonymus fortunei	Argenteo-variegata Auereo-marginata Colorata Emerald Gaiety	
		Emerald Gaiety	
		Emerald ëN Gold	
		Gold Edge	
		Gold Princess	
		Silver King	
		Tricolor	•••
Fan Palm, European	Chamaerops humilis	Vegetus	
Fan Palm, Mexican	Washingtonia robusta		
Fern (various)	Asparagus spp.		
Fescue	Festuca glauca	· · · · · · · · · · · · · · · · · · ·	
Fetterbush	Leucothoe fontanesiana	Rainbow	
Ficus	Ficus retusa	Nitidia	
Fir Fraser	Abies fraseri		
Forsythia	Forsythia X intermedia	Arnold Dwarf	
roisyina	i orsymia x intermedia	Bronxensis Dwarf	
		Lynwood Gold	
		Meadowlark	
	Spring Glory	Weeping	
Fountain grass	Pennisetum setaceum		
Fuchsia	Fuchsia spp.		
Galium	Galium ordoratum		
Gardenia	Gardenia jasminoides	Mystery Radicans	
Geranium	Pelargonium X hortorum		
Gum	Eucalyptus citriodora		
Hawthorn	Crataegus spp.	Cockspur White	
		Crimson Cloud	
		Enchantress	
		Jack Evans	
		Washington White	
Heather, Twisted	Erica cinerea	Mediterranean Pink	
Hemlock, Canada	Tsuga canadensis		
Hibiscus	Hibiscus spp.	Blue Bird	
		Brilliant	
Holly		Hula Girl Blue Boy	
	llex spp.	Blue Girl	
	llex X meserveae	Burfodii	(
	nex x meserveae	China Girl	
		Compacta	
		Forsteri	
		Hellerie	
		Japanese Northern Beauty	1
		Needlepoint	1
		Nellie R. Stevens	
	llex X attenuata	Savannah	
Holly, Chinese	llex cornuta		
Holly, Japanese	llex crenata		
Holly, Yaupon	llex vomitoria		

ĉ

Page 10

G2B / Dimension 0.250	G / Amend / 06-23-03		10 7 14 Page 10
Honeysuckle	Lonicera japonica	Claveyís Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta	
Hosta	Hosta spp. Hosta lancifolia	Albo Marginata	<u></u>
Ice Plant	Carpobrotus edulis		,, ,, ,, ,, ,,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, ,, , ,, , ,, , , , , , , , , , , , , , , , , , , ,
Impatiens	Impatiens spp.		• • • •
lris	Iris spp.	Dwarf Blue Wedgewood	
Ivy, English	Hedera helix	Bulgaria Thorndale	
Jasmine, Asian	Trachelospermum asiaticum		
Juniper	Juniperus spp.	Arcadia Armstrong Bar Harbor	
	Juniperus horizontalis	Blue Chip Blue Pacific Blue Rug	
	Juniperus horizontalis	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fruitlandi Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blue Sargent Blue Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan	
	Juniperus chinensis Juniperus sabina	Tamariscifolia Weberi	
		Youngstown Yukon Belle	
King Palm	Archontophoenix cunninghamiana		
Laurel, Australian	Pittosporum tobira		
Laurel, Mountain	Kalmia latifolia	······································	
Leucothoe	Leucothoe fontanesiana		
Ligustrum, Japanese	Ligustrum japonicum	1	]

.

 $\hat{}$ 

ť

(

Lily, African	Agapanthus africanus	Albus Peter Pan	
Lily, African Blue		Peter Pan	
Lily of the Valley	Pieris japonica	Mt. Fire	
Lilyturf	Liriope muscari	Evergreen Giant Lilac Beauty	· · · · · ·
Liriope, Green	Liriope spicata		
Maple, Japanese	Acer japonicum		
Maple, Norway	Acer platanoides		
Maple, Red	Acer rubrum		
Maple, Silver	Acer saccharinum		
Maple Sugar	Acer saccharum		
Marigold	Tagetes patula	Honeycomb Variegata Wheelerís Dwarf	
Mock Orange †	Philadelphus spp	Golden Snowflake Double White	
Mondo Grass	Ophiopogon japonicus		
Moss Rose	Portulaca grandiflora	Sunnyside	
Myrtle, Crape	Lagerstroemia indica	Faurei Langer Muskogee Standard Pink	
Myrtle, Wax	Myrica cerifer		
Nandina	Nandina domestica	Compacta Nana	
Narcissus	Narcissus spp.		
Oak, Laurel	Quercus laurifolia		
Oak, Pin	Quercus palustris		
Oak, Red	Quercus rubra		
Oak, Southern L	Quercus virginiana		
Oak, Willow	Quercus phellos		
Oleander	Nerium oleander	Hardy Red Petite Pink Sister Agnes	
Osteospermum	Osteospermum fruticosum	Wirligig	
Pachysandra	Pachysandra terminalis		
Palm, Bangalow			
Pampas Grass	Cortaderia selloana		
Pansy	Viola x wittrockiana		
Paper Flower	Bougainvillea glabra	Barbara Karst	
Peach †	Prunus persica		
Periwinkle, Dwarf	Vinca minor		
Petunia	Petunia X hybrida	Picoti	
Photinia, Red Tip	Photinia X fraseri		
Pieris	Pieris japonica		
Pine, Australian	Pinus nigra Pieris Taiwanensis		
	Piens Lawanensis		

.....

11 8 14

;;

G2B / Dimension 0.25G / Amend / 06-23-03

.....

٠

ç

Page 12

Pine, Loblolly	Pinus taeda			1
Pine, Longleaf	Pinus palustris			
Pine, Mugho	Pinus mugho	•	· · · · · ·	
Pine, Scotch	Pinus sylvestris		· · ·	
Pine, Slash	Pinus elliottii			
Pine, Swiss Mt.	Pinus mugo			
Pine, Virginia	Pinus virginiana		•	••••
Pine, White	Pinus strobus			
Pineapple, Guava	Feijoa sellowiana	· · · · · · · · · · · · · · · · · · ·	839	•••
Pittosporum, Japan	- r cijou sonovnana	······································		
Potentilla	Potentilla nepalensis	Abbotswood		
	Potentilla fruticosa			
Privet	Ligustrum japonicum	Golden Vicary Regal Texanum Wax Yellow Tipped		
Privet, Glossy	Ligustrum lucidum			
Pyracantha	Pyracantha spp.	Gnome Lalandei Victory		:
Queen Palm	Arecastrum rammanzoffianum			
Quince, Japanese †	······································			
Rhododendron	Rhododendron spp	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink		
Rhododendron, Carolina	Rhododendron carolinianum			
Rhododendron, Catawba	Rhododendron catawbiense			
Ribbon grass	Phalaris arundinacea			
Rockcress	Arabis spp.	Snowcap		
Rodie Max (Rosebay)	Rhododendron maximum			
Rose	Rosa banksiae	Luta		
Rosemary †	Rosmarinus officinalis			
Rosemary, Bog	Andromeda polifolia	Nana		
Salvia	Salvia farinacea	Rhea		
Sedum	Sedum spurium	Dragon Blood Red Red Carpet Yellow		
Snapdragon	Antirrhinum spp.			
Sourwood	Oxydendrum arboreum			
Spiraea	Astilbe X arendsii	Fanali		

r

13

Spiraea Spiraea spp. Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houtte White Spiraea, Garland Spiraea spp. Spruce, Black Hills Picea glauca Spruce, Colorado Blue Picea pungens Spruce, Norway Picea abies Spruce, White Picea glauca Conica Sweetflag, Grassyy-Le Sweetgum Liquidambar styraciflua Sycamore Platanus occidentalis Tree Fern (Tiki Fern) Asparagus virgatus Trumpet Flower, Gelsemium sempervirens Evening Apeldoorn Tulip Tulip, spp Tufted hairgrass Deschampsia caespitosa Verbena, Shrub Lantana sellowiana Vervain Verbena spp. St. Paul Viburnum Viburnum spp. American Cranberry Bush Arrowood **Common Snowball** European Cranberry Bush Linden Mohican Wright Vinca (Periwinkle) Vinca minor Windmill Palm Trachycarpus fortunei Xylosma Xylosma congestum Yarrow Achillea spp. Yaupon llex vomitoria Dwarf Yew Taxus cuspidata Denisiformis Taxus X media

<sup>†</sup> Ornamental species only: Do not use on food producing trees and ornamentals.

# **Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

# Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

# Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance,

Page 13

Page 14

or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

# **Limitation of Remedies**

The exclusive remedy for losses or damages resulting from this product (including claims based on \* \* contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of \* the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC EPA-accepted \_\_/\_\_/\_\_\_

ł