

62719-442

10-5-2004

1/17

F3B / Intrepid 2F / PropSec 3 Supp Strawberry / 02-11-04

file: R846-016 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Strawberry 11Feb04d.doc

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F for control or suppression of lepidoptera larvae infesting strawberry.

*Trademark of Dow AgroSciences LLC

2/17

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid* 2F

EPA Reg. No. 62719-442

For Control or Suppression of Lepidoptera Larvae Infesting Strawberry

ATTENTION


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Intrepid 2F (fl oz/acre)	Application Timing	Restrictions
armyworms corn earworm cutworms (suppression)	6 – 12 (0.09 – 0.19 lb ai/acre)	For early season applications only to young crops and small plants. Apply at first sign of feeding damage or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. For heavy infestations, continuous moth flights, and/or egg masses and larvae in all stages of development, a 10-14 day re-treatment interval is required to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> • Do not apply more than 12 fl oz per acre per application or 64 fl oz of Intrepid 2F (1 lb ai) per acre per season. • Pre-harvest Interval: Do not harvest within 3 days of application. • See Rotational Crop Restrictions.



Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult to cover crops to ensure thorough coverage. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days. **Note:** These rotational crop restrictions supercede prior restrictions of 356 days for strawberry and the prohibition for all other crops.

Note: This product may be hazardous to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R846-016

EPA-accepted ___/___/___

Initial printing.

F3B / Intrepid 2F / Prop Sec 3 Supp Leaves Root-Tuber Veggies / 01-22-04
file: R846-021 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Leaves Root-Tuber Veggies (Crop Group 2)
22Jan04d.doc

3/11

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F for control of lepidoptera larvae infesting leaves of root and tuber vegetables (crop group 2).

*Trademark of Dow AgroSciences LLC

4/17

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid* 2F

EPA Reg. No. 62719-442

For Control or Suppression of Lepidoptera Larvae Infesting Leaves of Root and Tuber Vegetables (Crop Group 2†)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

†Leaves of root and tuber vegetables (crop group 2) including, but not limited to: Bitter Cassava, Black Salsify, Carrot, Celeriac, Chicory, Dasheen, Edible Burdock, Garden Beet, Parsnip, Oriental Radish, Radish, Rutabaga, Sugarbeet, Sweet Cassava, Sweet Potato, Tanier, True Yam, Turnip, and Turnip-Rooted Chervil.

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Intrepid 2F (fl oz/acre)	Application Timing	Restrictions
beet armyworm cabbage looper cutworms (suppression) fall armyworm garden webworm imported cabbageworm southern armyworm true armyworm yellowstriped armyworm	4 – 8 (0.06 – 0.12 lb ai/acre)	For early season applications only to young crops and small plants. Apply at first sign of feeding damage or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities.	<ul style="list-style-type: none"> • Do not apply more than 16 fl oz per acre per application or 64 fl oz of Intrepid 2F (1 lb ai) per acre per season. • Pre-harvest interval: Do not harvest within 1 day of application. • See Rotational Crop Restrictions.

ACCEPTED

OCT 5 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-442

5/17

beet armyworm cabbage looper cabbage webworm cross-striped cabbageworm cutworms (suppression) fall armyworm garden webworm imported cabbageworm southern armyworm true armyworm yellowstriped armyworm	8 - 10 (0.12 – 0.16 lb ai/acre)	For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. For heavy infestations, continuous moth flights, and/or egg masses and larvae in all stages of development.	
diamondback moth (suppression only)	12 - 16 (0.19 – 0.25 lb ai/acre)	Infestations and crop damage are reduced when applied at initiation of egg laying.	

Specific Use Directions

Ground Application: It is recommended to apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. It is recommended to apply a minimum of 20 gallons per acre to densely foliated or difficult to cover crops to ensure thorough coverage. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days. **Note:** These rotational crop restrictions supercede prior restrictions of 356 days for leaves of root and tuber vegetables and the prohibition for all other crops.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R846-021

EPA-accepted ___/___/___

Initial printing.

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F

EPA Reg. No. 62719-442

For Control of Lepidoptera Larvae Infesting Spearmint and Peppermint

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid[®] 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F:

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworms cutworms loopers	10 - 16 (0.16 - 0.25 lb ai/acre)	Scout crops on a regular basis and treat as soon as economic thresholds have been met. Target small larvae and egg masses when possible. Use the higher rates in the rate range for high infestations and when extended residual is needed. Reapply at 14- to 21-day intervals when there are continuing infestations.	<ul style="list-style-type: none"> • Do not apply more than 16 fl oz per acre per application or 64 fl oz of Intrepid 2F (1.0 lb ai) per acre per season. • Pre-harvest Interval: Do not harvest within 14 days of application.

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 5 gallons per acre. Aircraft should be calibrated to assure uniform coverage of the target crop. Follow all guidelines on the main label.


Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R846-028

EPA-accepted: ___/___/___

Initial printing.



F3B / Intrepid 2F / Prop Sec 3 Supp Legume Veggies & Foliage (6A, 6B, 7A) / 02-11-04
file: R846-022 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Legume Veggies & Foliage (Subgroup 6A, 6B, 7A)
11Feb04d.doc

7/17

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F for control or suppression of lepidoptera larvae infesting legume vegetables (succulent) (crop subgroups 6A and 6B) and foliage of legume vegetables (except soybean) (crop subgroup 7A).

*Trademark of Dow AgroSciences LLC

8/17

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid* 2F EPA Reg. No. 62719-442

For Control or Suppression of Lepidoptera Larvae Infesting Legume Vegetables (Succulent) (Subgroup 6A, 6B[†]) and Foliage of Legume Vegetables (Except Soybean) (Subgroup 7A^{††})

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

[†]Legume vegetables (crop subgroup 6A and 6B) including, but not limited to: Asparagus Bean, Blackeyed Pea, Chinese Longbean, Cowpea, Dwarf Pea, Edible-Pod Pea, English Pea, Garden Pea, Green Lima Bean, Green Pea, Jackbean, Moth Bean, Pigeon Pea, Runner Bean, Snap Bean, Snow Pea, Soybean (Immature Seed), Southern Pea, Succulent Broad Bean, Sugar Snap Pea, Sword Bean, Wax Bean, Yardlong Bean

^{††}Foliage of legume vegetables (crop subgroup 7A) including, but not limited to any cultivar of bean and field pea (except soybean).

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Target Pests	Intrepid 2F (fl oz/acre)	Application Timing	Restrictions
alfalfa looper beet armyworm cabbage looper European corn borer fall armyworm southern armyworm tomato hornworm true armyworm yellowstriped armyworm western yellowstriped armyworm	4 - 8 (0.06 - 0.12 lb ai/acre)	For early season applications only to young crops and small plants. Apply at first sign of feeding damage or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities.	<ul style="list-style-type: none"> • Do not apply more than 16 fl oz per acre per application or 64 fl oz of Intrepid 2F (1 lb ai) per acre per season. • Do not make more than 4 applications per acre per season. • Pre-harvest Interval: Do not harvest within 7 days of application. • Re-treatment Interval: 7 days • See Rotational Crop Restrictions.
	8 - 16 (0.12 - 0.25 lb ai/acre)	For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. For heavy infestations, continuous moth flights, and/or egg masses and larvae in all stages of development, a 7-14 re-treatment interval is recommended to protect new growth until moth flights and/or larval infestations subside.	

ACCEPTED

OCT 5 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 62719-442

9/17

corn earworm <i>(Helioverpa/Heli othis)</i> (suppression only)	10 - 16 (0.16 – 0.25 lb ai/acre)	Apply at first sign of feeding damage or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities. May provide partial control when infestations reach high levels.	
tomato pinworm (suppression only)		Leafmining and infestations of leafmining phase are reduced when applied at initiation of egg laying.	

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult to cover crops to ensure thorough coverage. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre.

Spray Adjuvants: The addition of agricultural adjuvants to Intrepid 2F sprays will improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. Always add adjuvants last in the mixing process.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days. **Note:** These rotational crop restrictions supercede prior restrictions of 356 days for legume vegetables (succulent) and foliage of legume vegetables (except soybean) and the prohibition for all other crops.

Note: This product may be hazardous to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R846-022

EPA-accepted ___/___/___

Initial printing.

10/17

F3B / Intrepid 2F / Prop Sec 3 Supp Papaya et al / 02-27-04
file: R846-017 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Papaya et al 27Feb04d.doc

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F for control of lepidoptera larvae infesting black sapote, canistel, mamey sapote, mango, papaya, sapodilla, and star apple.

*Trademark of Dow AgroSciences LLC

11/17

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid* 2F

EPA Reg. No. 62719-442

For Control of Lepidoptera Larvae Infesting Black Sapote, Canistel, Mamey Sapote, Mango, Papaya, Sapodilla, and Star Apple

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworms citrus peelminer leafrollers orange tortrix	12 - 16 (0.19 - 0.25 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> • Do not apply more than 64 fl oz of Intrepid 2F (1 lb ai) per acre per season or make more than 5 applications per crop per year. • Re-treatment Interval: 10 days • Pre-harvest Interval: Do not harvest within 3 days of application.

Specific Use Directions

Ground Application: It is recommended to apply a minimum of 50 gallons per acre to trees 10 feet tall or less by conventional ground equipment. For trees greater than 10 feet tall, it is recommended to apply a minimum of 100 gallons per acre by conventional ground equipment. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.


Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R846-017

EPA-accepted ___/___/___

Initial printing.



12/17

F3B / Intrepid 2F / Prop Sec 3 Supp Cilantro Leaves / 01-22-04
file: R846-018 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Cilantro Leaves 22Jan04d.doc

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F to control or suppress lepidoptera larvae infesting cilantro leaves.

*Trademark of Dow AgroSciences LLC

13/17

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid* 2F EPA Reg. No. 62719-442

For Control or Suppression of Lepidoptera Larvae Infesting Cilantro Leaves

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
beet armyworm cabbage looper fall armyworm garden webworm imported cabbageworm southern armyworm true armyworm yellowstriped armyworm	4 - 8 (0.06 - 0.12 lb ai/acre)	For early season applications only to young crops and small plants. Apply at first sign of feeding damage or when infestations reach threshold levels as defined by a cooperative extension service or other qualified professional authorities.	<ul style="list-style-type: none"> • Do not apply more than 16 fl oz per acre per application or 64 fl oz of Intrepid 2F (1 lb ai) per acre per season. • Pre-harvest Interval: Do not harvest within 1 day of application. • See Rotational Crop Restrictions below.
beet armyworm cabbage looper cabbage webworm cross-striped cabbageworm fall armyworm garden webworm imported cabbageworm southern armyworm true armyworm yellowstriped armyworm	8 - 10 (0.12 - 0.16 lb ai/acre)	For mid- to late-season applications, heavier infestations, and under conditions in which thorough coverage is more difficult. For heavy infestations, continuous moth flights, and/or egg masses and larvae in all stages of development, a 10-14 day re-treatment interval is required to protect new growth until moth flights and/or hits subside.	
diamondback moth (suppression only)	12 - 16 (0.19 - 0.25 lb ai/acre)	Infestations and crop damage are reduced when applied at initiation of egg laying.	

ACCEPTED

OCT 5 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-442

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop or small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult to cover crops to ensure thorough coverage. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre.

Spray Adjuvants: The addition of agricultural adjuvants to Intrepid 2F sprays will improve initial spray

14
/ 17

deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. Always add adjuvants last in the mixing process.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days. **Note:** These rotational crop restrictions supercede prior restrictions of 356 days for cilantro leaves and the prohibition for all other crops.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R846-018

EPA-accepted ___/___/___

Initial printing.

15/17

F3B / Intrepid 2F / Prop Sec 3 Supp Root Veggies / 02-11-04
file: R846-019 Intrepid 2F Sec 3 Lepidoptera Larvae Control in Root Veggies (Subgroup 1A) 11Feb04d.doc

Intrepid* 2F

EPA Reg. No. 62719-442

Proposed Section 3 supplemental labeling (in support of IR-4 program) for use of Intrepid 2F for control of lepidoptera larvae infesting root vegetables (crop subgroup 1A).

*Trademark of Dow AgroSciences LLC

16/17

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid* 2F

EPA Reg. No. 62719-442

For Control or Suppression of Lepidoptera Larvae Infesting Root Vegetables (Subgroup 1A[†])

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid* 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

[†]Root vegetables (subgroup 1A) including, but not limited to: Black Salsify, Carrot, Celeriac, Chicory, Edible Burdock, Garden Beet, Ginseng, Horseradish, Parsnip, Oriental Radish, Radish, Rutabaga, Salsify, Skirret, Spanish Salsify, Sugarbeet, Turnip, Turnip-Rooted Chervil, and Turnip-Rooted Parsley.

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworms cabbageworms cutworm (suppression) loopers saltmarsh caterpillar webworms	6 - 16 (0.09 - 0.25 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> • Do not apply more than 64 fl oz of Intrepid 2F (1 lb ai) per acre per season. • Re-treatment Interval: 14 days • Pre-harvest Interval: Do not harvest sugarbeet within 7 days of application; do not harvest all other root vegetables within 14 days of application. • See Rotational Crop Restrictions below.

ACCEPTED

OCT 5 2004

 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-442

Specific Use Directions

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

17/17

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days. **Note:** These rotational crop restrictions supercede prior restrictions of 356 days for root vegetables and the prohibition for all other crops.

Note: This product is toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC
R846-019
EPA-accepted ___/___/___
Initial printing.