

62719-445

05/15/2006

1/28

Base Label:

(Logo) Dow AgroSciences

Dimension® Ultra 40WP

Specialty Herbicide

Dimension Ultra 40WP provides control of listed annual grasses and broadleaf weeds in established lawns, commercial sod farms, noncropland and industrial sites, ornamental turf (including golf course fairways, roughs, tee boxes), landscape, field grown and container ornamentals

In New York State, this product may only be used by commercial applicators at no more than 0.625 pounds per acre per year (10 ounces) or 2 water-soluble pouches per acre per year (0.25 pounds active ingredient).

Active Ingredient

dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)- S,S-dimethyl ester	40.0%
Inert Ingredients	60.0%
Total	100.0%

ACCEPTED

MAY 15 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Each 5 ounce water-soluble pouch contains 0.125 lb of active ingredient.
Product protected by U.S. Patent No. 4,692,184. Other patents pending

62719-445

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation •Harmful If Absorbed Through The Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers (other than mixer loaders) must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Mixer and Loaders must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-445

EPA Est. _____

©Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight 2.5 lb

3/28

8 x 5 oz Water-Soluble Pouches

4/28

Label Booklet:

(Logo) Dow AgroSciences

Dimension[®] Ultra 40WP

Specialty Herbicide

Dimension Ultra 40WP provides control of listed annual grasses and broadleaf weeds in established lawns, commercial sod farms, noncropland and industrial sites, ornamental turf (including golf course fairways, roughs, tee boxes), landscape, field grown and container ornamentals

In New York State, this product may only be used by commercial applicators at no more than 0.625 pounds per acre per year (10 ounces) or 2 water-soluble pouches per acre per year (0.25 pounds active ingredient).

Active Ingredient

dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)- S,S-dimethyl ester 40.0%

Inert Ingredients 60.0%

Total 100.0%

Each 5 ounce water-soluble pouch contains 0.125 lb of active ingredient.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for Precautionary Statements and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read **Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies** at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-445

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight 2.5 lb
8 x 5 oz Water-Soluble Pouches

5/28

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Use Directions for Professional Turf Use	-
Use Directions for Noncropland and Industrial Sites	-
Use Directions for Ornamental Plantings and Containers	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

6/28

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Causes Eye Irritation •Harmful If Absorbed Through The Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers (other than mixer loaders) must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Mixer and Loaders must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

REFORMULATION OR REPACKAGING OF THIS PRODUCT IS PROHIBITED.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements:

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

- **Keep unprotected persons out of treated area until sprays have dried.**

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F.

Product Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Container Disposal: Completely empty bags into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Spill: In case of spill, observe all protective and safety precautions when cleaning up spills. Dispose of material according to "product disposal" directions listed above.

8/28

General Information

This product is not for application by homeowners.

Dimension* Ultra 40WP specialty herbicide provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns, commercial sod farms, noncropland and industrial sites and ornamental turf (including golf course fairways, roughs, tee boxes), landscape ornamentals, field grown and container nursery ornamentals.

Except for control of crabgrass up to initiation of tillering, this product will not control emerged weeds. This stage of crabgrass growth generally corresponds to the time when seedlings are first visible in established turfgrasses. Applications to crabgrass after initiation of tillering will not provide satisfactory control. All other applications of this product should be made preemergence (prior to germination of target weeds).

This product is not effective until activated by 1/2 inch or more of rainfall or irrigation. Applications should be timed to ensure that activation has occurred prior to tillering stage of crabgrass development or prior to germination of all other weeds.

Note: In the state of New York State this product may be applied only by commercial applicators.

Mixing Instructions

Handling of Water Soluble Pouches: The enclosed pouches are water soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal over-wrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from over-wrap except to add directly to the spray tank.

Mixing Instructions

Dimension Ultra 40WP Alone with Water as the Carrier:

Be sure the sprayer is clean and not contaminated with other materials prior to use. Fill the mixing tank 1/2 to 3/4 full with clean water and begin agitation. Be certain that the agitation system is working properly and creates a rolling or rippling on the liquid surface. With the agitator running, add the required number of unopened water-soluble pouches into the tank and allow time for complete mixing prior to adding any additional materials to the tank. Depending on the water temperature and degree of agitation, disintegration and complete mixing of contents of water soluble pouches should occur within ten minutes after their addition to water. Finish by adding the remainder of the required amount of water to the spray tank. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the water source.

Maintain agitation of spray mixture during mixing and application to ensure uniformity of spray mixture and prevent settling.

9/28

Dimension Ultra 40WP Alone with Fluid Fertilizer as the Carrier

First, determine the compatibility of this product with the desired fluid fertilizer by mixing small proportional quantities in advance. Follow instructions in the "Compatibility Test" section of this label. Then follow the mixing procedure listed below for tank mixtures.

Tank Mixtures

First, determine the compatibility of this product and the desired tank mixture partner product(s) in the appropriate carrier (water or fluid fertilizer) by mixing small proportional quantities in advance. See the "Physical Compatibility Test" section of this label below. Then adhere to the following mixing procedure:

Note: Dimension Ultra 40WP is compatible with boron and spray oils; however, the water soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

Mixing Procedure:

1. Place a 20 to 35 mesh screen or wetting basket over the filling port.
2. Fill the previously cleaned sprayer half full with the appropriate carrier. Start agitation and continue agitation through mixing and spraying operations.
3. Add a compatibility agent if needed. Read and follow all of the information found on the product label for the selected compatibility agent. Check for adequate agitation.
4. Add Dimension Ultra 40WP to the tank and allow sufficient time for the water-soluble pouches to dissolve and contents to mix completely before adding other materials. Add any other wettable powder or dry flowable formulation. **When mixing this product or any other wettable powder or dry flowable product with fluid fertilizer, premix with water to form a slurry and then add slowly to the mixing tank.**
5. If a flowable pesticide formulation is used, premix with one part water, and add slowly to tank.
6. Add emulsifiable concentrate pesticide formulations to the tank.
7. Add water-soluble liquid pesticide formulations followed by surfactants, marker dyes or foams, or drift control additives while continuing the filling process.

Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, agitate thoroughly to re-suspend the mixture before resuming spraying.

Physical Compatibility Test

Before mixing this product with fluid fertilizers and/or other pesticides, it is recommended that the compatibility of the tank mixture be tested by mixing proportionate amounts of each component in a small glass jar according to the following instructions:

Compatibility Test Mixing Instructions

Pesticide Formulation	If	Amount of Pesticide added to Spray Carrier (assuming volume is 25 GPA) ADD:
	Rate per Acre is:	Level Teaspoons per Pint Jar of Carrier Solution
Dry	1 lb	1-1/2
Liquid	1 qt	1

This compatibility test is designed for 25 gallons of spray solution per acre. The table above gives general guidelines for use rate ratios of pesticides to be tank-mixed with this product. Determine the amount of pesticide to tank-mix by referring to the pesticide label(s). Then, calculate the amount of pesticide to add to jar based on use rate ratios in table. For a use rate of 1 pound per acre of dry pesticide add 1-1/2 teaspoons to the jar, and for a use rate of 1 quart per acre of liquid pesticide, add 1 teaspoon to the jar. This product should be added based on use rate ratios for liquid pesticides (for a use

10/28

rate of 1 quart per acre, add 1 teaspoon to the jar). For changes in spray volume or herbicide rate, make appropriate changes in the ingredients for the test. Shake well to mix.

If pesticide(s) does not form crystals, flakes, sludge, gels, oily films or layers, then the tested components are compatible. Incompatibility in any of the above-described forms will usually occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Repeat the above compatibility test with a suitable compatibility agent (one-half teaspoon per pint jar is equivalent to 2 pints per 100 gallons of spray solution). Do not use mixtures that show incompatible signs such as formation of crystals, flakes, sludge, gels, oil films or layers.

Weed Control in Turfgrasses

Use Precautions for Turf Use

Turfgrass Safety

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging before making the first application of this product. Use of this product on turf that is not well-established, or has been weakened by weather-, pest-, disease-, chemical-, mechanical or other related stress, may result in turf injury.

This product should only be applied to turf that is composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Common Name	Scientific Name
bentgrass, creeping ^(a)	<i>Agrostis palustris</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, fine ^(b)	<i>Festuca rubra</i>
fescue, tall	<i>Festuca arundinacea</i>
ryegrass, perennial	<i>Lolium perenne</i>

Warm-Season Grasses

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum</i>
bermudagrass ^(c)	<i>Cynodon dactylon</i>
buffalograss ^(d)	<i>Buchloe dactyloides</i>
carpetgrass	<i>Axonopus affinis</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
seashore paspalum	<i>Paspalum vaginatum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>

- Do not use on golf course putting greens.
- Use of this product is not recommended on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' as undesirable turfgrass injury may result. Not all varieties of creeping bentgrass have been tested.
Do not apply this product to colonial bentgrass (*Agrostis tenuis*) varieties.
- Use of this product on certain varieties of fine fescue is not recommended as use may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this

W/28

product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

- d) Use of this product on 'Tifgreen' (328) hybrid bermudagrass is not recommended as use may result in undesirable turfgrass injury.
- e) **Do not** use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Reseeding, Overseeding, or Sprigging

Reseeding, overseeding, or sprigging of treated areas within 10 weeks after a single application of this product, or 4 months after a split application program totaling more than 0.46 oz/1000 sq ft (1.25 lb/acre), may inhibit the establishment of desirable turfgrasses. However, overseeding of bermudagrass with perennial ryegrass 8 weeks after an application or as early as 6 weeks after application if slight injury to perennial ryegrass can be tolerated, is a recommended exception.

When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Sod Production

- It is recommended that sod be established for at least six (6) months before a Dimension Ultra 40WP application is made.
- **Do not** apply this product within three (3) months of harvest.

Other Use Precautions

Early postemergence applications of this product will control crabgrass only if applied prior to the fifth leaf (first tiller) stage of growth of crabgrass.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product.

- **Do not** apply this product until the grass has recovered from these cultural practices.
- **Do not** use clippings from treated turf for mulching around vegetables or fruit trees.
- **Do not** graze livestock or feed foliage cut from areas treated with this product.

Application Directions

Application Equipment and Instructions

Apply this product through conventional liquid application equipment in a sufficient volume of carrier solution to provide a uniform spray distribution. A hand held spray gun may be used. **Do not** apply this product through liquid application equipment that uses cluster spray nozzles or other boomless spray equipment due to variability in application use rates and spray patterns. Calibrate application equipment prior to usage. Avoid streaking, skips, or excess overlaps during application. The use of marker dyes or foams aids in making more accurate applications.

Chemigation: **Do not** apply this product through any type of irrigation system.

Control of Crabgrass

Preemergence and Early Postemergence Control.

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth.

12/28

after the crabgrass has emerged. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Early post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2 - 8 week longer period of time (depending on climatic conditions and crabgrass growth rates) to make applications and still control crabgrass.

Application Frequency and Timing.

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

Spring Applications

For applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1, depending on the user's location, the mowing height of the turfgrass, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

Use Program 2 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control up to crabgrass tillering at sites where turfgrass is cut relatively high (e.g., residential lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control up to crabgrass tillering at sites where turfgrass is cut relatively low (e.g., golf fairways). Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired. Split applications may also be made, with the rates in Table 1 being split across two applications made 5-10 weeks apart. Such applications may provide improved weed control.

Region	Application Rates	Program 1	Program 2	Program 3
All states and parts of states not listed in transition, south, coastal south or west.	Number of square feet per water-soluble pouch per treatment	29,040	21,780	14,520
	Number of pouches per acre	1.5	2	3
	Pounds per acre	0.47	0.625	0.95
	Pounds ai/per acre	0.188	0.25	0.38
Transition- DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI	Number of square feet per water-soluble pouch per treatment	21,780	14,520	10,900
	Number of pouches per treated acre	2	3	4
	Pounds per acre	0.625	0.95	1.25
	Pounds ai/per acre	0.25	0.38	0.5

B/208

South- AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	Number of square feet per water-soluble pouch per treatment	14,520	21,780	17,424
	Number of pouches per acre	3	2 + 2 ^c	2.5 + 2.5 ^{b,c}
	Pounds per acre	0.95	0.625 + 0.625	0.78 + 0.78
	Pounds ai/per acre	0.38	0.25 + 0.25	0.31 + 0.31
Coastal South- - HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	Number of square feet per water-soluble pouch per treatment	21,780	17,424	14,520
	Number of pouches per acre	2 + 2 ^c	2.5 + 2.5 ^{b,c}	3 + 3 ^c
	Pounds per acre	0.625 + 0.625	0.78 + 0.78	0.95 + 0.95
	Pounds ai/per acre	0.25 + 0.25	0.31 + 0.31	0.38 + 0.38
WEST - AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	Number of square feet per water-soluble pouch per treatment	29,040 - 21,780	21,780 - 14,520	21,780
	Number of pouches per acre	1.5 - 2 ^b	2 - 3	2 + 2 ^c
	Pounds per acre	0.47 - 0.625	0.625 - 0.95	0.625 + 0.625
	Pounds ai/per acre	0.18 - 0.25	0.25 - 0.38	0.25 + 0.25

- In New York State, do not apply more than 2 water-soluble pouches per acre or 0.23 oz/1000 sq ft (0.625 lb/acre) per year.
- The water-soluble bag should not be opened. The water-soluble bag is not designed to be re-measured or subdivided. Treat an appropriate number of acres to use a whole number of bags.
- Preemergence applications totaling more than 0.5 lb ai/acre (greater than 1.25 lb/acre of Dimension Ultra 40WP) must be applied as a split application. The recommended interval for split applications is 5 to 10 weeks apart. Early postemergence applications are limited to 0.5 lb ai/acre (1.25 lb of Dimension Ultra 40WP) per application.

Maximum Use Rates (All Turf Uses)

- Do not apply more than 0.5 lb ai/acre per application or more than 1.5 lb ai/acre per year using split or sequential applications. This is equivalent to 4 water-soluble pouches per acre or 0.46 oz/1000 sq ft (1.25 lb/acre of product) per application, and 12 water-soluble pouches per acre or 1.38 oz/1000 sq ft (3.75 lb/acre of product) per year.
- In New York State, do not apply more than 0.25 lb ai per year. This is equivalent to 2 water-soluble pouches (0.625 lb/acre) per acre or 0.23 oz/1000 sq ft of Dimension Ultra 40WP.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. Do not exceed maximum use rates as described above.

The Fall Application may be followed by an appropriately timed spring application to provide season-long weed control provided the maximum use rate per year is not exceeded.

Tank Mixtures for Postemergence Control of Crabgrass

Application of this product alone provides early postemergence control of crabgrass when treated prior to the tillering stage of growth. For postemergence control of tillered crabgrass up to 3 tillers, this product may be applied in tank mix combination with either MSMA or Acclaim.

If preemergence herbicides have been applied prior to a postemergence application, use the applicable Program 1 or Program 2 use rate of Dimension Ultra 40WP for your area; otherwise, apply the Program 3 use rate.

14/28

A compatibility test (see the "Physical Compatibility Test" in "Mixing Instructions" section) is recommended before tank mixing this product with fluid fertilizers and/or either MSMA or Acclaim herbicide.

Refer to the labels for MSMA or Acclaim for information on tolerance of specific turfgrass species and observe all limitations, precautionary statements, and use restrictions on the respective labels when using them in tank mixtures. The addition of a nonionic surfactant may improve control. Always read and follow the surfactant manufacturer's label recommendations.

Control of Other Grass and Broadleaf Weeds

Weeds Controlled

Used as directed, Dimension Ultra 40WP will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence.

Note: This product will not control emerged broadleaf weeds or grasses, except for crabgrass up to tillering stage of development. Therefore, the area to be treated should be free of weeds prior to application.

Numbers in parentheses (-) refer to footnotes.

Grasses

barley	<i>Hordeum</i> spp.
barnyardgrass	<i>Echinochloa crus-galli</i>
bluegrass, annual (1)	<i>Poa annua</i>
brome	<i>Bromus</i> spp.
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
crabgrass, Southern	<i>Digitaria ciliaris</i>
crowfootgrass	<i>Dactyloctenium aegyptium</i>
dallisgrass (seedling)	<i>Paspalum dilatatum</i>
goosegrass	<i>Eleusine indica</i>
foxtail, giant	<i>Setaria faberi</i>
foxtail, green	<i>Setaria verdi</i>
foxtail, yellow	<i>Pennisetum clandestinum</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
Mary's grass	<i>Microstegium vimineum</i>
oats, wild	<i>Avena fatua</i>
ryegrass (annual & perennial)	<i>Lolium</i> spp.
sandbur	<i>Cenchrus</i> spp.
smutgrass	<i>Sporobolus indicus</i>

- (1) **For Extended Control of Annual Poa (*Poa annua*) in South and Coastal South Regions:** An initial application at 1.25 lb/acre (0.5 lb ai) is recommended 6 weeks before overseeding followed by a second application of 0.625 to 1.25 lb/acre (0.25 to 0.5 lb ai) 120 days after overseeding. Some injury to overseeded perennial ryegrass may occur (See "Reseeding, Overseeding, or Sprigging" precautions under "Use Precautions for Turf Use").

Broadleaf Weeds

bittercress	<i>Cardamine</i> spp.
carpetweed	<i>Mollugo verticillata</i>
chickweed	<i>Stellaria</i> spp.
coneflower, purple	<i>Rudbeckia purpurea</i>
geranium, Carolina	<i>Geranium carolinianum</i>

15/28

henbit	<i>Lamium</i> spp.
knotweed, prostrate	<i>Polygonum aviculare</i>
lespedeza, common	<i>Lespedeza striata</i>
marestail	<i>Conyza canadensis</i>
medic, black	<i>Medicago lupulina</i>
mustard	<i>Brassica</i> spp.
oxalis, buttercup	<i>Oxalis pes-caprae</i>
pineappleweed	<i>Matricaria matricarioides</i>
pigweed, redroot	<i>Amaranthus retroflexus</i>
parsley-piert	<i>Alchemilla arvensis</i>
purslane, common	<i>Portulaca oleracea</i>
rocket, London	<i>Sisymbrium irio</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>
speedwell, corn	<i>Veronica arvensis</i>
spurge, garden	<i>Euphorbia hirta</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Use Directions for Noncropland and Industrial Sites

Dimension Ultra 40WP may be applied for preemergence control of listed annual grasses and broadleaf weeds in terrestrial noncrop areas such as farm yards, fence rows, highway or roadside utility and railroad rights-of-way, airports, recreation areas, campgrounds, and industrial sites such as lumber yards, tank farms, and storage areas. Refer to "Control of other Grasses and Broadleaf Weeds" section above for a listing of weeds controlled.

Applied preemergence, Dimension Ultra 40WP controls weeds as they germinate. This product will not control established weeds. Make applications prior to germination of target weeds or to bare ground. The best weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide.

To be effective, Dimension Ultra 40WP must be activated by 1/2 inch or more of rainfall or irrigation prior to germination of target weeds. Once the treatment is activated, avoid mixing of the soil surface that will expose untreated soil.

Application Rates:

Dimension Ultra 40WP		
Rate per acre	Rate per 1000 sq ft	Square feet per water soluble pouch
20 oz	0.46 oz	10,890

Sequential applications may be made at 3 to 4 month intervals for extended preemergence weed control. Do not exceed maximum use rates per year.

Maximum Use Rates

- Do not apply more than 0.5 lb ai/acre per application or more than 1.5 lb ai/acre per year. This is equivalent to 4 water-soluble pouches per acre or 0.46 oz/1000 sq ft (1.25 lb/acre) per application, and 12 water-soluble pouches per acre or 1.375 oz/1000 sq ft (3.75 lb/acre) per year.
- In New York State, do not apply more than 0.25 lb ai per year. This is equivalent to 2 water-soluble pouches (0.625 lb/acre) per acre or 0.23 oz/1000 sq ft of Dimension Ultra 40WP.

16/208

Use Precautions for Noncrop Areas

- Do not apply when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not graze livestock or feed forage cut from areas treated with this product.
- For ornamentals within non-crop areas, apply only after transplanting when soil around roots has been thoroughly settled by rainfall or irrigation or injury may result.

Use Directions For Ornamentals

Dimension* Ultra 40WP specialty herbicide provides preemergence control of listed annual grass and broadleaf weeds in areas planted with tolerant ornamental plants listed on this label. It is intended for use on plants being grown for aesthetic purposes in landscaped areas. When applied as directed, the ornamental plants listed on this label have shown tolerance to post-directed applications of Dimension Ultra 40WP herbicide. It is impossible, however, to evaluate tolerance to this product on all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, it is strongly recommended that a limited area involving only a few plants be treated for tolerance evaluation prior to large-scale applications.

Treatment of Turf or Ornamental Species not listed on the Label for Dimension Ultra 40WP

Although this label contains a large number of ornamental species, it is not possible to include all of the ornamental plants that may be encountered in nursery or landscape settings. Users who wish to use this product on a plant species not listed on this label may determine the suitability for such use by treating a small area or small number of plants at a recommended rate. Prior to treatment of larger areas, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. **The user assumes the responsibility for any plant damage or other liability resulting from use of Dimension Ultra 40WP on species not recommended on this label.**

Chemigation: Do not apply through any type of irrigation system.

Use Rates

Make applications prior to germination of target weed species in spring, summer or fall. Sequential applications may be made at 3 to 4 month intervals to maintain weed control. Do not exceed maximum use rates per year

Apply Dimension Ultra 40WP with a calibrated sprayer that will assure accurate, uniform, spray distribution. In general, Dimension Ultra 40WP should be thoroughly mixed with clean water at 4 water-soluble pouches per acre or 0.46 oz of product /1000 square feet (1.25 lb/acre) per application and applied at 20 to 40 psi in a minimum of one gallon of water per 1000 square feet. One 5 ounce water-soluble pouch treats 10,890 square feet. This product may be applied as a single application or as a split application prior to weed seed germination.

Application Rates:

Dimension Ultra 40WP		
Rate per acre	Rate per 1000 sq ft	Square feet per water soluble pouch
20 oz	0.46 oz	10,890

Maximum Use Rates

- Do not apply more than 0.5 lb ai/acre per application or more than 1.5 lb ai/acre per year using split or sequential applications. This is equivalent to 4 water-soluble pouches per acre or 0.46 oz/1000 sq ft (1.25 lb/acre) per application, and 12 water-soluble pouches per acre or 1.375 oz/1000 sq ft (3.75 lb/acre) per year.

M/20

- In New York State, do not apply more than 0.25 lb ai per year. This is equivalent to 2 water-soluble pouches (0.625 lb/acre) per acre or 0.23 oz/1000 sq ft of Dimension Ultra 40WP.

Application Timing and Recommendations

Apply Dimension Ultra 40WP as a post-directed or over the top spray around established ornamentals as directed on the tolerance ornamentals table. When making directed spray applications, direct spray to the soil at the base of the ornamentals avoiding contact or drift to foliage.

Dimension Ultra 40WP is a preemergence herbicide that will control germinating weeds. It will not control weeds that are established. Make applications prior to weed seed germination. The best weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, existing vegetation should be controlled by hand weeding, by cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of this product and plant roots.

Precautions

- Apply this product only to established ornamentals (after transplanting when soil around roots has been thoroughly settled by rainfall or irrigation).
- **Do not** apply this product to bare roots of ornamental plants as injury may result.
- **Do not** incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- **Do not** apply to or around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or diseases).
- Drift or direct sprays of this product may injure foliage of non-target plants. **Do not** make applications when weather conditions favor drift to non-target areas.
- **Do not** apply this product on grasses grown for seed.
- **Do not** graze livestock or feed forage cut from areas treated with this product.
- **Do not** apply this product in enclosed structures and greenhouses.

Tolerant Ornamentals

When applied under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Under your growing conditions, it is strongly recommended that a limited number of plants be treated for tolerance evaluation, prior to initiating large-scale applications. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

Common Name	Botanical Name	Tolerant Cultivars	Recommended Application Method O = Over the top, D = Directed [†]	Recommended Approved Treatment Method C = Container Grown, F = Field
abelia, dwarf	<i>Abelia X grandiflora</i>	nana	D	F
abyssinian red banana	<i>Ensete ventricosum</i>	maurelii	O,D	C,F

18/208

ajuga	<i>Ajuga reptans</i> <i>Ajuga genevensis</i>	bronze bronze beauty	D	F
almond, flowering	<i>Prunus glandulosa</i>		D	F
apple †	<i>Malus pumila</i>		D	F
arborvitae	<i>Thuja occidentalis</i>	nigra	D	F
		pyramidalis	D	F
		smaragh	D	F
		techny	D	F
		woodwardii	D	F
arborvitae, golden	<i>Thuja orientalis</i>	aurea nana	O,D	C, F
aster, Chinese	<i>Callistephus chinensis</i>	dwarf queen	D	F
ash, green	<i>Fraxinus pennsylvanica</i>		D	F
ash, mountain	<i>Sorbus aucuparia</i>		D	F
ash, purple	<i>Fraxinus americana</i>		D	F
azalea	<i>Rhododendron</i> spp.	brilliant	D	F
		buccaneer	D	F
		carror	D	F
		chimes (belgian)	D	F
		elsie lee	D	F
		exbury	D	F
		fashion	D	F
		hardijzer beauty	D	F
		hershey red	D	F
		higasa	D	F
		hinocrimson	D	F
		holland (hybrid)	D	F
		marion lee	D	F
		northern lights	D	F
		orange cup	D	F
orchid lights	D	F		
snow	D	F		
southern charm	D	F		
azalea, flame	<i>Rhodocendron calendulaceum</i>		D	F
azalea, Kirishima			D	F
bamboo, heavenly			D	F
barberry	<i>Berberis thunbergii</i>	aurea	D	F
		dwarf pigmy	D	F
		green	D	F
		kobold	D	F
		pygmy red	D	F
		rose glow	D	F
barberry, purple		atropurpurea	D	F
basket flower	<i>Gaillardia grandiflora</i>		D	F

19/28

beach grass	<i>Ammophila breviligulata</i>		D	F
bearberry (common)	<i>Arctostaphylos uva-ursti</i>	Massachusetts	D	F
bee balm	<i>Monarda didyma</i>		D	F
begonia	<i>Begonia spp.</i>		D	F
birch, river	<i>Betula nigra</i>		D	F
blackeyed susan	<i>Rudbeckia hirta</i>	goldstrum	D	F
blanket flower	<i>Gaillardia spp.</i>		D	F
blueberry †	<i>Vaccinium spp.</i>	bluecrop blue jay jersey north blue northland	D D D D D	F F F F F
blue fescue	<i>Festuca ovina</i>		D	F
bottlebrush	<i>Callistemon citrinus</i>		D	F
boxwood, Japanese		japonica	D	F
boxwood, weller	<i>Buxus sempervirens</i>	winter gem	O,D	C,F
broom	<i>Cytisus spp. Genista pilosa</i>	moonlight Vancouver gold	D D	F F
bugle carpet			D	F
camellia	<i>Camellia japonica</i> <i>Camellia sasanqua</i>	debutante mathotiana supreme chansonette setsukgekka	D O,D D O,D	F C,F F C,F
candy tuft	<i>Iberis spp.</i>	Snow White	D	F
carex, variegated	<i>Carex</i>		D	F
cedar, red	<i>Juniperus virginiana</i>		D	F
celosia	<i>Celosia spp.</i>		D	F
centaura	<i>Centaurea montana</i>		D	F
cockscomb, plumosa	<i>Celosia cristata</i>	scarlet plumosa	D	F
coleus	<i>Coleus blumei</i>	red kewpie	D	F
columbine	<i>Aquilegia spp.</i>		D	F
coneflower, purple	<i>Echinacea purpurea</i>		D	F
copper leaf	<i>Acalypha wilkesiana</i>		D	F
coreopsis	<i>Coreopsis spp.</i>	moonbeam	D	F
corn flower	<i>Centaurea spp.</i>		D	F
cotoneaster	<i>Cotoneaster apiculatus</i>		D	F
coyotebrush	<i>Baccharis pilularis</i>		D	F
cycads	<i>Cycas revoluta</i>		D	F
cypress, bald	<i>Taxodium distichum</i>		D	F

20/28

cypress, Italian	<i>Cupressus sempervirens</i>	glauca	D	F
cypress, Japanese false	<i>Chamaecyparis obtusa</i>	gracilis	D	F
cypress, leyland	<i>Cupressocyparis leylandii</i>		D	F
daffodil	<i>Narcissus</i> spp.	King Alfred	D	F
daylilly	<i>Hemerocallis</i> spp.	spring glory	OD	C,F
		aztec gold	D	F
		bright yellow (hybrid)	D	F
		single gold (evergreen)	D	F
		Wilson's yellow	D	F
dianthus (sweet william)	<i>Dianthus barbatus</i>		D	F
delphinium	<i>Delphinium elatum</i>	magic fountain	D	F
dogwood	<i>Cornus florida</i>		D	F
dogwood, American	<i>Cornus sericea</i>	flavarimaea	D	F
douglas fir	<i>Pseudotsuga menziesii</i>		D	F
dusty miller	<i>Senecio cineraria</i>		D	F
elm, drake	<i>Ulmus parvifolia</i>		D	F
euryops	<i>Europs pectinatus</i>	viridis	O,D	C,F
eulaliagrass	<i>Miscanthus sinensis</i>	maiden grass, gracillimus	O,D	C,F
euonymus	<i>Euonymus fortunei</i>	argenteo-variegata	D	F
		auereo-marginata	D	F
		colorata	D	F
		emerald gaiety	D	F
		emerald 'n gold	D	F
		gold edge	D	F
		gold princess	D	F
		silver king	D	F
		tricolor	D	F
		vegetus	D	F
fan palm, European	<i>Chamaerops humilis</i>		D	F
fan palm, Mexican	<i>Washingtonia robusta</i>		D	F
fern(various)	<i>Asparagus</i> spp.		D	F
fescue	<i>Festuca glauca</i>		D	F
fetterbush	<i>Leucothoe fontanesiana</i>	rainbow	D	F
figus	<i>Ficus retusa</i>	nitidia	D	F
fir fraser	<i>Abies fraseri</i>		D	F
forsythia	<i>Forsythia X intermedia</i>	arnold dwarf	D	F
		bronxensis dwarf	D	
		lynwood gold	D	
		meadowlark	D	
		weeping	D	
	<i>Spring Glory</i>			
fountain grass	<i>Pennisetum</i>	rubrum	O,D	C, F

21/28

	<i>setaceum</i>			
fuchsia	<i>Fuchsia</i> spp.		D	F
galium	<i>Galium odoratum</i>		D	F
garlic, variegated society	<i>Thunbergia violacea</i>	variegata	O,D	C,F
gardenia	<i>Gardenia jasminoides</i>	white gem mystery radicans	O,D D	C,F F
geranium	<i>Pelargonium X hortorum</i>		D	F
gum	<i>Eucalyptus citriodora</i>		D	F
hawthorn	<i>Crataegus</i> spp.	cockspur white crimson cloud enchantress Jack Evans Washington white	D D D D D	F F F F F
heather, twisted	<i>Erica cinerea</i>	Mediterranean pink	D	F
hemlock, Canada	<i>Tsuga canadensis</i>		D	F
hibiscus	<i>Hibiscus</i> spp.	blue bird brilliant hula girl	D D D	F F F
holly	<i>Ilex</i> spp. <i>Ilex X meserveae</i> <i>Ilex X attenuata</i>	blue boy blue girl burfordii china girl compacta forsteri hellerie Japanese northern beauty needlepoint nellie r. stevens savanah	D D D D D D D D D D D D	F F F F F F F F F F F F
holly, Chinese	<i>Ilex cornuta</i>		D	F
holly, Japanese	<i>Ilex crenata</i>		O,D	C,F
holly, yaupon	<i>Ilex vomitoria</i>		D	F
honeysuckle	<i>Lonicera japonica</i>	clavey's dwarf halliana tatarian canadian white zebelli red hosta	D D D D D	F F F F F
hosta	<i>Hosta</i> spp. <i>Hosta lancifolia</i>	albo marginata	D D	F F
ice plant	<i>Carpobrotus edulis</i>		O,D	C,F
ice plant, white trailing	<i>Delosperma alba</i> CT		O,D	C,F
impatiens	<i>Impatiens</i> spp.			
iris	<i>Iris</i> spp.	dwarf blue wedgewood	D	F
ivy, English	<i>Hedera helix</i>	bulgaria	D	F

23/28

			D D D D D D D D D	F F F F F F F F
	<i>Juniperus sabina</i>	weberi youngstown yukon belle	D O,D	F C,F
king palm	<i>Archontophoenix cunninghamiana</i>		D D D	F F F
laurel, Australian	<i>Pittosporum tobira</i>		O,D	C,F
laurel, mountain	<i>Kalmia latifolia</i>		O,D	C,F
leucothoe	<i>Leucothoe fontanesiana</i>		D	F
ligustrum, Japanese	<i>Ligustrum japonicum</i>		D	F
lily, African	<i>Agapanthus africanus</i>	albus Peter Pan streamline	D	F
lily, African blue			D D O,D	F F C,F
lily of the valley	<i>Pieris japonica</i>	Mt. Fire	D	F
lilyturf	<i>Liriope muscari</i>	evergreen giant lilac beauty majestic monroe white silvery sunproof variegata	D	F
liriope, green	<i>Liriope spicata</i>		D D O,D D O,D D	F F C,F F C,F F
maple, amur	<i>Acer ginnala</i>	emerald elf	D	F
maple, Japanese	<i>Acer japonicum</i>		O,D	C,F
maple, Norway	<i>Acer platanoides</i>		D	F
maple, red †	<i>Acer rubrum</i>	red sunset	D	F
maple, silver	<i>Acer saccharinum</i>		O,D D	C,F F
maple sugar †	<i>Acer saccharum</i>		D	F
marigold	<i>Tagetes patula</i>	honeycomb variegata wheeler's dwarf	D	F
mock orange †	<i>Philadelphus spp</i>	golden	D	F

2/28

		snowflake double white	D D	F F
mondo grass	<i>Ophiopogon japonicus</i>		D	F
moss rose	<i>Portulaca grandiflora</i>	sunnyside	D	F
myrtle, crape	<i>Lagerstroemia indica</i>	faurei langer muskogee standard pink	D	F
myrtle, wax	<i>Myrica cerifer</i>		D D D D	F F F F
nandina	<i>Nandina domestica</i>	compacta nana	D	F
narcissus	<i>Narcissus Spp.</i>		D D	F F
oak, laurel	<i>Quercus laurifolia</i>		D	F
oak, pin	<i>Quercus palustris</i>		D	F
oak, red	<i>Quercus rubra</i>		D	F
oak, southern l	<i>Quercus virginiana</i>		D	F
oak, willow	<i>Quercus phellos</i>		D	F
oleander	<i>Nerium oleander</i>	hardy red petite pink sister agnes	D	F
osteospermum	<i>Osteospermum fruticosum</i>	wirrigig	D D D	F F F
pachysandra	<i>Pachysandra terminalis</i>		D	F
palm, bangalow			D	F
pampas grass	<i>Cortaderia selloana</i>		D	F
pansy	<i>Viola x wittrockiana</i>		D	F
paper flower	<i>Bougainvillea glabra</i>	Barbara Karst James Walker	D	F
peach †	<i>Prunus persica</i>		O,D	C,F
periwinkle, dwarf	<i>Vinca minor</i>		D	F
petunia	<i>Petunia X hybrida</i>	picoti	D	F
photinia, red tip	<i>Photinia X fraseri</i>		D	F
pieris	<i>Pieris japonica</i>		D	F
pine, Afghan	<i>pinus eldarica</i>		D	F
pine, Australian	<i>Pinus nigra</i> <i>Pieris Taiwanensis</i>		O,D	C,F
pine, Japanese black	<i>Pinus thunbergiana</i>		O,D	C,F
pine, loblolly	<i>Pinus taeda</i>		D	F
pine, longleaf	<i>Pinus palustris</i>		D	F
pine, mugo	<i>Pinus mugho</i>		D	F
pine, scotch	<i>Pinus sylvestris</i>		D	F

25/28

pine, slash	<i>Pinus elliotii</i>		D	F
pine, Swiss mt.	<i>Pinus mugo</i>		D	F
pine, Virginia	<i>Pinus virginiana</i>		D	F
pine, white	<i>Pinus strobus</i>		D	F
pineapple, guava †	<i>Feijoa sellowiana</i>		O,D	C,F
plumbago, cape	<i>Plumbago auriculata</i>	Royal cape	D	F
pittosporum, Japan			O,D	C,F
potentilla	<i>Potentilla nepalensis</i> <i>Potentilla fruticosa</i>	abbotswood	D	F
privet	<i>Ligustrum japonicum</i>	golden vicary regal texanum wax yellow tipped	D D	F F
privet, glossy	<i>Ligustrum lucidum</i>		D D O,D D D	F F C,F F F
pyracantha	<i>Pyracantha koidzumii</i>	gnome lalandei victory C,F	D	F
queen palm	<i>Arecastrum rammanzoffianum</i>		D D O,D	F F
quince, Japanese †			D D	F F
rhododendron	<i>Rhododendron spp</i>	album Cunningham white fashion hardy pjm purple gem silvery pink	D	F
rhododendron, Carolina	<i>Rhododendron carolinianum</i>		D D D D D D	F F F F F F
rhododendron, catawba	<i>Rhododendron catawbiense</i>		D D	F F
ribbon grass	<i>Phalaris arundinacea</i>		D D	F F
rockcress	<i>Arabis spp.</i>	snowcap	D	F
rhodie max (rosebay)	<i>Rhododendron maximum</i>		D	F
rose †	<i>Rosa banksiae</i>	luta	D	F
rose, knockout	<i>Rosa spp. hybrid</i>	knockout	D	F

shrub				
rosemary †	<i>Rosmarinus officinalis</i>		O,D	C,F
rosemary, bog	<i>Andromeda polifolia</i>	nana	D	F
salvia	<i>Salvia farinacea</i>	rhea	D	F
smoketree, royal purple	<i>Cotinus coggryia</i>	royal purple	D	F
sedum	<i>Sedum spurium</i>	dragon blood red red carpet yellow	O,D	C,F
snapdragon	<i>Antirrhinum</i> spp.		D D D	F F F
sourwood	<i>Oxydendrum arboreum</i>		D	F
spiraea	<i>Astilbe X arendsii</i>	fanall	D	F
spiraea	<i>Spiraea</i> spp.	Anthony waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white	D	F
spiraea, garland	<i>Spiraea</i> spp.		D D D D D	F F F F F
spruce, black hills	<i>Picea glauca</i>		D	F
spruce, Colorado blue	<i>Picea pungens</i>	glauca	D	F
spruce, Norway	<i>Picea abies</i>		O,D	C,F
spruce, white	<i>Picea glauca</i>	conica	D	F
sweetflag, grassy-le			D	F
sweetgum	<i>Liquidambar styraciflua</i>		D	F
sycamore	<i>Platanus occidentalis</i>		D	F
tree fern (tiki fern)	<i>Asparagus virgatus</i>		D	F
trumpet flower, evening	<i>Gelsemium sempervirens</i>		D	F
tulip	<i>Tulip</i> , spp	apeldoorn	D	F
tufted hairgrass	<i>Deschampsia caespitosa</i>		D	F
verbena, shrub	<i>Lantana sellowiana</i>		D	F
vervain	<i>Verbena</i> spp.	St. Paul	D	F
viburnum	<i>Viburnum</i> spp.	opulus 'sterile'	D	F

27/08

		American cranberry bush arrowood common snowball European cranberry bush linden mohican Wright		
vinca (periwinkle)	<i>Vinca minor</i>		O,D D D D D D D	C,F F F F F F F
windmill palm	<i>Trachycarpus fortunei</i>		D	F
yarrow	<i>Achillea</i> spp.		D	F
xylosma	<i>Xylosma congestum</i>		D	F
yaupon	<i>Ilex vomitoria</i>	dwarf	D	F
yew	<i>Taxus cuspidata</i> <i>Taxus cuspidata</i> <i>Taxus X media</i>	capitata denisiformis	D	F
			O,D D D	C,F F F

¹ Directed means with the spray mixture directed to the soil around and under an ornamental plant minimizing contact with the plant's foliage.

[†] Ornamental species only. DO NOT USE ON FOOD-PRODUCING TREES AND PLANTS.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

28/28

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

EPA accepted: __/__/__