

62719-447

12-10-2007

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AND

OFFICE OF
PREVENTION, PESTICIDES
TOXIC SUBSTANCES

12-10-07

Dow AgroSciences
9330 Zionsville Road
Indianapolis, IN 46268-1054

Subject: EPA Reg. 62719-424/Goal 2XL Label Amendment

The above mentioned labeling is acceptable provided you make the following changes to the label:

1. Remove "Recommended" from the label.
2. To the label add "Do not apply more than 0.5 lb. ai per year."

If you have any questions please call Erik Kraft at 703-308-9358.

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft", written over a horizontal line.

Erik Kraft
Acting Product Manager 23
Herbicide Branch
Registration Division (7505P)

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Goal[®] 2XL

EPA Reg. No. 62719-424

For Weed Control In Fallow Beds To Be Planted To Field Corn

(For Distribution and Use Only Within the State of California)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Goal[®] 2XL herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Goal 2XL according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Goal 2XL.

Directions for Use

General Information

Goal[®] 2XL herbicide is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate, paraquat or other registered postemergence herbicides for the control of annual broadleaf weeds in fallow beds to be planted to field corn.

Before planting field corn, treated soil must be thoroughly mixed to a depth of at least 2.5 inches. Failure to achieve thorough soil mixing may lead to crop injury and/or stand loss. Weed control should not be expected after soil incorporation.

At least 0.25 inch of irrigation or rainfall is recommended to activate Goal 2XL and should occur within 3 to 4 weeks after application.

A tank mix with glyphosate, paraquat or another registered postemergence herbicide is recommended if the treatment area contains dense weed populations, oversized weed seedlings, volunteer grains or annual grasses.

Refer to the main product label for a list of weeds controlled, mixing directions and other details.

Exercise extreme care to avoid herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result.

Goal 2XL Used Alone: Goal 2XL may be applied at 1 to 2 pints (0.25 to 0.5 lb. active) per acre. The lower rate (1.0 pint per acre) should provide up to four weeks of preemergence control of susceptible weeds and provide postemergence control of susceptible weeds up to four-leaf stage. The higher rate (2.0 pints per acre) should provide preemergence control of susceptible weeds for up to eight weeks and postemergence control of susceptible weeds up to six leaf-stage.

ACCEPTED W/comm

12-10-07

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

62719-424

2/3

Tank Mixes with Goal 2XL: When tank mixing, always read and follow all individual manufacturer's labels. In interpreting all labels for the tank mix, the most restrictive situations must apply. Goal 2XL can be tank mixed with glyphosate, paraquat or other registered postemergence herbicides for postemergence control of annual weed grasses; volunteer grains and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lb. active) of Goal 2XL with labeled rates of glyphosate, paraquat or other registered postemergence products. Apply at the recommended rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

Ground Application: Goal 2XL should be applied in a minimum of 20 gallons of water per acre. The volume of water used should be increased as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

Aerial application: Goal 2XL may be aerially applied for weed control on fallow beds. Follow requirements for aerial application in the general information section of the main product label.

Avoid Drift: When applying to fallow beds, extreme care must be exercised to prevent spray drift, which could result in damage to other crops or desirable vegetation.

Crop injury may be enhanced if newly seeded crops are under stress due to drought, flooding, excessive fertilizer or soil salts, low soil temperatures, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.

Restrictions

- Follow all applicable use directions, precautions, restrictions, and Worker Protection Standard requirements on the EPA registered label.
- **Maximum application rate:** Do not apply more than 2 pints (0.5 lb active ingredient) per acre, per fallow season. Do not apply more than 0.5 lb of oxyfluorfen active ingredient per acre per fallow season as a result of single or multiple applications of this or other oxyfluorfen formulations.
- Do not apply Goal 2XL within 60 days prior to planting of field corn.
- Before planting field corn, treated soil must be thoroughly mixed to a depth of at least 2.5 inches.
- **Chemigation:** For uses described in the supplemental labeling, do not apply this product through any type of irrigation system.
- Do not use on sweet corn.
- A 7-day interval from treatment to planting is specified; the use of any plants from a treated field for green chop, ensilage, forage or fodder or the feeding or grazing of animals on any treated area is prohibited; PHI is 60 days.
- Application may be made in a minimum of 20 gal of water/A using ground equipment or 5 gal/A by air. Applications may be made alone or as a tank mix with other herbicides.

®Trademark of Dow AgroSciences LLC

R843-087

EPA accepted: __/__/__

Initial printing.