



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Ray S. Brinkmeyer
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

FEB 29 2008

Dear Mr. Brinkmeyer:

Subject: Eagle 0.62G
EPA Reg. No. 62719-462
Your Submission of October 15, 2007

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. A stamped copy of the label is enclosed for your records

Submit one copy of the final printed label before you release the product for shipment.

Sincerely yours,

A handwritten signature in cursive script that reads "Carl Grable for".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

7505P:CGrable:cg:2/27/08

Eagle[®] 0.62G

EPA Reg. No. 62719-462

The master label for Eagle 0.62G consists of separate, self-contained sub labels for professional use and residential use, respectively.

Registration Notes:

Source label text based on EPA-accepted copy dated October 24, 2006.

Proposed changes by amendment:

1. Restructured label into master label format with separate labels for professional and residential uses.
2. Ingredients statement: (1) Changed "inert" to "other"; (2) revised statement "Contains 0.31 lb of myclobutanil per 50 lb bag."
3. Precautionary Statements, Personal Protective Equipment and First Aid Statements: Revised these statements based upon the following acute tox categories:

Acute Toxicity Category by Route of Exposure:

Route of Exposure	Eagle 0.62G	MRID No.
Oral LD ₅₀	Cat IV	46886701
Dermal LD ₅₀	Cat IV	46886702
Inhalation	Cat IV	current label
Ocular Irritation	Cat III	46886703
Dermal Irritation	Cat IV	46886704
Dermal Sensitization	Non-sensitizer	46886705

4. Revised referral statement on base label to read "Refer to label booklet for Directions for Use."
5. Added "Produced for" before registrant name and address.
6. Storage and Disposal: Added section to base label.
7. Directions for Use: Added "Read all directions carefully before applying this product." directly underneath heading.
8. Storage and Disposal: Revised section per Chap 13 of the Label Review Manual and Federal Register Notice dated August 16, 2006.
9. Added gray leafspot to list of diseases in table.


Sub Label A: Professional Use

10. Added resistance management language from PR Notice 2001-5.
11. Personal Protective Equipment: Revised PPE for what is required for WPS uses and what is required for non-WPS uses.
12. Added Agricultural Use Requirements box to base label and label booklet cover.
13. Non-Agricultural Use Requirements: Revised section based upon Chap 10 of the Label Review Manual.
14. General Information: Revised section for consistency with other DAS labels.
15. Application Equipment and Application Techniques: Added heading and revised section for consistency with other DAS labels.
16. Application Timing: Added heading and revised section for consistency with other DAS labels.
17. Broadcast Application Rate: Added heading and revised section for consistency with other DAS labels.

Sub Label B: Residential Use

18. Revised sale copy.
19. Directions for Use: (1) Added exclusionary statement "Not for use on turfgrass being...or for research purposes."; (2) deleted "Do not apply this product...agency responsible for pesticide regulation."
20. How This Product Works: Changed heading from "General Information" and revised section for consistency with other DAS labels.

4 8 32



62719-462

[Sub Label A: Professional Uses]

(Base label for rigid containers 5 gal or less):

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

Group	3	FUNGICIDE
--------------	----------	------------------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-
1-H-1,2,4 triazole-propanenitrile..... 0.62%

Other Ingredients 99.38%

Total 100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use.** If terms are unacceptable, return at once unopened.

6 7 32

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight __

(Base label for nonrigid containers any size):

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

Group	3	FUNGICIDE
--------------	----------	------------------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile.....	0.62%
Other Ingredients	99.38%
Total	100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

8 8 32

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight __

9 8 32

(Base label for refillable rigid containers greater than 5 gal):

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile.....	0.62%
Other Ingredients	99.38%
Total	100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

10 8 32

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

11 8 32

A7A / Eagle 0.62G / MSTR Amend / 09-10-07

Page 8

®Trademark of Dow AgroSciences LLC
Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight __

(Base label for nonrefillable rigid containers larger than 5 gal):

Eagle[®] 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

Group	3	FUNGICIDE
--------------	----------	------------------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-
1-H-1,2,4 triazole-propanenitrile..... 0.62%

Other Ingredients 99.38%

Total 100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

13 8 32

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

14 8 32

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight _____

(Label booklet cover):

Eagle[®] 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile.....	0.62%
Other Ingredients	99.38%
Total	100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%

Keep Out of Reach of Children

CAUTION

<p>Agricultural Use Requirements Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.</p>

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC
Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight __

16 8 32

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Causes Moderate Eye Irritation**

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

17 8 32

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made from any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until dusts have settled.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this

18 8 32

container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

Eagle® 0.62G specialty fungicide is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established turfgrass. Examples of such sites include, but are not limited to: residential and commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, golf courses (fairways, roughs, tees, greens), and sod farms. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program.

Use Eagle 0.62G in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

General Use Precautions

Fungicide Resistance Management

Eagle 0.62G belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Eagle 0.62G should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Application Equipment and Application Techniques

Apply Eagle 0.62G uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

Application Timing

Reduce the interval between applications of Eagle 0.62G when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of Eagle 0.62G and shorter application interval. Under light to moderate disease pressure, apply Eagle 0.62G at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F, apply only to dry foliage.

Broadcast Application Rate

In non-residential turfgrass, optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 to 5 lb per 1000 sq ft. In residential turfgrass, optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 lb per 1000 sq ft. See the following table for specific application rates for various diseases. See the table below for specific application rates for various diseases.

[Supplemental distributor should include the following bullet if product is formulated on fertilizer carrier.]

Do not adjust application rates to meet fertilizer requirements.

[Supplemental distributor may include appropriate spreader equipment, suggested calibration settings and/or pictures showing proper application techniques.]

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)		
	2.5	3.5	5
XXXX	X	X	X

Eagle 0.62G covers approximately 9500 sq ft at the application rate of 1 lb of nitrogen (5.3 lb of fertilizer) per 1000 sq ft.

Non-Residential Turfgrass¹

Diseases	Eagle 0.62G (lb/1000 sq ft)	Application Interval/Timing (Days)	Use Recommendations	Restrictions
anthracnose red thread septoria leaf spot	2.5	14 - 21	Apply when conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not apply more than 29 lb of Eagle 0.62G (0.14 lb myclobutanil) per 1000 sq ft per year. For Nassau and Suffolk counties in New York State, use of this product is limited to 7.2 lb of Eagle 0.62G per 1000
brown patch	2.5 - 5	14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	
copper spot zonate leaf spot	2.5		Apply when conditions are favorable for disease development.	

crown rot leaf spot melting-out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.
dollar spot	2 - 5	14 - 28	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.
fusarium blight	2.5 - 5	14 - 21	Apply when conditions are favorable for disease development.
fusarium patch (pink snow mold)	2.5 - 5	fall - winter	Apply prior to snow cover.
gray leafspot	2.5 - 5	14	Apply when conditions are favorable for disease development.
leaf smuts	2.5		Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.
necrotic ring spot	2.5 - 5	spring: 28	Make applications on a preventative basis in early to mid-spring.
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant. Apply 5 lb per 1000 sq ft followed by a second application one month later.
powdery mildew rusts	2.5	14 - 28	Apply when conditions are favorable for disease development.
summer patch	2.5 - 5		Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.

sq ft per year (1.95 lb myclobutanil per acre).

take-all patch	5	spring/fall: 28 days	To reduce the severity, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F and 1 to 2 spring applications in April and May depending upon local recommendations.
zoysia large patch	5	fall: 28 days	Make applications in fall before turfgrass becomes dormant. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.

¹Including, but not limited to: commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, golf courses (fairways, roughs, tees, greens), and sod farms.

Residential Turfgrass

Diseases	Eagle 0.62G (lb/1000 sq ft)	Application Interval/Timing (Days)	Use Recommendations	Restrictions
anthracnose red thread septoria leaf spot	2.5	14 - 21	Apply when conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not apply more than 29 lb of Eagle 0.62G per 1000 sq ft per year. For Nassau and Suffolk counties in New York State, use of this product is limited to 7.2 lb of Eagle 0.62G per 1000 sq ft per year.
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	
crown rot leaf spot melting-out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	
dollar spot		14 - 21	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight			Apply when conditions are favorable for disease development.	

fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.
gray leafspot		14	Apply when conditions are favorable for disease development.
leaf smuts			Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.
summer patch			Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted: __/__/__

[Sub Label B: Residential Uses]

(Base label):

Eagle[®] 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established home lawns and in ornamental turfgrass

and/or
Provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot

and/or
Provides controlled release nitrogen feeding

and/or
For control of turfgrass diseases on home lawns

and/or
Systemic [action] [formula] [protectant]

and/or
Cures up to _____ sq ft

and/or
Treats up to _____ sq ft

and/or
Cures and prevents all major lawn diseases [including brown patch, dollar spot, and other listed lawn diseases]

and/or
Prevents over 10 major lawn diseases

and/or
[One application] Protects for up to 4 weeks

and/or
Use throughout the entire year to prevent [and control] disease

and/or
Rainproof – once absorbed, can't wash off

and/or
Controls [anthracnose], [brown spot], [copper spot], [dollar spot], [fusarium blight], [fusarium patch], [crown rot], [leaf smuts], [necrotic ring spot], [powdery mildew], [summer patch] and more

and/or
Do-it-yourself disease control

and/or
For use anytime [during the year] to prevent and control lawn disease[s]

and/or
Use anytime to prevent and control lawn disease[s]

and/or
For use on all types of home lawns [and turfgrass]

and/or
Easy [spreader] application

and/or
Can be used on all turfgrass species

and/or
Ready-to-spread granules

and/or

Up to six applications can be made per year
and/or

For more information on complete line of [lawn and garden] products, visit our web site at
www.dowagro.com

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile.....	0.62%
Other Ingredients	99.38%
Total	100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for
fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Total Nitrogen (N).....	X.XX%
Ammoniacal Nitrogen.....	X.XX%
Urea Nitrogen	X.XX%
Available Phosphate (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O)	X.XX%
Sulfur (S).....	X.XX%
Free Sulfur (S).....	X.XX%
Chlorine (Cl) Max	X.XX%

May include the following:

Poly coated sulfur coated urea, urea, ammonium phosphate, muriate of potash.

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

**Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes and socks.
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or
using tobacco.**

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact
lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or
doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for
treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms
in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to inter-

26 7 32

tidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight _____

(Label booklet cover):

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established home lawns and in ornamental turfgrass

- and/or
- Provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot**
- and/or
- Provides controlled release nitrogen feeding**
- and/or
- For control of turfgrass diseases on home lawns**
- and/or
- Systemic [action] [formula] [protectant]**
- and/or
- Cures up to _____ sq ft**
- and/or
- Treats up to _____ sq ft**
- and/or
- Cures and prevents all major lawn diseases [including brown patch, dollar spot, and other listed lawn diseases]**
- and/or
- Prevents over 10 major lawn diseases**
- and/or
- [One application] Protects for up to 4 weeks**
- and/or
- Use throughout the entire year to prevent [and control] disease**
- and/or
- Rainproof – once absorbed, can't wash off**
- and/or
- Controls [anthracnose], [brown spot], [copper spot], [dollar spot], [fusarium blight], [fusarium patch], [crown rot], [leaf smuts], [necrotic ring spot], [powdery mildew], [summer patch] and more**
- and/or
- Do-it-yourself disease control**
- and/or
- For use anytime [during the year] to prevent and control lawn disease[s]**
- and/or
- Use anytime to prevent and control lawn disease[s]**
- and/or
- For use on all types of home lawns [and turfgrass]**
- and/or
- Easy [spreader] application**
- and/or
- Can be used on all turfgrass species**
- and/or
- Ready-to-spread granules**
- and/or
- Up to six applications can be made per year**
- and/or

For more information on complete line of [lawn and garden] products, visit our web site at www.dowagro.com

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile.....	0.62%
Other Ingredients	99.38%
Total	100.00%

Contains 0.31 lb of myclobutanil per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Total Nitrogen (N).....	X.XX%
Ammoniacal Nitrogen.....	X.XX%
Urea Nitrogen.....	X.XX%
Available Phosphate (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%
Sulfur (S).....	X.XX%
Free Sulfur (S).....	X.XX%
Chlorine (Cl) Max.....	X.XX%

May include the following:

Poly coated sulfur coated urea, urea, ammonium phosphate, muriate of potash.

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Net Weight _____

29 8 32

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Causes Moderate Eye Irritation**

Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on turfgrass being grown for sale or other commercial use as sod, or for commercial seed productions, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

How This Product Works

Eagle® 0.62G specialty fungicide is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established home lawns and ornamental turfgrass. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program.

Use Eagle 0.62G in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

How to Apply

Apply Eagle 0.62G uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

When to Apply

Reduce the interval between applications of Eagle 0.62G when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of Eagle 0.62G and shorter application interval. Under light to moderate disease pressure, apply Eagle 0.62G at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F; apply only to dry foliage.

How Much to Apply

Optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 lb per 1000 sq ft. See the table below for specific application rates for various diseases.

[Supplemental distributor should include the following if product is formulated on fertilizer.]
Do not adjust application rates to meet fertilizer requirements.

[Supplemental distributor may include appropriate spreader equipment, suggested calibration settings and/or pictures showing proper application techniques.]

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)		
	2.5	3.5	5
XXXX	X	X	X

Diseases	Eagle 0.62G (lb/1000 sq ft)	Application Interval/Timing (Days)	Use Recommendations	Restrictions
anthracnose red thread septoria leaf spot brown patch	2.5	14 - 21	Apply when conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not apply more than 29 lb of Eagle 0.62G per 1000 sq ft per year. For Nassau and Suffolk counties in New York State, use of this product is limited to 7.2 lb of Eagle 0.62G per 1000 sq ft per year.
		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	

crown rot leaf spot melting-out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.
dollar spot		14 - 21	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.
fusarium blight		14 - 21	Apply when conditions are favorable for disease development.
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.
gray leafspot		14	Apply when conditions are favorable for disease development.
leaf smuts		14	Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.
summer patch			Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted: __/__/__