

62719-462

03-12-2010

1/29



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms Cindy Loy
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 12 2010

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated December 9, 2009 for:

EPA Registration 62719-462

Eagle 0.62G

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number Dow AgroSciences LLC/62719-462	2. EPA Product Manager Shaja B. Joyner	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/Eagle 0.62G	PM# 20	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
MAR 12 2010

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container
1 lb, 50 lb, 500 lb

5. Location of Label Directions
 On label

6. Manner in Which Label is Affixed to Product
 Lithograph Paper glued Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Raymond S. Brinkmeyer, Ph.D.	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4694 (fax: 317-337-4649)
--------------------------------------	-----------------------------	-------------------------------------------------------------------------

Certification
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Regulatory Manager	6. Date Application Received (Stamped)
4. Typed Name Raymond S. Brinkmeyer, Ph.D. (rsbrinkmeyer@dow.com)	5. Date January 28, 2010	

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

308/2E
December 9, 2009



Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U. S. Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

EAGLE 0.62G (A.I. MYCLOBUTANIL)
EPA REGISTRATION NUMBER: 62719-462
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICE 2007-4

Enclosed please find labeling for the notification action of Eagle® 0.62G specialty fungicide. The following change has been made by notification:

- I. Storage and Disposal: Revised section per PR Notice 2007-4.

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Contents of Submission


- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled Eagle 0.62G (A7A/Eagle 0.62G/MSTR Notif/12-09-09) (25 Pages plus Registration Notes) (1 Copy)

If you require further information, please contact Cindy Loy, Regulatory Specialist at (317) 337-4655.

Sincerely,

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
(317) 337-4694
(317) 337-4649 (FAX)

Enclosures



Eagle[®] 0.62G

EPA Reg. No. 62719-462

The master label for Eagle 0.62G consists of separate, self-contained sub labels for professional use and residential use, respectively.


Registration Notes:

Source label text based on EPA-accepted copy dated February 29, 2008 and notification coded "A7A / Eagle 0.62G / MSTR Notif / 07-31-08."

Change by notification:

1. Storage and Disposal: Revised section per PR Notice 2007-4.

®Trademark of Dow AgroSciences LLC



5
29

[Sub Label A: Professional Uses]

NOTIFICATION

(Base label):

MAR 12 2010

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

[Editor's note: Supplemental distributor will add one of these statements if product is formulated with fertilizer.]

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-
1-H-1,2,4 triazole-propanenitrile 0.62%

Other Ingredients 99.38%

Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation


Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT



covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

(Storage and Disposal for rigid containers 5 gal or less)**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

7
29

(Storage and Disposal for nonrigid containers any size)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

1
29

(Label booklet cover):

Eagle[®] 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass

[Editor's note: Supplemental distributor will add one of these statements if product is formulated with fertilizer.]

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-

1-H-1,2,4 triazole-propanenitrile 0.62%

Other Ingredients 99.38%

Total 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N) X.XX%

Available Phosphoric Acid (P₂O₅) X.XX%

Soluble Potash (K₂O) X.XX%

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

®Trademark of Dow AgroSciences LLC


Produced for

EPA Est. _____

12
29

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Weight __



(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Causes Moderate Eye Irritation**

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear

- Long-sleeved shirt and long pants
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

12
29

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made from any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until dusts have settled.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this

container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Information

Eagle® 0.62G specialty fungicide is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established turfgrass. Examples of such sites include, but are not limited to: residential and commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, golf courses (fairways, roughs, tees, greens), and sod farms. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program.

Use Eagle 0.62G in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

General Use Precautions

Fungicide Resistance Management

Eagle 0.62G belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Eagle 0.62G should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if product is formulated with fertilizer and is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.

14
29

- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.
- Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Application Equipment and Application Techniques

Apply Eagle 0.62G uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

Application Timing

Reduce the interval between applications of Eagle 0.62G when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of Eagle 0.62G and shorter application interval. Under light to moderate disease pressure, apply Eagle 0.62G at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F, apply only to dry foliage.

Broadcast Application Rate


In non-residential turfgrass, optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 to 5 lb per 1000 sq ft. In residential turfgrass, optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 lb per 1000 sq ft. See the following table for specific application rates for various diseases.

[Supplemental distributor should include the following bullet if product is formulated on fertilizer carrier.]

Do not adjust application rates to meet fertilizer requirements.

[Supplemental distributor may include appropriate spreader equipment, suggested calibration settings and/or pictures showing proper application techniques.]

Spreader Settings as a Guide for Calibration



16
29

		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant. Apply 5 lb per 1000 sq ft followed by a second application one month later.
powdery mildew rusts	2.5	14 - 28	Apply when conditions are favorable for disease development.
summer patch	2.5 - 5		Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.
take-all patch	5	spring/fall: 28	To reduce the severity, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F and 1 to 2 spring applications in April and May depending upon local recommendations.
zoysia large patch	5	fall: 28	Make applications in fall before turfgrass becomes dormant. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.

¹Including, but not limited to: commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, golf courses (fairways, roughs, tees, greens), and sod farms.

Residential Turfgrass

Diseases	Eagle 0.62G (lb/1000 sq ft)	Application Interval/Timing (Days)	Use Recommendations	Restrictions
anthracnose red thread septoria leaf spot	2.5	14 - 21	Apply when conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not apply more than 29 lb of Eagle 0.62G per 1000 sq ft per year. For Nassau and Suffolk counties in New York State, use
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	

CCCC
CCCC
CCCC
CCCC

17
29

copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	of this product is limited to 7.2 lb of Eagle 0.62G per 1000 sq ft per year.
crown rot leaf spot melting-out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	
dollar spot		14 - 21	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight			Apply when conditions are favorable for disease development.	
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	
gray leafspot		14	Apply when conditions are favorable for disease development.	
leaf smuts			Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.	
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.	
summer patch			Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	

18
29

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies


To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted 02/29/08



19
29

[Sub Label B: Residential Uses]

(Base label):

Eagle® 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established home lawns and in ornamental turfgrass

[Editor's note: Supplemental distributor will this statement if product is formulated with fertilizer and the product is registered in Florida.]

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

[Editor's note: Supplemental distributor will add one of these statements if product is formulated with fertilizer.]

**Cures up to _____ sq ft
and/or**

Treats up to _____ sq ft

**Provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot
and/or**

**Provides controlled release nitrogen feeding
and/or**

**For control of turfgrass diseases on home lawns
and/or**

**Systemic [action] [formula] [protectant]
and/or
and/or**

**Cures and prevents all major lawn diseases [including brown patch, dollar spot, and other listed lawn diseases]
and/or**

**Prevents over 10 major lawn diseases
and/or**

**[One application] Protects for up to 4 weeks
and/or**

**Use throughout the entire year to prevent [and control] disease
and/or**


**Rainproof – once absorbed, can't wash off
and/or**

**Controls [anthracnose], [brown spot], [copper spot], [dollar spot], [fusarium blight], [fusarium patch], [crown rot], [leaf smuts], [necrotic ring spot], [powdery mildew], [summer patch] and more
and/or**

**Do-it-yourself disease control
and/or**

**For use anytime [during the year] to prevent and control lawn disease[s]
and/or**

**Use anytime to prevent and control lawn disease[s]
and/or**



29
29

For use on all types of home lawns [and turfgrass]

and/or

Easy [spreader] application

and/or

Can be used on all turfgrass species

and/or

Ready-to-spread granules

and/or

Up to six applications can be made per year

and/or

For more information on complete line of [lawn and garden] products, visit our web site at www.dowagro.com

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-
1-H-1,2,4 triazole-propanenitrile 0.62%

Other Ingredients 99.38%

Total..... 100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Total Nitrogen (N)	X.XX%
Ammoniacal Nitrogen	X.XX%
Urea Nitrogen	X.XX%
Available Phosphate (P ₂ O ₅).....	X.XX%
Soluble Potash (K ₂ O).....	X.XX%
Sulfur (S).....	X.XX%
Free Sulfur (S)	X.XX%
Chlorine (Cl) Max.....	X.XX%

May include the following:

Poly coated sulfur coated urea, urea, ammonium phosphate, muriate of potash.

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation

Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for

21
29

treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.
Container Handling: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.


EPA Reg. No. 62719-462

EPA Est. _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Net Weight __



22
29

(Label booklet cover):

Eagle[®] 0.62G

Specialty Fungicide

A granular [fertilizer] product containing a systemic protectant and curative fungicide for the control of diseases in established home lawns and in ornamental turfgrass

[Editor's note: Supplemental distributor will this statement if product is formulated with fertilizer and the product is registered in Florida.]

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

[Editor's note: Supplemental distributor will add one of these statements if product is formulated with fertilizer.]

Cures up to _____ sq ft
and/or

Treats up to _____ sq ft

Provides systemic prevention and control of turfgrass diseases including benzimidazole tolerant dollar spot
and/or

Provides controlled release nitrogen feeding
and/or

For control of turfgrass diseases on home lawns
and/or

Systemic [action] [formula] [protectant]
and/or

Cures and prevents all major lawn diseases [including brown patch, dollar spot, and other listed lawn diseases]
and/or

Prevents over 10 major lawn diseases
and/or

[One application] Protects for up to 4 weeks
and/or

Use throughout the entire year to prevent [and control] disease
and/or

Rainproof – once absorbed, can't wash off
and/or


Controls [anthracnose], [brown spot], [copper spot], [dollar spot], [fusarium blight], [fusarium patch], [crown rot], [leaf smuts], [necrotic ring spot], [powdery mildew], [summer patch] and more
and/or

Do-it-yourself disease control
and/or

For use anytime [during the year] to prevent and control lawn disease[s]
and/or

Use anytime to prevent and control lawn disease[s]
and/or

For use on all types of home lawns [and turfgrass]
and/or



23
29

Easy [spreader] application

and/or

Can be used on all turfgrass species

and/or

Ready-to-spread granules

and/or

Up to six applications can be made per year

and/or

For more information on complete line of [lawn and garden] products, visit our web site at www.dowagro.com

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)- 1-H-1,2,4 triazole-propanenitrile	0.62%
Other Ingredients	99.38%
Total	100.00%

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used if product is formulated with fertilizer.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Total Nitrogen (N)	X.XX%
Ammoniacal Nitrogen	X.XX%
Urea Nitrogen	X.XX%
Available Phosphate (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%
Sulfur (S)	X.XX%
Free Sulfur (S)	X.XX%
Chlorine (Cl) Max	X.XX%

May include the following:

Poly coated sulfur coated urea, urea, ammonium phosphate, muriate of potash.

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**


In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-462

®Trademark of Dow AgroSciences LLC


**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**



24

29

Net Weight __



25
29

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION****Causes Moderate Eye Irritation**

Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This pesticide is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on turfgrass being grown for sale or other commercial use as sod, or for commercial seed productions, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

How This Product Works

Eagle® 0.62G specialty fungicide is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established home lawns and ornamental turfgrass. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program.

Use Eagle 0.62G in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing

26
29

the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

How to Apply

Apply Eagle 0.62G uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

When to Apply

Reduce the interval between applications of Eagle 0.62G when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of Eagle 0.62G and shorter application interval. Under light to moderate disease pressure, apply Eagle 0.62G at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F; apply only to dray foliage.

How Much to Apply

Optimum disease control is achieved when Eagle 0.62G is applied in a preventative disease control program at a rate of 2.5 lb per 1000 sq ft. See the table below for specific application rates for various diseases.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if product is formulated with fertilizer and is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

[Supplemental distributor should include the following if product is formulated on fertilizer.]

Do not adjust application rates to meet fertilizer requirements.

[Supplemental distributor may include appropriate spreader equipment, suggested calibration settings and/or pictures showing proper application techniques.]

Spreader Settings as a Guide for Calibration

	Calibration Settings at Specified Rate (lb/1000 sq ft)
--	---------------------------------------------------------------

27
29

Brand and Model of Spreader	2.5	3.5	5
XXXX	X	X	X

Diseases	Eagle 0.62G (lb/1000 sq ft)	Application Interval/Timing (Days)	Use Recommendations	Restrictions
anthracnose red thread septoria leaf spot	2.5	14 - 21	Apply when conditions are favorable for disease development.	<ul style="list-style-type: none"> Do not apply more than 29 lb of Eagle 0.62G per 1000 sq ft per year. For Nassau and Suffolk counties in New York State, use of this product is limited to 7.2 lb of Eagle 0.62G per 1000 sq ft per year.
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	
crown rot leaf spot melting-out			Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	
dollar spot		14 - 21	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight		14 - 21	Apply when conditions are favorable for disease development.	
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	
gray leafspot		14	Apply when conditions are favorable for disease development.	
leaf smuts			Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.	
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	

28
29

powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.	
summer patch			Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

29
29

®Trademark of Dow AgroSciences LLC
EPA accepted 02/29/08

