

62719-473

6/18/2002

1/1

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

ACCEPTED
JUN 18 2002
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-473

MACH 2* 1.5G

Dow AgroSciences EPA Reg. No. 62719-473
RohMid EPA Reg. No. 69075-4

A Single Application at Maximum Labeled Use Rate for Optimum Control of White Grubs on Turfgrass

- ATTENTION**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 - This labeling must be in the possession of the user at the time of application.
 - Read the label affixed to the container for MACH 2* 1.5G specialty insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
 - Use of MACH 2 1.5G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for MACH 2 1.5G.

Directions for Use

Refer to product label for General Use Precautions and Application Instructions.

Pest	Amount of MACH 2 1.5G	Comments
White grub larvae such as: Japanese beetle, <i>Popillia japonica</i> Northern masked chafer, <i>Cyclocephala borealis</i> , Southern masked chafer, <i>Cyclocephala lurida</i> , May/June beetle, <i>Phyllophaga</i> spp., Black turfgrass ataenius, <i>Ataenius spretulus</i> , Green June beetle, <i>Cotinus nitida</i> , Annual bluegrass weevil larvae, <i>Hyperodes</i> spp., Billbugs, <i>Sphenophorus</i> spp., Aphodius beetle, <i>Aphodius</i> spp., European chafer, <i>Rhizotrogus majalis</i> , Oriental beetle, <i>Exomala orientalis</i>	133 lb/acre (3 lb/1000 sq ft)	MACH 2 1.5G may be used as either a preventative or an early curative treatment (see application timing instructions). Make only one application.

Restrictions:

- Do not apply more than 133 lb (2 lb active ingredient) per acre per year regardless of pests controlled.
- Use of this product on "Tifdwarf" Bermudagrass greens may result in short term discoloration of the turfgrass. Since many agronomic factors may influence this response it is recommended that users treat a small area at recommended rates prior to initiating large-scale use.
- **For Sod Farms:** Allow at least 7 days to elapse between application and harvest of sod.

*Trademark of Dow AgroSciences LLC D06-158-002 EPA-accepted 03/21/02 Initial printing.	
---	--