

62719-487

09/29/2004

1/18

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 05-31-98



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number: Dow AgroSciences/62719-487
2. EPA Product Manager: Joanne I. Miller (7505C)
3. Proposed Classification: [X] None [ ] Restricted
4. Company/Product (Name): Dow AgroSciences/Crab-Buster™ 0.008% FG
PM#: PM / 23
5. Name and Address of Applicant: Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268
6. Expedited Review: EPA Reg. No., Product Name

Section - II

Amendment - Explain below. Resubmission in response to Agency letter dated. Notification - Explain below. Final printed labels in response to Agency letter dated. "Me Too" Application. Other - Explain below. NOTIFICATION SEP 29 2004

Explanation: Use additional page(s) if necessary. Labeling notification based on EPA-accepted copy (Notice of Registration) dated March 24, 2003 and notification coded "G2B/Crab-Buster 0.08%% FG/ Notification With Edits/06-17-04".

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging, 2. Type of Container: Metal, Plastic, Glass, Paper, Other. 3. Location of Net Contents Information: Label, Container. 4. Size(s) Retail Container. 5. Location of Label Directions: On Label, On Labeling accompanying product. 6. Manner in Which Label is Affixed to Product: Lithograph, Paper glued, Stenciled, Other.

Section - IV

1. Contact Point /Complete items directly below for identification of individual to be contacted, if necessary, to process this application. Name: Kimberly S. Gilbert, Title: Regulatory Manager, Telephone No. (317) 337-468; fax no. (317) 337-4649

Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. 2. Signature: Kimberly S. Gilbert, 3. Title: Regulatory Manager, 4. Typed Name: RM: Kimberly S. Gilbert, 5. Date: September 3, 2004, 8. Date Application Received (Stamped)



[Front Panel of Bag]

(Logo) Dow AgroSciences

NOTIFICATION

SEP 29 2004

# Crab-Buster™ 0.08% FG

## Lawn and Ornamental Herbicide

For control of crabgrass and control or suppression many other annual grass and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf.

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

*In the state of New York, do not exceed 7.2 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).*

### Active Ingredient

dithiopyr: 3,5- pyridinedicarbothioic acid, 2-(difluoromethyl)- 4-(2-methylpropyl)-6- (trifluoromethyl)-S,S-dimethyl ester.....	0.08%
Inert Ingredients .....	99.92%
Total .....	100.00%

This product contains 0.04 pound of the active ingredient dithiopyr per 50 pound bag. Product protected by U.S. Patent No. 4,692,184. Other patents pending

### Guaranteed Fertilizer Analysis: XX-XX-XX †

Water-Insoluble Nitrogen .....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen.....	XX%
Urea Nitrogen.....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Calcium (Ca) .....	XX%
Magnesium (Mg).....	XX%
Copper (Cu) .....	XX%
Iron (Fe) .....	XX%
Manganese (Mn).....	XX%
Zinc (Zn).....	XX%

† This product may be formulated with one or more of the following fertilizer materials: urea, methylene urea, sulfur coated urea, ammoniated phosphates, sulfate of potash, muriate of potash, copper oxide, iron oxide, manganese oxide, zinc oxide or limestone.

**Keep Out of Reach of Children**

**CAUTION**

4/18

**[Front Panel of Bag - Continued]**

**[Optional Label Claims]**

- A Selective Herbicide Plus Fertilizer Combination Product that Provides Control of Crabgrass and many Other Weeds in Established Lawns and Ornamental Turfgrasses and Residential Landscapes.
- **Premium Granular Fertilizer** - Providing the Nutrients Needed for Beautiful Healthy Lawn and Ornamentals.
- Contains **Crab-Buster™ Herbicide** - A Highly Effective Herbicide for Crabgrass Control!
- Can Be Applied **Up to 4 Weeks Later** Than Other Preemergence Crabgrass Preventers.
- This Bag Treats Up To \_\_\_\_\_ Square Feet

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals**

**CAUTION: Causes Eye Irritation • Harmful If Inhaled**

**Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.**

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, the give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

**Environmental Hazards**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

**Refer to back/side panel of bag for Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-487

EPA Est. \_\_\_\_\_

™Trademark of Dow AgroSciences LLC  
**Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.**

**Net Weight \_\_ lb**

**[Back Panel of Bag]****Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Storage and Disposal**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Storage:** Store this product only in its original container in a dry, cool, secured storage area.

**Disposal Instructions:**

**If empty:** Do not reuse this container. Place in trash or offer recycling if available.

**If partially filled:** Call you local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**[Editor's Note: Supplemental distributor may insert own toll free number.]**

**Use Directions for Turf**

This turf fertilizer product, containing Crab-Buster™ herbicide, can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control.

Crab-Buster 0.08% FG provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

<b>Cool-Season Grasses</b>	<b>Warm-Season Grasses</b>
Bentgrass, creeping	Bahiagrass
Bluegrass, Kentucky	Bermudagrass
Fescue, fine	Buffalograss
Fescue, tall	Carpetgrass
Ryegrass, perennial	Centipedegrass
	St. Augustinegrass
	Zoysiagrass

**Use Precautions for Turf**

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

- Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. **Do not** apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

6/18

- **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 14.3 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

#### Other Use Precautions

- **Do not** use clippings from treated turf for mulching around vegetables or fr487 trees.
- **Do not** allow people or pets to enter treated area until dust has settled.

## Application Timing

### Spring Application

**Control of Crabgrass:** Crab-Buster 0.08% FG provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

### Fall Application

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **do not** reseed, overseed or sprig turfgrasses within 12 weeks after application.

### Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

## Weeds Controlled or Suppressed

Crab-Buster 0.08% FG will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

### Grasses

barley  
barnyardgrass  
bluegrass, annual  
brome  
crabgrass, large  
crabgrass, smooth  
crabgrass, Southern  
crowfootgrass †  
dallisgrass  
goosegrass  
foxtail, green  
foxtail, yellow  
kikuyugrass †  
oats, wild  
ryegrass (annual & perennial)  
sandbur  
smutgrass

### Broadleaf Weeds

bittercress †  
carpetweed  
chickweed †  
geranium, Carolina †  
henbit  
knotweed, prostrate  
lespedeza, common †  
marestail  
medic, black  
mustard  
oxalis, buttercup  
pineappleweed  
pigweed, redroot  
parsley-piert †  
purslane, common  
rocket, London  
shepherdspurse  
speedwell, corn †  
spurge, garden  
spurge, prostrate  
spurge, spotted  
woodsorrel, creeping  
woodsorrel, yellow


† Suppression only

## Length of Control

Crab-Buster 0.08% FG will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. **Refer to the map below to determine the region of the U.S. that includes your location.** Use the table below to determine the amount of Crab-Buster 0.08% FG to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.

9/18

**Use "Regions" within the United States:**



**Recommended Use Rates**

Amount of Crab-Buster 0.08% FG to Apply (Pounds of product to apply per 1,000 square feet of turfgrass <sup>†</sup> )			
Region of U.S.	----- Length of Crabgrass Control -----		
	2 - 3 months	3 - 4 months	Early Postemergence Control
North	3.7 lb/1000 sq ft (0.128 lb ai/acre)	5.0 lb/1000 sq ft (0.174 lb ai/acre)	6.3 - 14.3 lb/1000 sq ft <sup>††</sup> (0.22 - 0.5 lb ai/acre)
South	5.5 lb/1000 sq ft (0.192 lb ai/acre)	6.3 - 7.3 lb/1000 sq ft <sup>††</sup> (0.22-0.256 lb ai./acre)	6.3 - 14.3 lb/1000 sq ft <sup>††</sup> (0.22 - 0.5 lb ai/acre)

<sup>†</sup> 1,000 square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.

<sup>††</sup> Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

**Maximum Application Rates:**

- Do not apply more than 14.3 lb of this product per 1,000 sq ft per application and no more than 43.0 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 7.2 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).

**Application Instructions and Equipment**

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.



9/19

[Editor's note: Supplemental Distributor may add a table listing recommended application equipment and calibration settings.]

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)					
	3.7	5.0-	5.5	6.3	7.3	14.3
XXXXXXX	XX	XX	XX	XX	XX	

### Use Directions for Landscape Ornamentals

Crab-Buster 0.08% FG provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. For application instructions and equipment, see Application Instructions and Equipment section above.

### Use Rates

Apply Crab-Buster 0.08% FG with a properly calibrated spreader or other means that will assure uniform distribution. This product may be applied as a single application or as a split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided the maximum application rates per year given below are not exceeded.

Pounds of Product per 1000 Square Feet <sup>b</sup>	Ounces of Product per 100 Square Feet <sup>†</sup>
14.3	1.43 (22.9 oz)

<sup>†</sup> Rate based on an area of 100 sq ft for treatment of small areas.

#### Maximum Application Rates:

- **Do not** apply more than 14.3 lb of this product per 1,000 sq ft per application and no more than 43.0 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).
- In the state of New York, do not exceed 7.2 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).

### Application Timing and Recommendations

Crab-Buster 0.08% FG is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination. Bare ground or mulched areas may be treated. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

## Use Precautions

- Apply this product only to established ornamentals.
- **Do not** apply this product directly to bare roots of ornamental plants as injury may result.
- **Do not** incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- **Do not** apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- **Do not** apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.

## Weeds Controlled or Suppressed

Used as directed, Crab-Buster 0.08% FG will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. **Refer to "Use Directions for Turf" section above for a complete listing of weeds controlled or suppressed.** This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

## Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to label directions. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, it is recommended that a few plants in a limited area be treated and observed for tolerance under local growing conditions prior to large-scale use.

Common Name	Botanical Name	Tolerant Cultivars
Abelia, Dwarf	<i>Abelia X grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i> <i>Ajuga genevensis</i>	Bronze Bronze Beauty
Almond, Flowering	<i>Prunus glandulosa</i>	
Apple <sup>1</sup>	<i>Malus pumila</i>	
Arborvitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus aucuparia</i>	
Ash, Purple	<i>Fraxinus americana</i>	

11/18

Azalea	<i>Rhododendron</i> spp.	Brilliant Buccaneer Carror Chimes (Belgian) Elsie Lee Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow
Barberry, Purple		Atropurpurea
Basket flower	<i>Gaillardia grandiflora</i>	
Bearberry (common)	<i>Arctostaphylos uva-ursi</i>	Massachusetts
Bee Balm	<i>Monarda didyma</i>	
Begonia	<i>Begonia</i> spp.	
Birch, River	<i>Betula nigra</i>	
Blackeyed Susan	<i>Rudbeckia hirta</i>	Goldstrum
Blanket Flower	<i>Gaillardia</i> spp.	
Blueberry <sup>†</sup>	<i>Vaccinium</i> spp.	Bluecrop Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinus</i>	
Boxwood, Japanese		Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus</i> spp.	Moonlight
	<i>Genista pilosa</i>	Vancouver Gold
Bugle Carpet		
Camellia	<i>Camellia japonica</i>  <i>Camellia sasanqua</i>	Debutante Mathotiana Supreme Chansonette
Candy Tuft	<i>Iberis</i> spp.	Snow White
Carex, Variegated	<i>Carex</i>	
Cedar, Red	<i>Juniperus virginiana</i>	
Celosia	<i>Celosia</i> spp.	
Centaurea	<i>Centaurea montana</i>	
Cockscomb, Plumosa	<i>Celosia cristata</i>	Scarlet Plumosa
Coleus	<i>Coleus blumei</i>	Red Kewpie
Columbine	<i>Aquilegia</i> spp.	
Copper leaf	<i>Acalypha wilkesiana</i>	
Coreopsis	<i>Coreopsis</i> spp.	Moonbeam,

12/18

Corn Flower	<i>Centaurea</i> spp.	
Cotoneaster	<i>Cotoneaster apiculatus</i>	
Coyotebrush	<i>Baccharis pilularis</i>	
Cycads	<i>Cycas revoluta</i>	
Cypress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glauca
Cypress, Japanese False	<i>Chamaecyparis obtusa</i>	Gracilis
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Daffodil	<i>Narcissus</i> spp.	King Alfred
Daylily	<i>Hemerocallis</i> spp.	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dianthus (Sweet William)	<i>Dianthus</i> spp.	
Delphinium	<i>Delphinium</i> spp.	Magic fountain
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	Flavarium
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Aureo-marginata Colorata Emerald Gaiety Emerald & N Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fern (various)	<i>Asparagus</i> spp.	
Fescue	<i>Festuca glauca</i>	
Fetterbush	<i>Leucothoe fontanesiana</i>	Rainbow
Ficus	<i>Ficus retusa</i>	Nitidia
Fir Fraser	<i>Abies fraseri</i>	
Forsythia	<i>Forsythia X intermedia</i>  <i>Spring Glory</i>	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Fuchsia	<i>Fuchsia</i> spp.	
Galium	<i>Galium odoratum</i>	
Gardenia	<i>Gardenia jasminoides</i>	Mystery Radicans
Geranium	<i>Pelargonium X hortorum</i>	
Gum	<i>Eucalyptus citriodora</i>	
Hawthorn	<i>Crataegus</i> spp.	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White
Heather, Twisted	<i>Erica cinerea</i>	Mediterranean Pink
Hemlock, Canada	<i>Tsuga canadensis</i>	
Hibiscus	<i>Hibiscus</i> spp.	Blue Bird Brilliant Hula Girl

Holly	<i>Ilex</i> spp. <i>Ilex X meserveae</i>  <i>Ilex X attenuata</i>	Blue Boy Blue Girl Burfodii China Girl Compacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savannah
Holly, Chinese	<i>Ilex cornuta</i>	
Holly, Japanese	<i>Ilex crenata</i>	
Holly, Yaupon	<i>Ilex vomitoria</i>	
Honeysuckle	<i>Lonicera japonica</i>	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta
Hosta	<i>Hosta</i> spp. <i>Hosta lancifolia</i>	Albo Marginata
Ice Plant	<i>Carpobrotus edulis</i>	
Impatiens	<i>Impatiens</i> spp.	
Iris	<i>Iris</i> spp.	Dwarf Blue Wedgewood
Ivy, English	<i>Hedera helix</i>	Bulgaria Thorndale
Jasmine, Asian	<i>Trachelospermum asiaticum</i>	
Juniper	<i>Juniperus</i> spp.	Arcadia Armstrong Bar Harbor
	<i>Juniperus horizontalis</i>	Blue Chip Blue Pacific Blue Rug
	<i>Juniperus horizontalis</i>	Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fr487landi Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blue Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	<i>Juniperus chinensis</i>	Tamariscifolia

	<i>Juniperus sabina</i>	Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Lily, African	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red †	<i>Acer rubrum</i>	
Maple, Silver	<i>Acer saccharinum</i>	
Maple Sugar †	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb Variegata Wheeleris Dwarf
Mock Orange †	<i>Philadelphus spp</i>	Golden Snowflake Double White
Mondo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	Sunnyside
Myrtle, Crape	<i>Lagerstroemia indica</i>	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	<i>Myrica cerifera</i>	
Nandina	<i>Nandina domestica</i>	Compacta Nana
Narcissus	<i>Narcissus spp.</i>	
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oleander	<i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes
Osteospermum	<i>Osteospermum fruticosum</i>	Wirligig
Pachysandra	<i>Pachysandra terminalis</i>	
Palm, Bangalow		
Pampas Grass	<i>Cortaderia selloana</i>	
Pansy	<i>Viola x wittrockiana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Peach †	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picoti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	

Pine, Japanese Black	<i>Pinus thunbergiana</i>	
Pine, Loblolly	<i>Pinus taeda</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugo	<i>Pinus mugo</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus elliotii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, White	<i>Pinus strobus</i>	
Pineapple, Guava †	<i>Feijoa sellowiana</i>	
Pieris	<i>Pieris taiwanensis</i>	
Pittosporum, Japan		
Potentilla	<i>Potentilla nepalensis</i> <i>Potentilla fruticosa</i>	Abbotswood
Privet	<i>Ligustrum japonicum</i>	Golden Vicary Regal Texanum Wax Yellow Tipped
Privet, Glossy	<i>Ligustrum lucidum</i>	
Pyracantha	<i>Pyracantha koidzumii</i>	Gnome Lalandei Victory
Queen Palm	<i>Arecastrum rammanzoffianum</i>	
Quince, Japanese †		
Rhododendron	<i>Rhododendron spp</i>	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rockcress	<i>Arabis spp.</i>	Snowcap
Rhodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose †	<i>Rosa banksiaæ</i>	Luta
Rosemary †	<i>Rosmarinus officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow
Snapdragon	<i>Antirrhinum spp.</i>	
Sourwood	<i>Oxydendrum arboreum</i>	
Spiraea	<i>Astilbe X arendsii</i>	Fanall
Spiraea	<i>Spiraea spp.</i>	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houtte White
Spiraea, Garland	<i>Spiraea spp.</i>	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	Conica
Sweetflag, Grassyy-Le		
Sweetgum	<i>Liquidambar styraciflua</i>	

Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip, spp</i>	Apeldoorn
Verbena, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena spp.</i>	St. Paul
Viburnum	<i>Viburnum spp.</i>	American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden Mohican Wright
Vinca ( Periwinkle)	<i>Vinca minor</i>	
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yarrow	<i>Achillea spp.</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Densiformis

† Ornamental species only: Do not use on food producing trees and ornamentals.

### Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

### Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

### Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.



The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

<sup>TM</sup>Trademark of Dow AgroSciences LLC  
EPA-accepted date: 03-24-03

308/2E  
September 3, 2004



Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
U. S. Environmental Protection Agency  
Room 266A, Crystal Mall 2  
1801 South Bell Street  
Arlington, VA 22202

CRAB-BUSTER 0.08% FG (AI: DITHIOPYR)  
EPA REGISTRATION NUMBER: 62719-487  
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICE 98-10

Enclosed please find labeling for the notification action of Crab-Buster™ 0.08% FG lawn and ornamental herbicide. The following changes have been made by notification based on EPA-accepted copy (Notice of Registration) dated March 24, 2003 and notification coded "G2B / Crab-Buster 0.08% FG / Notification With Edits / 06-17-04":

1. Corrected trademark symbols throughout the label.
2. At the request of the state of New York, revised the following statement "In the state of New York, do not exceed 14.3 lb/1000 sq ft per year (equivalent to 0.5 pounds of active ingredient per acre). In Nassau and Suffolk Counties, do not exceed 7.2 lb/1000 sq ft per year (equivalent to 0.25 pounds of active per acre)." to read "In the state of New York, do not exceed 7.2 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre)."

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Contents of Submission**

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled Crab-Buster 0.08% FG (G2A/Crab-Buster 0.08% FG/Notif/09-02-04)  
(15 Pages plus Registration Notes) (5 Copies)

If you require further information, please contact Richard Bjerregaard, Regulatory Specialist at 317-337-4674 or Chris Hutchison, Registration Assistant for this product, at 317-337-4976.

Sincerely,

A handwritten signature in cursive script that reads "Kimberly S. Gilbert".

Kimberly S. Gilbert  
Regulatory Leader  
Regulatory Success – Americas  
317-337-4685  
317-337-4649 (FAX)  
ch/  
Enclosures