



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

16 MAY 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Kimberly Gilbert
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

Dear Ms. Gilbert:

SUBJECT: Label Amendment (Commercial and Residential)
Dimension 2EW
EPA Registration No. 62719-542
Your Application Dated February 20, 2008

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) copy of your final printed labeling before you release the product for shipment. This amended labeling supersedes all previously accepted ones.

Sincerely yours,

A handwritten signature in cursive script that reads "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

2 8 61

[Sub Label A: Commercial Use]

(Base label for rigid containers 5 gal or less):

Dimension[®] 2EW

Specialty Herbicide

ACCEPTED

16 MAY 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-542

Provides control of listed annual grasses and broadleaf weeds in:

- Established lawns
- Commercial sod farms
- Non-cropland and industrial sites
- Ornamental turf (including but not limited to golf course fairways, roughs, tee boxes)
- Container grown ornamentals
- Field-grown ornamentals
- Landscape ornamentals.

In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Active Ingredient

dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates

Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear protective clothing and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

3 8 61

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

WPS Uses: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks

WPS Uses: Mixers and loaders must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks
- Chemical-resistant apron

Non-WPS Uses: Mixers and loaders who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not

4761

apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Net Contents__

5 7 61

(Base label for refillable rigid containers larger than 5 gal):

Dimension[®] 2EW

Specialty Herbicide

Provides control of listed annual grasses and broadleaf weeds in:

- Established lawns
- Commercial sod farms
- Non-cropland and industrial sites
- Ornamental turf (including but not limited to golf course fairways, roughs, tee boxes)
- Container grown ornamentals
- Field-grown ornamentals
- Landscape ornamentals.

In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Active Ingredient

dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate.....	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates

Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear protective clothing and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

6 7 61

WPS Uses: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks

WPS Uses: Mixers and loaders must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks
- Chemical-resistant apron

Non-WPS Uses: Mixers and loaders who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents__

(Base label for nonrefillable rigid containers larger than 5 gal):

Dimension[®] 2EW

Specialty Herbicide

Provides control of listed annual grasses and broadleaf weeds in:

- Established lawns
- Commercial sod farms
- Non-cropland and industrial sites
- Ornamental turf (including but not limited to golf course fairways, roughs, tee boxes)
- Container grown ornamentals
- Field-grown ornamentals
- Landscape ornamentals.

In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Active Ingredient	
dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates
Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.
Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear protective clothing and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

WPS Uses: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks

WPS Uses: Mixers and loaders must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks
- Chemical-resistant apron

Non-WPS Uses: Mixers and loaders who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Chemical-resistant gloves >14 mils such as barrier laminate or butyl rubber

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

10 8 61

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
 9330 Zionsville Road
 Indianapolis, IN 46268

Net Contents__

11 7 61

(cover):

Dimension[®] 2EW

Specialty Herbicide

Provides control of listed annual grasses and broadleaf weeds in:

- Established lawns
- Commercial sod farms
- Non-cropland and industrial sites
- Ornamental turf (including but not limited to golf course fairways, roughs, tee boxes)
- Container grown ornamentals
- Field-grown ornamentals
- Landscape ornamentals.

In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Active Ingredient

dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates

Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

12 8 61

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Net Contents__

13 8 61

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING**

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear protective clothing and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

WPS Uses: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks

WPS Uses: Mixers and loaders must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks
- Chemical-resistant apron

Non-WPS Uses: Mixers and loaders who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear:

- Chemical-resistant gloves ≥ 14 mils such as barrier laminate or butyl rubber

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

14 8 61

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

REFORMULATION OR REPACKAGING OF THIS PRODUCT IS PROHIBITED.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves \geq 14 mils such as barrier laminate or butyl rubber
- Chemical-resistant footwear plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

- Keep unprotected persons out of treated area until sprays have dried.

15 8 61

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers larger than 5 gallons:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi

General Information

Dimension® 2EW specialty herbicide provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns, commercial sod farms, non-cropland and industrial sites, ornamental turf (including but not limited to golf course fairways, roughs, tee boxes), container-grown ornamentals, field-grown ornamentals, and landscape ornamentals.

16 7 61

This product will not control established weeds, except for emerged crabgrass, up to initiation of tillering. This stage of growth for crabgrass generally corresponds to the time when crabgrass seedlings are first visible in established turfgrasses and individual plants have five leaves or less. Applications to crabgrass after initiation of tillering will not provide satisfactory control. All other applications of this product should be made preemergence (prior to germination of target weeds).

This product is not effective until activated by 1/2 inch or more of rainfall or irrigation. Applications should be timed to ensure that activation has occurred prior to tillering stage of crabgrass development or prior to germination of all other weeds.

Note: In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Chemigation: Do not apply this product through any type of irrigation system.

Mixing Directions

Dimension 2EW Alone with Water as the Carrier

Fill a previously cleaned spray tank with water to about three-fourths of the desired volume. Add the recommended amount of Dimension 2EW to the tank. Complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the water source.

Dimension 2EW Alone with Fluid Fertilizer as the Carrier

Determine the compatibility of Dimension 2EW with the desired fluid fertilizer by mixing small proportional quantities in advance. See the Physical Compatibility Test section of this label. Then follow the mixing procedure listed below for tank mixtures.

Tank Mixtures

Dimension 2EW may be applied in tank mix combination with labeled rates of fluid fertilizers or other herbicides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product. Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels. Tank mix with Gallery[®] 75 DF or other herbicides to control a broader spectrum of broadleaf weeds. When tank mixing Dimension 2EW with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. See the Compatibility Test Mixing Instructions section of this label. The addition of a nonionic surfactant may improve control. Always read and follow the surfactant manufacturer's label recommendations.

Mixing Order for Tank Mixes: Place a 20 to 35 mesh screen or wetting basket over the filling port. Fill the spray tank 1/2 full with the appropriate carrier. Start agitation. Slowly add different formulation types in the order indicated below, allowing time for complete dispersion and mixing after addition of each product.

1. Compatibility agent (if needed)
2. Wettable powder or water dispersible granules (if used)
3. Suspension concentrates
4. Dimension 2EW and liquid (emulsifiable concentrate or liquid concentrate) pesticide (if used)
5. Water solutions

During the final filling of the tank, add water soluble liquid pesticide formulations (if used), followed by surfactants, marker dyes or foams, or drift control additives. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed.

Premixing: Dry and flowable formulations should be premixed with water (slurried) and added to the spray tank through a 20 to 35 mesh screen. This procedure assures good initial dispersion of these formulation types. If a liquid (emulsifiable concentrate, liquid concentrate, flowable) pesticide formulation is used, premix it with one part water before adding to the spray tank.

Physical Compatibility Test

Before mixing this product with fluid fertilizers and/or other pesticides, test compatibility by mixing all the components in a small jar in proportionate quantities.

Compatibility Test Mixing Instructions

Pesticide Formulation	If	Amount of Pesticide added to Spray Carrier (assuming volume is 25 gpa) ADD:
	Rate per Acre is:	Level Teaspoons per Pint Jar of Carrier Solution
Dry	1 lb	1 1/2
Liquid	1 qt	1

This compatibility test is designed for 25 gallons of spray solution per acre. The table above gives general guidelines for use rate ratios of pesticides to be tank mixed with this product. Determine the amount of pesticide to tank mix by referring to the pesticide label(s). Then, calculate the amount of pesticide to add to the jar based on use rate ratios in table. For a use rate of 1 lb per acre of dry pesticide, add 1 1/2 teaspoons to the jar. For a use rate of 1 quart per acre of liquid pesticide, add 1 teaspoon to the jar. Dimension 2EW should be added based on use rate ratios for liquid pesticides (for a use rate of 1 quart per acre, add 1 teaspoon to the jar). For changes in spray volume or herbicide rate, make appropriate changes in the ingredients for the test. Shake well after mixing.

If pesticide(s) does not form crystals, flakes, sludge, gels, oily films or layers, then the tested components are compatible. Incompatibility in any of the above-described forms will usually occur within 5 minutes after mixing. If components are incompatible, a compatibility agent should be used. Repeat the above compatibility test with a suitable compatibility agent (1/2 teaspoon per pint jar is equivalent to 2 pints per 100 gallons of spray solution). Do not use mixtures that show incompatible signs such as formation of crystals, flakes, sludge, gels, oil films or layers.

Grass and Broadleaf Weeds Controlled by Dimension 2EW

Used as directed, Dimension 2EW controls annual grass and broadleaf weeds listed in the table below if applied preemergence. This product will not control emerged broadleaf weeds or grasses (except for crabgrass prior to tillering stage of growth). Therefore, the area to be treated must be free of weeds prior to application.

Common Name	Scientific Name
Grasses	
barley	<i>Hordeum</i> spp.
barnyardgrass	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
brome	<i>Bromus</i> spp.
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>

crabgrass, southern	<i>Digitaria ciliaris</i>
crowfootgrass	<i>Dactyloctenium aegyptium</i>
dallisgrass (seedling)	<i>Paspalum dilatatum</i>
foxtail, giant	<i>Setaria faberi</i>
foxtail, green	<i>Setaria verdi</i>
foxtail, yellow	<i>Pennisetum glaucum</i>
goosegrass	<i>Eleusine indica</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
Mary's grass	<i>Microstegium vimineum</i> (Trin.) A. Camus var. <i>imberbe</i>
oats, wild	<i>Avena fatua</i>
ryegrass (annual & perennial)	<i>Lolium spp.</i>
sandbur	<i>Cenchrus spp.</i>
smutgrass	<i>Sporobolus indicus</i>

Broadleaf Weeds

bittercress	<i>Cardamine spp.</i>
carpetweed	<i>Mollugo verticillata</i>
chickweed	<i>Stellaria spp.</i>
geranium, Carolina	<i>Geranium carolinianum</i>
henbit	<i>Lamium spp.</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lespedeza, common	<i>Lespedeza striata</i>
maretail	<i>Conyza canadensis</i>
medic, black	<i>Medicago lupulina</i>
mulberry weed	<i>Fatoua villosa</i>
mustard	<i>Brassica spp.</i>
oxalis, buttercup	<i>Oxalis pes-caprae</i>
parsley-piert	<i>Alchemilla arvensis</i>
pigweed, redroot	<i>Amaranthus retroflexus</i>
pineappleweed	<i>Matricaria matricarioides</i>
purslane, common	<i>Portulaca oleracea</i>
rocket, London	<i>Sisymbrium irio</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>
speedwell, corn	<i>Veronica arvensis</i>
spurge, garden	<i>Euphorbia hirta</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Uses

Turfgrass

Use Dimension 2EW on seeded, sodded, or sprigged lawns and ornamental turfgrass that are well established. Newly established turf must have developed a good root system and a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging before making the first application of this product. Use of this product on turf that is not well-established, or has been weakened by weather, pest, disease, chemical, mechanical or other related stress, may result in turf injury.

Specific Use Precautions and Restrictions:

- Do not apply this product to golf course putting greens.

- Dimension 2EW will prevent the germination of annual bluegrass. Dimension 2EW will not affect established annual bluegrass. If maintenance of annual bluegrass is desired, using this product during the time of annual bluegrass germination is not recommended. In the states of AZ, CA, NV, OR, WA, NM, ID, MT and UT, Dimension 2EW may contribute to the thinning or stand reduction in established stands of annual bluegrass.
- It is recommended that sod be established for at least 6 months before applying Dimension 2EW.
- Do not apply this product within 3 months of harvesting sod.
- Early postemergence applications of this product will control crabgrass only if applied prior to the fifth leaf (first tiller) stage of growth.
- For best results, cultural practices that disturb the soil, such as verticutting and core-, spike-, or hydro-aerification, should be done before applying this product.
- Do not apply this product until the turfgrass has recovered from cultural practices such as verticutting or core-, spike-, or hydro-aerification.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not apply more than 0.73 oz of Dimension 2EW per 1000 sq ft (2 pints per acre) per application.
- Do not apply more than 2.2 oz of Dimension 2EW per 1000 sq ft (6 pints per acre) per year using split or sequential applications.
- In New York State, do not apply more than 2 pints of Dimension 2EW (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Dimension 2EW should only be applied to turf that is composed of the following turfgrass species that have been determined to be tolerant to application of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Established Cool Season Turfgrasses

Common Name	Scientific Name
bentgrass, creeping [†]	<i>Agrostis palustris</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, fine ^{††}	<i>Festuca rubra</i>
fescue, tall	<i>Festuca arundinacea</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turfgrasses

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum</i>
Bermudagrass ^{†††}	<i>Cynodon dactylon</i>
buffalograss ^{††††}	<i>Buchloe dactyloides</i>
carpetgrass	<i>Axonopus affinis</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
seashore paspalum	<i>Paspalum vaginatum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>

[†] Do not use this product on certain varieties of creeping bentgrass, such as cohansey, carmen, seaside, and Washington as undesirable turfgrass injury may result. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (*Agrostis tenuis*) varieties.

^{††} Do not use this product on certain varieties of fine fescue as undesirable turf injury may result. The following fine fescue varieties have been found to be sensitive to this product: Atlanta, banner, beauty, bilgart, CF-2, enjoy, HF-93, highlight, ivalo, Jamestown, koket, majenta, Mary, pennlawn, Tamara, Tatjana, waldorf, and waldina. Not all varieties of fine fescue have been tested.

^{†††} Do not use this product on Tifgreen (328) hybrid Bermudagrass as undesirable turfgrass injury may result. Other common and hybrid Bermudagrass varieties are tolerant.

20 8 61

†††† Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Reseeding, Overseeding, or Sprigging

Reseeding, overseeding, or sprigging of treated areas within 3 months after a single application of this product, or within 4 months after a split application program totaling more than 0.73 oz per 1000 sq ft (2 pints per acre), may inhibit the establishment of desirable turfgrasses. However, overseeding of Bermudagrass with perennial ryegrass 8 weeks after an application or as early as 6 weeks after application if slight injury to perennial ryegrass can be tolerated is a recommended exception.

When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

Application Directions

Apply Dimension 2EW through conventional liquid application equipment in a sufficient volume of carrier solution to provide a uniform spray distribution. Applying this product through cluster nozzles or other boomless spray equipment may not give satisfactory results due to variability in application volume and spray pattern. A hand held spray gun may be used. Calibrate application equipment prior to usage. Avoid streaking, skips, or excess overlaps during application. The use of marker dyes or foams aids in making more accurate applications.

Crabgrass Control

Preemergence and Early Postemergence Control

This product provides preemergence control of crabgrass (including the large, smooth, and southern species) when applied prior to crabgrass emergence in established lawns and ornamental turf. This product also provides early postemergence control of crabgrass during early crabgrass seedling growth following emergence. However, it is often difficult to observe very small crabgrass seedlings at early stages of growth in well-established turfgrass. Early postemergence crabgrass control is obtained only when this product is applied prior to the tillering of crabgrass (less than 5 leaves per plant), which generally corresponds to the time when crabgrass seedlings are easily observed in lawn or turf. The practical benefit of this product's additional, early postemergence activity is a "window of application" during which control of crabgrass can be achieved; that is 2 to 8 weeks longer than products that provide only preemergence control. Depending upon climatic conditions and rate of growth, the time from crabgrass emergence until tillering will vary from 2 to 8 weeks. The addition of a nonionic surfactant at 0.5% by volume (2 qt per 100 gallons of spray) may improve early postemergence control. Read and follow the surfactant manufacturer's label recommendations.

Tank Mixtures for Early Postemergence Control:

Application of this product alone provides early postemergence control of crabgrass when treated prior to reaching the tillering stage of growth. If applied in combination with either MSMA herbicide or Acclaim herbicide, control of crabgrass with up to 3 tillers may be achieved.

If preemergence herbicides have been applied prior to a postemergence application, use the applicable Program 1 or Program 2 use rate of Dimension 2EW for your area; otherwise, apply the Program 3 use rate in combination with MSMA or Acclaim.

A compatibility test (see Compatibility Test in Mixing Directions section) is suggested before tank mixing this product with fluid fertilizers and/or either MSMA or Acclaim. Refer to the labels for MSMA or Acclaim for information on tolerance of specific turfgrass species. Observe all limitations, precautionary statements, and use restrictions on the respective labels when using them in tank mixtures.

Application Frequency and Timing

21 7 61

This product may be applied as a single application, a split application, or a sequential application for crabgrass control in the spring, summer, or fall.

Early Summer or Spring Application: For applications made in the spring or early summer, apply Dimension 2EW at the appropriate rate corresponding to one of the three control programs listed in the table below. The rate used depends upon the user's location, the mowing height of the turfgrass, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending upon weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Late Summer or Fall Application: This product can also be applied in the late summer or early fall (late August through November) at the Program 3 use rates listed in the table below. A fall application followed by an appropriately timed spring application provides season-long weed control. Do not exceed the maximum use rate per year. For *Poa annua* control see Section titled "Poa annua (Annual Bluegrass) Control" below Recommended Use Rate table.

Program 1: Use Program 1 for preemergence control in turfgrass that is cut relatively high (e.g., residential lawns). This program provides 3 to 5 months of preemergence crabgrass control and also provides early postemergence control of crabgrass up to the 3-leaf stage at sites where turfgrass is cut relatively high. Split applications may also be made, with the rates in the Recommended Use Rates table being split across two applications made 5 to 10 weeks apart. Split applications may provide improved weed control.

Program 2: Use Program 2 for preemergence control in turf: (a) where turfgrass is cut relatively low (e.g., golf fairways); and (b) where turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control up to crabgrass tillering at sites where turfgrass is cut relatively high. Split applications may also be made, with the rates in the Recommended Use Rates table being split across two applications made 5 to 10 weeks apart. Split applications may provide improved weed control.

Program 3: Use Program 3 for preemergence control in turf: (a) where turfgrass is cut relatively low (e.g., golf fairways); and (b) where turf maintenance or weed control was not conducted during the previous year. This program provides 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control up to crabgrass tillering at sites where turfgrass is cut relatively low (e.g., golf fairways). Make subsequent, sequential pre and/or postemergence applications where longer periods of control are desired. Split applications may also be made, with the rates in the Recommended Use Rates table being split across two applications made 5 to 10 weeks apart. Split applications may provide improved weed control.

Recommended Use Rate Table

Note: 1 oz equals 6 teaspoons

Region	Application Rates	Program 1	Program 2	Program 3
All states, except NY [†] and parts of states not listed in transition, south, coastal south or west	pt/acre	.75	1	1.5
	oz/1000 sq ft	0.28	0.37	0.55
	lb ai/acre	0.188	0.25	0.38
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH,	pt/acre	1	1.5	2 [†]
	oz /1000 sq ft	0.37	0.55	0.73 [†]
	lb ai/acre	0.25	0.38	0.5 [†]

coastal areas of CT, & RI				
South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, TX	pt/acre	1.5	1 + 1	1.2 + 1.2
	oz /1000 sq ft	0.55	0.37 + 0.37	0.46 + 0.46
	lb ai/acre	0.38	0.25 + 0.25 [†]	0.31 + 0.31 [†]
Coastal South: HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, TX	pt/acre	1 + 1	1.25 + 1.25	1.5 + 1.5
	oz /1000 sq ft	0.37 + 0.37	0.46 + 0.46	0.55 + 0.55
	lb ai/acre	0.25 + 0.25 [†]	0.31 + 0.31 [†]	0.38 + 0.38
West: AZ, CA, NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	pt/acre	0.75 + 1	1 + 1.5	1 + 1
	oz /1000 sq ft	0.28 + 0.37	0.37 + 0.55	0.37 + 0.37
	lb ai/acre	0.18 + 0.25	0.25 + 0.38	0.25 + 0.25

[†] In New York State, this product may be used by commercial applicators only at no more than 2 pints (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Maximum Use Rates (All Turf Uses)

Do not apply more than 0.73 oz of Dimension 2EW per 1000 sq ft (2 pints per acre) per application. Do not apply more than 2.2 oz of Dimension 2EW per 1000 sq ft (6 pints per acre) per year using split or sequential applications.

Poa Annua (annual bluegrass) Control

Apply Dimension 2EW for preemergence control of *Poa annua* (annual bluegrass) at a rate of 0.38 to 0.5 lbs active ingredient per acre. This is equivalent to 1.5 to 2 pints per acre.

- Apply 6 to 8 weeks before overseeding bermudagrass with perennial ryegrass. Specific to perennial ryegrass; not recommended for *Poa trivialis* or bentgrass.
- Minimum seeding rate of perennial ryegrass is 400 lbs per acre.
- Use limited to fairways and roughs
- Perennial varieties of *Poa annua* (var. repens) may not be controlled as well as the true annual.
- Do not apply earlier than 16 weeks after over-seeding unless injury to the ryegrass can be tolerated.
- A follow-up treatment 16 weeks after overseeding offers an early season crabgrass treatment and helps suppress some winter annual broadleaf weeds.

Use Directions for Noncropland and Industrial Sites

Apply Dimension 2EW for preemergence control of annual grasses and broadleaf weeds listed in Grass and Broadleaf Weeds Controlled by Dimension 2EW in terrestrial non-crop areas including farm yards, fence rows, highway, utility and railroad rights-of-way, airports, recreation areas, campgrounds, and industrial sites (lumber yards, tank farms, and storage areas).

Applied preemergence, Dimension 2EW controls weeds as they germinate. This product will not control established weeds. Apply prior to germination of target weeds or to bare ground. The best weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to applying, control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide.

To be effective, Dimension 2EW must be activated by 1/2 inch or more of rainfall or irrigation prior to germination of target weeds. Once the treatment is activated, avoid excessive soil disruption that may

break down the herbicide barrier. Minimal surface disruption such as raking should not break down the herbicide barrier.

Specific Use Precautions and Restrictions:

- Do not apply when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not graze livestock or feed forage cut from areas treated with this product.
- For ornamentals within non-crop areas, apply only after transplanting when soil around roots has been thoroughly settled by rainfall or irrigation and no cracks are present or injury will result.

Equivalent Application Rates:

Equivalent Rates of Dimension 2EW			
(pt/acre)	(oz/1000 sq ft)	(oz/100 sq ft)	(mL/100 sq ft)
2	0.73	0.073	0.36

Make sequential applications at 3 to 4 month intervals for extended preemergence weed control. Do not exceed maximum use rates per year

Maximum Use Rates

- **Split or sequential applications:** Do not use more than 0.73 oz of Dimension 2EW per 1000 sq ft (2 pints per acre) per application or more than 2.2 oz of Dimension 2EW per 1000 sq ft (6 pints per acre) per year.

In New York State, do not apply more than 2 pints of Dimension 2EW (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Use Directions For Ornamentals (Landscape, Field Grown, and Container Grown)

Dimension[®] 2EW specialty herbicide provides preemergence control of listed annual grasses and broadleaf weeds in areas planted with tolerant ornamental plants listed on this label. It is intended for use on plants grown for aesthetic purposes in landscaped areas or in container or field grown production nurseries. When applied as directed, the ornamental plants listed on this label have shown tolerance to over-the-top applications of Dimension 2EW.

Specific Use Precautions and Restrictions:

- Apply Dimension 2EW to established ornamentals only.
- Do not apply this product to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into the soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, heat stress, nematodes or diseases).
- Do not apply when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants unless they are listed on this label.
- Do not apply this product to grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.
- Do not apply this product directly to plants that are grown for food (e.g., fruit trees or maple trees tapped for syrup).
- Do not apply this product in enclosed structures and greenhouses.
- Do not apply more than 0.73 oz of Dimension 2EW per 1000 sq ft (2 pints per acre) per application and no more than 2.2 oz of Dimension 2EW per 1000 sq ft (6 pints per acre) per year using split or sequential applications.

24 8 61

- In New York State, do not apply more than 2 pints of Dimension 2EW (0.5 lb active ingredient) per acre per year. In Nassau and Suffolk counties of New York, do not exceed 1 pint per year of this product (equivalent to 0.25 lb of active ingredient per acre).

Treatment of Ornamental Species Not Listed on the Label for Dimension 2EW: It is impossible to evaluate tolerance to this product on all ornamental plant species or varieties or under all possible growing conditions. Users who wish to use Dimension 2EW on ornamental species not recommended on this label may determine the suitability for use by treating a small number of ornamental plants at a recommended rate. Prior to treatment of larger areas, treated plants should be observed for any symptoms of herbicidal injury, such as foliar damage, reduced vigor or stand reduction, during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

Application Directions

Apply Dimension 2EW as a directed spray in established ornamentals or as a broadcast over-the-top spray to certain established ornamentals (see ornamental plant listing). Make directed sprays to the soil at the base of the ornamentals.

To reduce injury potential:

- Apply product with calibrated equipment using a minimum of 2 gallons of water per 1000 sq. ft.
- Shortly after application apply overhead irrigation to activate the herbicide and wash Dimension 2EW from plant surface onto soil surface.
- In the spring when buds are rapidly growing and expanding, over the top application of Dimension 2EW may temporarily injure new growth of desirable plants. To reduce the possibility of injury at this time, wait to apply Dimension 2EW over the top of newly emerged vegetation until it has hardened off, unless your experience indicates that the ornamental plant will not be injured by the over the top application.
- Do not apply to plants that are under stress such as heat, drought or frost damage.

Dimension 2EW is a preemergence herbicide that controls weeds during germination. Dimension 2EW does not control emerged broadleaf or grass weeds except crabgrass up to tillering stage of growth. Apply prior to germination of target weeds to bare ground. Optimum weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to applying, control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide. After applying Dimension 2EW, excessive soil disruption may breakdown the herbicide barrier. Minimal surface disruption such as raking should not break down the herbicide barrier. Following transplanting, care must be taken that soil or planting mixes have settled firmly through irrigation, rainfall or packing and that there are no cracks that would allow direct contact of this product to the plant roots or plant injury may occur.

Application Rates

Apply Dimension 2EW prior to germination of target weed species. Make sequential applications at 3 to 4 month intervals for extended preemergence weed control. Do not exceed maximum use rates per year.

When treating a small area, apply Dimension 2EW with a calibrated sprayer that assures accurate, uniform spray distribution. In general, Dimension 2EW should be thoroughly mixed with clean water at 0.5 to 0.73 oz of product per 1000 sq ft per application and applied at 20 to 40 psi in a minimum of 1 gallon of water per 1000 sq ft.

Equivalent Rates of Dimension 2EW			
(pt/acre)	(oz/1000 sq ft)	(oz tsp/100 sq ft)	(oz mL/100 sq ft)
2	0.73	0.073	0.36

Tolerant Ornamentals

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials.

		Acceptable Application Method Noted by a (X)		Application to Field/Landscape or Container Grown Ornamentals Noted by a (X)	
Name	Tolerant Cultivars	Over the Top	Directed	Container Grown	Field/Landscape Grown
abelia, dwarf (<i>Abelia x grandiflora</i>)	nana		x		x
abyssinian red banana (<i>Ensete ventricosum</i>)	maureli	x	x	x	x
Ajuga carpet bugle (<i>Ajuga reptans</i>)	bronze		x		x
(<i>Ajuga genevensis</i>)	bronze beauty		x		x
almond, flowering (<i>Prunus gladiosa</i>)			x		x
apple [†] (<i>Malus pumila</i>)			x		x
arborvitae (<i>Thuja occidentalis</i>)	nigra		x		x
	pyramidalis		x		x
	smaragh		x		x
	techny		x		x
	woodwardii		x		x
arborvitae, dwarf golden (<i>Thuja orientalis</i>)	aurea nana	x	x	x	x
ash, green (<i>Fraxinus pennsylvanica</i>)			x		x
ash, autumn purple (<i>Fraxinus americana</i>)	autumn purple		x		x
aster, Chinese (<i>Callistephus chinensis</i>)	dwarf queen		x		x
azalea (<i>Rhododendron spp.</i>)	brilliant		x		x
	buccaneer		x		x
	carror		x		x
	chimes (Belgian)		x		x
	Elsie Lee		x		x
	exbury		x		x
	fashion		x		x
	hardijzer beauty		x		x
	hershey red		x		x
	higasa		x		x
	hinocrimson		x		x
	Holland (hybrid)		x		x
	Marion Lee		x		x
	northern lights		x		x
	orange cup		x		x
	orchid lights		x		x
	snow		x		x
	southern charm		x		x
azalea, flame (<i>Rhododendron calendulaceum</i>)			x		x
azalea, kirishima (<i>Rhododendron obtusum</i>)			x		x

bamboo, heavenly (<i>Nandina domestica</i>)	compacta nana		x x		x x
barberry, Japanese (<i>Berberis thunbergii</i>)	aurea dwarf pygmy green kobold pygmy red rose glow		x x x x x x		x x x x x x
barberry, purple (<i>Berberis thunbergii</i> var <i>atropurpurea</i>)	atropurpurea		x		x
basket flower (<i>Gaillardia grandiflora</i>)			x		x
beach grass (<i>Ammophila breviligulata</i>)			x		x
bearberry (common) (<i>Arctostaphylos uva-ursi</i>)	Massachusetts		x		x
bee balm (<i>Monarda didyma</i>)			x		x
begonia (<i>Begonia</i> spp.)			x		x
birch, river (<i>Betula nigra</i>)			x		x
blackeyed Susan (<i>Rudbeckia hirta</i>)	goldstrum		x		x
blanket flower (<i>Gaillardia</i> spp.)			x		x
blue fescue (<i>Festuca ovina</i>)			x		x
blueberry [†] (<i>Vaccinium</i> spp.)	bluecrop blue jay Jersey north blue northland		x x x x x		x x x x x
bottlebrush (<i>Callistemon citrinus</i>)	Little John	x	x	x	X
bougainvillea (<i>Bougainvillea spectabilis</i>)	James Walker Scarlet O' Hara	x	x x	x	x x
boxwood (<i>Buxus microphylla</i>)	japonica		x		X
boxwood, green beauty (<i>Buxus microphylla</i> <i>japonica</i>)	green beauty	x	x	x	x
boxwood, welleri (<i>Buxus sempervirens</i>)	winter gem	x	x	x	x
broom (<i>Cytisus scoparius</i>) (<i>Genista pilosa</i>)	moonlight Vancouver gold		x x		x x
camellia (<i>Camellia japonica</i>) (<i>Camellia sasanqua</i>)	debutante mathotiana supreme chansonette setsukgekka	x x	x x x x	x x	x x x x

candytuft (<i>Iberis sempervirens</i>)	snow white		x		x
carex, variegated (<i>Carex spp.</i>)			x		x
carpet bugle (<i>Ajuga reptans</i>)	bronze		x		x
(<i>Ajuga genevensis</i>)	bronze beauty		x		x
cedar, red (<i>Juniperus virginiana</i>)			x		x
celosia (<i>Celosia spp.</i>)			x		x
centaura (<i>Centaurea montana</i>)			x		x
cockscomb, plumosa (<i>Celosia cristata</i>)	scarlet plumosa		x		x
coleus (<i>Coleus blumei</i>)	red kewpie		x		x
columbine (<i>Aquilegia spp.</i>)			x		x
coneflower, purple (<i>Echinacea purpurea</i>)	magnus purple	x	x	x	x
			x		x
copper leaf (<i>Acalypha wilkesiana</i>)			x		x
coreopsis (<i>Coreopsis spp.</i>)	moonbeam		x		x
corn flower (<i>Centaurea spp.</i>)			x		x
cotoneaster (<i>Cotoneaster apiculatus</i>)			x		x
coyotebrush (<i>Baccharis pilularis</i>)			x		x
cycad (<i>Cycas revoluta</i>)			x		x
cypress, bald (<i>Taxodium distichum</i>)			x		x
cypress, Italian (<i>Cupressus sempervirens</i>)	glauca		x		x
cypress, Japanese false (<i>Chamaecyparis obtusa</i>)	gracilis		x		x
cypress, leyland (<i>Cupressocyparis leylandii</i>) hybrid			x		x
daffodil (<i>Narcissus spp.</i>)	King Alfred		x		x
daylilly (<i>Heemerocallis spp.</i>)	Aztec gold		x		x
	bright yellow (hybrid)		x		x
	single gold (evergreen)		x		x
	Wilson's yellow		x		x
dianthus (sweet William) (<i>Dianthus spp.</i>)			x		x
delphinium (<i>Delphinium spp.</i>)	magic fountain		x		x
dogwood (<i>Cornus florida</i>)			x		x

dogwood, American (<i>Cornus sericea</i>)	flavarimea		x		x
Douglas fir (<i>Pseudotsuga menziesii</i>)			x		x
dusty miller (<i>Senecio cineraria</i>)		x	x	x	x
elm (<i>Ulmus parvifolia</i>)	drake		x		x
eulaliagrass (<i>Miscanthus sinensis</i>)	maiden grass	x	x	x	x
euonymus (<i>Euonymus japonicus</i>)	Argenteo-variegata		x		x
	Auereomarginata		x		x
	colorata		x		x
(<i>Euonymus fortunei</i>)	emerald gaiety		x		x
	emerald n' gold		x		x
	gold edge		x		x
	gold princess		x		x
	Manhattan	x	x	x	x
	silver king		x		x
	tricolor		x		x
	vegetus		x		x
euryps, green leaved (<i>Euryops pectinatus</i>)	viridis	x	x	x	x
fan palm, European (<i>Chamaerops humilis</i>)			x		x
fan palm, Mexican (<i>Washingtonia robusta</i>)			x		x
fern (various) (<i>Asparagus spp.</i>)			x		x
fescue (<i>Festuca glauca</i>)			x		x
fetterbush (<i>Leucothoe fontanesiana</i>)	rainbow		x		x
fig (<i>Ficus retusa</i>)	nitidia		x		x
fir fraser (<i>Abies fraseri</i>)			x		x
Forsythia (<i>Forsythia</i> x 'Arnold Dwarf')	arnold dwarf		x		x
(<i>Forsythia viridissima</i>)	bronxensis dwarf		x		x
(<i>Forsythia xintermedia</i>)	lynwood gold		x		x
(<i>Forsythia</i> x 'Meadowlark')	meadowlark		x		x
(<i>Forsythia x intermedia</i>)	spring glory	x	x	x	x
(<i>Forsythia suspensa</i>)	weeping		x		x
fountain grass, purple (<i>Pennisetum setaceum</i>)	rubrum	x	x	x	x
fuchsia (<i>Fuchsia spp.</i>)			x		x

galium (<i>Galium odoratum</i>)			x		x
gardenia (<i>Gardenia jasminoides</i>)	mystery radicans white gem		x x x		x x x
Garlic, variegated society (<i>Thulbaghia violacea</i>)	variegata	x	x	x	x
geranium (<i>Pelargonium x hortorum</i>)			x		x
gum (<i>Eucalyptus citriodora</i>)			x		x
hawthorn (<i>Crataegus spp.</i>)	cockspur white crimson cloud enchantress Jack Evans Washington white		x x x x x		x x x x x
hawthorn, Indian (<i>Rhaphiolepis indica</i>)	ballerina	x	x	x	x
heather, twisted (<i>Erica cinerea</i>)	Mediterranean pink		x		x
hemlock, Canada (<i>Tsuga canadensis</i>)			x		x
hibiscus (<i>Hibiscus spp.</i>)	blue bird brilliant hula girl		x x x		x x x
holly (<i>Ilex x 'Nellie R. Stevens'</i>) (<i>Ilex x attenuata</i>)	Nellie R. Stevens fosteri Savannah	x	x x x	x	x x x
holly, blue (<i>Ilex x meserveae</i>)	blue boy blue girl China girl		x x x		x x x
holly, cassine (<i>Ilex cassine</i>)		x	x	x	x
holly, Chinese (<i>Ilex cornuta</i>)	burfordii needlepoint		x x		x x
holly, Japanese (<i>Ilex crenata</i>)	compacta hellerie Japanese northern beauty steeds		x x x x		x x x x
holly, yaupon (<i>Ilex vomitoria</i>)			x		x
honeysuckle (<i>Lonicera xylosteum</i>) (<i>Lonicera japonica</i>) (<i>Lonicera tatarica</i>)	Claveyí dwarf halliana Canadian white Zabelli		x x x x		x x x x
hosta (<i>Hosta sieboldii</i>) (<i>Hosta lancifolia</i>)	albo marginata		x x		x x
ice plant (<i>Carpobrotus edulis</i>)		x	x	x	x
ice plant, white trailing (<i>Delosperma alba</i>)		x	x	x	x

impatiens (<i>Impatiens spp.</i>)			x		x
iris (<i>Iris spp.</i>)	dwarf blue		x		x
	wedgewood		x		x
ivy, English (<i>Hedera helix</i>)	Bulgaria		x		x
	thorndale		x		x
jasmine, Asian (<i>Trachelospermum asiaticum</i>)			x		x
jasmine, star (<i>Trachelospermum jasminoides</i>)			x		x
juniper (<i>Juniperus spp.</i>)	Arcadia		x		x
	Armstrong		x		x
	bar harbor		x		x
	blue chip		x		x
	blue Pacific		x		x
	blue rug		x		x
	blue star		x		x
	broadmoor		x		x
	buffalo		x		x
	calgary carpet		x		x
	emerald sea		x		x
	emerald spreader		x		x
	endora compacta		x		x
	fruitlandi		x		x
	green		x		x
	gold tip		x		x
	hetzi		x		x
	hughes		x		x
	Manhattan blue		x		x
	parsoni		x		x
	pfitzeriana		x		x
	plumosa		x		x
	Prince of Wales		x		x
	procumbens dwarf		x		x
	San Jose		x		x
	sargent blue		x		x
sargent green		x		x	
scandia		x		x	
scopulorum moonglow		x		x	
spartan		x		x	
tamariscifolia		x		x	
weberi		x		x	
Youngstown		x		x	
Yukon belle		x		x	
juniper (<i>Juniperus chinensis</i>)	gold coast	x	x	x	x
	torulosa – Hollywood twisted	x	x	x	x
juniper (<i>Juniperus procumbnes nana</i>)	prostrate Japanese garden	x	x	x	x

juniper (<i>Juniperus sabina</i> 'tamariscifolia')	tamarix	x	x	x	x
juniper (<i>Juniperus scopulorum</i>)	scopulorum skyrocket	x	x	x	x
king palm (<i>Archontophoenix</i> <i>cunninghamiana</i>)		x	x	x	x
laurel, Carolina cherry (<i>Prunus caroliniana</i>)	bright n' tight	x	x	x	x
laurel, mountain (<i>Kalmia latifolia</i>)			x		x
leucothoe (<i>Leucothoe fontanesiana</i>)			x		x
ligustrum, Japanese (<i>Ligustrum japonicum</i>)			x		x
lily (<i>Agapanthus spp.</i>)	streamline		x		x
Lily of the Nile (<i>Agapanthus africanus</i>)	albus Peter Pan		x x		x x
lilyturf (<i>Liriope muscari</i>)	evergreen giant lilac beauty majestic monroe white silvery sunproof variegata	x x	x x x x x	x x	x x x x x
lilyturf, creeping (<i>Liriope spicata</i>)			x		x
maple, amur (<i>Acer ginnala</i>)	emerald elf	x	x	x	x
maple, Japanese (<i>Acer palmatum</i>)			x		x
maple, Norway (<i>Acer platanoides</i>)			x		x
maple, red [†] (<i>Acer rubrum</i>)	red sunset	x	x	x	x
maple, silver (<i>Acer saccharinum</i>)			x		x
maple sugar [†] (<i>Acer saccharum</i>)			x		x
marigold (<i>Tagetes patula</i>)	honeycomb variegata wheeleris dwarf		x x x		x x x
mock orange [†] (<i>Philadelphus spp</i>)	golden snowflake double white		x x		x x
mondo grass (<i>Ophiopogon japonicus</i>)			x		x
moss rose (<i>Portulaca grandiflora</i>)	sunnyside		x		x
mountainash (<i>Sorbus aucuparia</i>)			x		x

myrtle, crape (<i>Lagerstroemia indica</i>)	faurei langer muskogee standard pink		x x x x		x x x x
myrtle, wax (<i>Myrica californica</i>)			x		x
narcissus (<i>Narcissus spp.</i>)			x		x
oak, laurel (<i>Quercus laurifolia</i>)			x		x
oak, pin (<i>Quercus palustris</i>)			x		x
oak, red (<i>Quercus rubra</i>)			x		x
oak, southern (<i>Quercus virginiana</i>)			x		x
oak, willow (<i>Quercus phellos</i>)			x		x
oleander (<i>Nerium oleander</i>)	hardy red petite pink sister agnes		x x x		x x x
osteospermum (<i>Osteospermum fruticosum</i>)	whirligig		x		x
pachysandra (<i>Pachysandra terminalis</i>)			x		x
palm, bangalow (<i>Archontophoenix cunninghamiana</i>)			x		x
palm, queen (<i>Syagrus romanzoffianum</i>)		x	x	x	x
pampas grass (<i>Cortaderia selloana</i>)			x		x
pansy (<i>Viola x wittrockiana</i>)			x		x
paper flower (<i>Bougainvillea glabra</i>)	Barbara Karst	x	x	x	x
peach [†] (<i>Prunus persica</i>)			x		x
periwinkle, dwarf (<i>Vinca minor</i>)			x		x
petunia (<i>Petunia x hybrida</i>)	picoti		x		x
photinia, red tip (<i>Photinia x fraseri</i>)			x		x
pieris (<i>Pieris taiwanensis</i>)			x		x
pieris, Japanese (<i>Pieris japonica</i>)	mountain fire		x		x
pine, Afghan (<i>Pinus eldarica</i>)		x	x	x	x
pine, Austrian black (<i>Pinus nigra</i>)		x	x	x	x

pine, Japanese black (<i>Pinus thunbergia</i>)			x		x
pine, loblolly (<i>Pinus taeda</i>)			x		x
pine, longleaf (<i>Pinus palustris</i>)			x		x
pine, mugo or Swiss Mt. (<i>Pinus mugo</i>)			x		x
pine, Scotch (<i>Pinus sylvestris</i>)			x		x
pine, slash (<i>Pinus elliotii</i>)			x		x
pine, Virginia (<i>Pinus virginiana</i>)			x		x
pine, white (<i>Pinus strobus</i>)		x	x	x	x
pineapple, guava† (<i>Feijoa sellowiana</i>)			x		x
pittosporum, tobira (<i>Pittosporum tobira</i>)	Wheeler's dwarf	x	x	x	x
plumbago, cape (<i>Plumbago auriculata</i>)	royal cape	x	x	x	x
potentilla (<i>Potentilla fruticosa</i>)) (<i>Potentilla nepalensis</i>)	abbotswood		x x		x x
privet (<i>Ligustrum x vicaryii</i>) (<i>Ligustrum japonicum</i>)	golden vicary regal texanum yellow tipped		x x x x	x	x x x x
privet, glossy (<i>Ligustrum lucidum</i>)			x		x
Pyracantha or firethorn (<i>Pyracantha x 'Gnome'</i>) (<i>Pyracantha coccinea</i>) (<i>Pyracantha koidzumii</i>)	gnome lalandei victory	x	x x x	x	x x x
queen palm (<i>Arecastrum rammanzoffianum</i>)			x		x
quince, Japanese† (<i>Chaenomeles japonica</i>)			x		x
Rhododendron (<i>Rhododendron spp</i>)	album Cunningham white PJM purple gem silvery pink		x x x x x		x x x x x
rhododendron, Carolina (<i>Rhododendron carolinianum</i>)			x		x
rhododendron, catawba (<i>Rhododendron catawbiense</i>)			x		x

rhododendron, rhodie max - rosebay (<i>Rhododendron maximum</i>)			x		x
ribbon grass (<i>Phalaris arundinacea</i>)			x		x
rockcress (<i>Arabis caucasica</i>)	snowcap		x		x
rose† (<i>Rosa banksiae</i>)	luta		x		x
rose, knockout shrub (<i>Rosa</i> spp. hybrid)	knockout	x	x	x	x
rosemary† (<i>Rosmarinus officinalis</i>)			x		x
rosemary, bog (<i>Andromeda polifolia</i>)	nana		x		x
salvia (<i>Salvia farinacea</i>)	rhea		x		x
sedum (<i>Sedum spurium</i>)	dragon blood red red carpet yellow		x x x		x x x
smoketree, royal purple (<i>Cotinus coggygria</i>)	royal purple		x		x
snapdragon (<i>Antirrhinum</i> spp.)			x		x
snowball, common (<i>Viburnum opulus</i>)	sterile	x	x	x	x
sourwood (<i>Oxydendrum arboreum</i>)			x		x
spirea (<i>Astilbe X arendsii</i>)	fanall		x		x
spirea (<i>Spiraea</i> spp.)	Anthony Waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white		x x x x x x		x x x x x x
spirea, garland (<i>Spiraea x arguta</i> .)			x		x
spruce, Black Hills (<i>Picea glauca var densata</i>)			x		x
spruce, Colorado blue (<i>Picea pungens</i>)	glauca	x	x	x	x
spruce, Norway (<i>Picea abies</i>)			x		x
spruce, white (<i>Picea glauca</i>)	conica		x		x
sweetflag grass (<i>Acorus calimus</i>)			x		x
sweetgum (<i>Liquidambar styraciflua</i>)			x		x
sycamore, American (<i>Platanus occidentalis</i>)			x		x

35 8 61

tea tree, New Zealand (<i>Leptospermum scoparium</i>)	ruby glow	x	x	x	x
tree fern (tiki fern) (<i>Asparagus virgatus</i>)			x		x
trumpet flower or Carolina Jessamine (<i>Gelsemium sempervirens</i>)			x		x
tulip (<i>Tulip spp</i>)	apeldoorn		x		x
tufted hairgrass (<i>Deschampsia caespitosa</i>)			x		x
verbena, shrub (<i>Lantana sellowiana</i>)			x		x
Verbena, St. Paul's (<i>Verbena peruviana.</i>)	St. Paul		x		x
viburnum (<i>Viburnum spp.</i>)	American cranberry bush arrowwood European cranberry bush linden Mohican wright		x x x x x x		x x x x x x
vinca (periwinkle) (<i>Vinca minor</i>)			x		x
windmill palm (<i>Trachycarpus fortunei</i>)			x		x
xylosma (<i>Xylosma congestum</i>)			x		x
yarrow (<i>Achillea spp.</i>)			x		x
yaupon (<i>Ilex vomitoria</i>)	dwarf		x		x
yew (<i>Taxus cuspidata</i>) (<i>Taxus x media</i>)	capitata denisiformis	x	x x	x	x x

† Ornamental species only. Do not use on plants grown for food.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

EPA accepted: __/__/__

[Sub Label B: Residential Use]

(Base Label:)

Dimension[®] 2EW

Specialty Herbicide

Provides control of listed annual grasses and broadleaf weeds in:

- **Established lawns**
- **Ornamental turf**
- **Container and landscape ornamentals**

In New York State, do not use more than 1 pint of this product per acre per year (equivalent to 0.25 lb of active ingredient per acre).

Controls [tough] weeds [such as crabgrass] [barnyardgrass] [annual bluegrass] [barley] [brome] [crowfootgrass] [seedling dallisgrass] [giant foxtail] [green foxtail] [yellow foxtail] [goosegrass] [kikuyugrass] [Mary's grass] [wild oats] [annual and perennial ryegrass] [sandbur] [smutgrass] [bittercress] [carpetweed] [chickweed] [Carolina geranium] [henbit] [prostrate knotweed] [common lespedeza] [maretail] [black medic] [mulberry weed] [mustard] [oxalis] [pineappleweed] [parsley-piert] [redroot pigweed] [common purslane] [London rocket] [shepherdspurge] [corn speedwell] [garden spurge] [prostrate spurge] [spotted spurge] [creeping woodsorrel] [yellow woodsorrel]

and/or

Season Long Crabgrass Control [non-staining]

and/or

Season Long Crabgrass Prevention [non-staining]

and/or

Easy to use

and/or

Apply [early Spring], [Spring], [late Spring]

and/or

Apply in the fall for annual bluegrass, chickweed, henbit and other winter weeds

and/or

Use for a [beautiful] lawn [non-staining]

and/or

Provides early post emergent crabgrass control [non-staining]

and/or

Better crabgrass control / prevention than other leading brands [non-staining]

and/or

Effective crabgrass control /prevention leads to [thicker], [healthier] lawns [non-staining]

and/or

Prevents crabgrass and other weeds like [spurge] ,[oxalis] from germinating [non-staining]

and/or

[Prevents] [Controls] crabgrass [non-staining]

and/or

For use on all turf types

and/or

For use on cool season and warm season turf grass

and/or

Just one application needed

and/or

More than a crabgrass preventer, also a [spurge] [oxalis] preventer

Active Ingredient

dithiopyr: 3,5- pyridinedicarbothioic acid, 2-(difluoromethyl)- 4-(2-methylpropyl)-6- (trifluoromethyl)-S,S-dimethyl ester.....	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates

Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear long-sleeved shirt and long pants, chemical-resistant gloves \geq 14 mils (such as barrier laminate or butyl rubber), and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC
Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents__

(cover)

Dimension[®] 2EW

Specialty Herbicide

Provides control of listed annual grasses and broadleaf weeds in:

- Established lawns
- Ornamental turf
- Container and landscape ornamentals

In New York State, do not use more than 1 pint of this product per acre per year (equivalent to 0.25 lb of active ingredient per acre).

[Editor's note: One or more of the following statements may be included on the label.]

Controls [tough] weeds [such as crabgrass] [barnyardgrass] [annual bluegrass] [barley] [brome] [crowfootgrass] [seedling dallisgrass] [giant foxtail] [green foxtail] [yellow foxtail] [goosegrass] [kikuyugrass] [Mary's grass] [wild oats] [annual and perennial ryegrass] [sandbur] [smutgrass] [bittercress] [carpetweed] [chickweed] [Carolina geranium] [henbit] [prostrate knotweed] [common lespedeza] [maretail] [black medic] [mulberry weed] [mustard] [oxalis] [pineappleweed] [parsley-piert] [redroot pigweed] [common purslane] [London rocket] [shepherdspurse] [corn speedwell] [garden spurge] [prostrate spurge] [spotted spurge] [creeping woodsorrel] [yellow woodsorrel]

and/or

Season Long Crabgrass Control [non-staining]

and/or

Season Long Crabgrass Prevention [non-staining]

and/or

Easy to use

and/or

Apply [early Spring], [Spring], [late Spring]

and/or

Apply in the fall for annual bluegrass, chickweed, henbit and other winter weeds

and/or

Use for a [beautiful] lawn [non-staining]

and/or

Provides early post emergent crabgrass control [non-staining]

and/or

Better crabgrass control / prevention than other leading brands [non-staining]

and/or

Effective crabgrass control /prevention leads to [thicker], [healthier] lawns [non-staining]

and/or

Prevents crabgrass and other weeds like [spurge] ,[oxalis] from germinating [non-staining]

and/or

[Prevents] [Controls] crabgrass [non-staining]

and/or

For use on all turf types

and/or

For use on cool season and warm season turf grass

and/or

Just one application needed

and/or

More than a crabgrass preventer, also a [spurge] [oxalis] preventer

Active Ingredient

dithiopyr: 3,5- pyridinedicarbothioic acid, 2-(difluoromethyl)- 4-(2-methylpropyl)-6- (trifluoromethyl)-S,S-dimethyl ester.....	24%
Other Ingredients	76%
Total	100%

Contains petroleum distillates

Contains 240 grams per liter or 2 lb active ingredient per U.S. gallon.

Product protected by U.S. Patent No. 4,692,184. Other patents pending

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.
If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-542

EPA Est. _____

®Trademark of Dow AgroSciences LLC
Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents__

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes Skin Irritation • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or on clothing. Avoid contact with eyes. Wear long-sleeved shirt and long pants, chemical-resistant gloves ≥ 14 mils (such as barrier laminate or butyl rubber), and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Not for use on plants being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or on lawns and grounds.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area. Store this product above 32°F to avoid crystallization. If crystals form or product freezes, move product to area with ambient temperature above 32°F and shake well until crystals have dissolved.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

General Information

Dimension® 2EW specialty herbicide provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns, ornamental turf, and container and landscape ornamentals.

This product will not control established weeds, except for emerged crabgrass, up to initiation of tillering. This stage of growth for crabgrass generally corresponds to the time when crabgrass seedlings are first visible in established turfgrasses and individual plants have five leaves or less. Applications to crabgrass after initiation of tillering will not provide satisfactory control. All other applications of this product should be made preemergence (prior to germination of target weeds).

This product is not effective until activated by 1/2 inch or more of rainfall or sprinkler irrigation. Applications should be timed to ensure that activation has occurred prior to tillering stage of crabgrass development or prior to germination of all other weeds.

Note: In New York State, do not use more than 1 pint of this product per acre per year (equivalent to 0.25 lb of active ingredient per acre).

Keep unprotected persons out of treated area until sprays have dried.

Mixing Directions

Apply Dimension 2EW with a hand-held pump-up or backpack sprayer in 1 to 2 gallons of water per 1000 sq ft. Fill the clean spray tank to about three-fourths of the required volume of water for the area being treated. Add the required amount of Dimension 2EW to the spray tank and complete the filling process.

Recommended Use Rates

Application Rate (active ingredient/acre)	Equivalent Rate per 1000 sq ft [†]		
	Pint (pt)	Teaspoons (tsp)	Ounces (oz)
0.25 lb	0.02	2.2	0.32
0.5 lb	0.04	4.4	0.64

[†] 1000 sq ft is equal to an area 100 ft X 10 ft or square area about 32 ft on a side.

Equivalent Rates for Treatment of Small Areas:

Rate (pt per 1000 sq ft)	Equivalent Rate per 500 sq ft			Equivalent Rate per 100 sq ft		
	(pt)	(tsp)		(pt)	(tsp)	(oz)
0.04	0.02	1.9	0.32	0.004	0.38	.064
0.02	0.01	0.9	0.16	0.002	0.19	.032

Grass and Broadleaf Weeds Controlled by Dimension 2EW

Used as directed, Dimension 2EW controls annual grass and broadleaf weeds listed below if applied preemergence. This product will not control emerged broadleaf weeds or grasses (except for crabgrass

prior to tillering stage of growth). Therefore, the area to be treated must be free of weeds prior to application.

Grasses

barley	foxtail, green
barnyardgrass	foxtail, yellow
bluegrass, annual*	goosegrass
brome	kikuyugrass
crabgrass, large	Mary's grass
crabgrass, smooth	oats, wild
crabgrass, southern	ryegrass (annual & perennial)
crowfootgrass	sandbur
dallisgrass (seedling)	smutgrass
foxtail, giant	

Broadleaf Weeds

bittercress	parsley-piert
carpetweed	pigweed, redroot
chickweed	pineappleweed
geranium, Carolina	purslane, common
henbit	rocket, London
knotweed, prostrate	shepherdspurge
lespedeza, common	speedwell, corn
marestail	spurge, garden
medic, black	spurge, prostrate
mulberry weed	spurge, spotted
mustard	woodsorrel, creeping
oxalis, buttercup	woodsorrel, yellow

*Extended Control of Annual bluegrass in South and Coastal South Regions

For extended control of annual bluegrass in the south and coastal south regions, an initial application of 4.4 teaspoons of Dimension 2EW per 1000 sq ft (0.5 lb active ingredient) 6 to 8 weeks before overseeding with perennial ryegrass may be followed by a second application of 2.2 to 4.4 teaspoons of Dimension 2EW per 1000 sq ft (0.25 to 0.5 lb active ingredient) 120 days after overseeding. Minimum seeding rate of 9 lbs perennial ryegrass seed per 1000 sq. ft. Some injury to overseeded perennial ryegrass may occur (see Reseeding, Overseeding, or Sprigging precautions under Turfgrass).

Uses

Turfgrass

Use Dimension 2EW on seeded, sodded, or sprigged lawns and ornamental turfgrass that are well established.

Specific Use Precautions and Restrictions:

Newly established turf following seeding, sodding, or sprigging must have developed a good root system and a uniform stand, and should have received at least two mowings before the first application of this product. Use of this product on turf that is not well-established, or has been weakened by weather, pest, disease, chemical, mechanical or other related stress, may result in turf injury.

- Dimension will prevent the germination of annual bluegrass. Dimension will not affect established annual bluegrass. If maintenance of annual bluegrass is desired, using this product during time of annual bluegrass germination is not recommended. In the states of AZ, CA, NV, OR, WA, NM, ID,

MT and UT Dimension may contribute to the thinning or stand reduction in established stands of annual bluegrass.

- It is recommended that sod be established for at least 6 months before applying Dimension 2EW
- Early postemergence applications of this product will control crabgrass only if applied prior to the fifth leaf (first tiller) stage of growth.
- For best results, cultural practices that disturb the soil, such as verticutting and core-, spike-, or hydro-aerification, should be done before applying this product.
- Do not apply this product until the turfgrass has recovered from cultural practices such as verticutting or core-, spike-, or hydro-aerification.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not apply more than 4.4 teaspoons (0.73 oz) of Dimension 2EW per 1000 sq ft per application.
- Do not apply more than 13.2 teaspoons (2.2 oz) of Dimension 2EW per 1000 sq ft per year using split or sequential applications.
- In New York State, do not use more than 1 pint of this product per acre per year (equivalent to 0.25 lb of active ingredient per acre).

Dimension 2EW should be used only on the following turfgrass species that have been determined to be tolerant to this product:

Established Cool Season Turfgrasses

bentgrass, creeping[†]
bluegrass, Kentucky
fescue, fine^{††}
fescue, tall
ryegrass, perennial

Established Warm Season Turfgrasses

bahiagrass
Bermudagrass^{†††}
buffalograss^{††††}
carpetgrass
centipedegrass
kikuyugrass
seashore paspalum
St. Augustinegrass
zoysiagrass

[†] Do not use this product on certain varieties of creeping bentgrass, such as cohansey, carmen, seaside, and Washington as undesirable turfgrass injury may result. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (*Agrostis tenuis*) varieties.

^{††} Do not use this product on certain varieties of fine fescue as undesirable turf injury may result. The following fine fescue varieties have been found to be sensitive to this product: Atlanta, banner, beauty, bilgart, CF-2, enjoy, HF-93, highlight, ivalo, Jamestown, koket, majenta, Mary, pennlawn, Tamara, Tatjana, waldorf, and waldina. Not all varieties of fine fescue have been tested.

^{†††} Do not use this product on Tifgreen (328) hybrid Bermudagrass as undesirable turfgrass injury may result. Other common and hybrid Bermudagrass varieties are tolerant.

^{††††} Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Reseeding, Overseeding, or Sprigging

Reseeding, overseeding, or sprigging of treated areas within 3 months after a single application of this product, or within 4 months after a split application program totaling more than 4.5 teaspoons per 1000 sq ft, may inhibit the establishment of desirable turfgrasses. However, overseeding of Bermudagrass with perennial ryegrass 8 weeks after an application or as early as 6 weeks after application if slight injury to perennial ryegrass can be tolerated is a recommended exception.

Application Directions

Apply uniformly to the treatment area as a coarse spray. Avoid streaking, skips, or excess overlaps during application. For control or suppression of grass and broadleaf weeds, apply Dimension 2EW as a preventative (preemergence) treatment prior to their germination and emergence in the spring, summer or fall.

Crabgrass Control

Spring Application: Dimension 2EW provides preemergence control of crabgrass (including the large, smooth, and southern species) when applied prior to crabgrass emergence in established lawns and ornamental turf. This product also provides early postemergence control of crabgrass during early crabgrass seedling growth following emergence. However, it is often difficult to observe very small crabgrass seedlings at early stages of growth in well-established turfgrass. Early postemergence crabgrass control is obtained only when this product is applied prior to the tillering (spreading) of crabgrass (less than 5 leaves per plant), which generally corresponds to the time when crabgrass seedlings are easily observed in lawn or turf. The practical benefit of this product's additional, early postemergence activity is a "window of application" during which control of crabgrass can be achieved. That is 2 to 8 weeks longer than products that provide only preemergence control. Depending upon climatic conditions and rate of growth, the time from crabgrass emergence until tillering will vary from 2 to 8 weeks.

Fall Application: This product can also be applied in the late summer or early fall (late August through November) for preemergence control of all listed weeds. A fall application followed by an appropriately timed spring application provides season-long weed control. Do not exceed the maximum use rate per year.

Ornamentals (Landscape and Container Grown) Use

Dimension 2EW provides preemergence control of listed annual grasses and broadleaf weeds in areas planted with tolerant ornamental plants listed on this label. It is intended for use on plants grown for aesthetic purposes in landscaped areas.

Specific Use Precautions and Restrictions:

- Apply Dimension 2EW to established ornamentals only.
- Do not apply this product to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into the soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, heat stress, nematodes or diseases).
- Do not apply when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants unless they are listed on this label.
- Do not apply this product directly to plants that are grown for food (e.g., fruit trees or maple trees tapped for syrup).
- Do not apply this product in enclosed structures and greenhouses.
- Do not apply more than 4.4 teaspoons (0.73 oz) of Dimension 2EW per 1000 sq ft per application and no more than 13.2 (2.2 oz) teaspoons of Dimension 2EW per 1000 sq ft per year using split or sequential applications.
- In New York State, do not use more than 1 pint of this product per acre per year (equivalent to 0.25 lb of active ingredient per acre).

Treatment of Ornamental Species Not Listed on the Label for Dimension 2EW: It is impossible to evaluate tolerance to this product on all ornamental plant species or varieties or under all possible

growing conditions. Users who wish to use Dimension 2EW on ornamental species not recommended on this label may determine the suitability for use by treating a small number of ornamental plants at a recommended rate. Prior to treatment of larger areas, treated plants should be observed for any symptoms of herbicidal injury, such as foliar damage, reduced vigor or stand reduction, during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

Application Directions

Apply Dimension 2EW as a spray in established ornamentals or as a broadcast over-the-top spray to certain established ornamentals (see ornamental plant listing). Make directed sprays to the soil at the base of the ornamentals.

To reduce injury potential:

- Apply product with calibrated equipment using a minimum of 2 gallons of water per 1000 sq. ft.
- Shortly after application apply overhead irrigation to activate the herbicide and wash Dimension 2EW from plant surface onto soil surface.
- In the spring when buds are rapidly growing and expanding, over the top application of Dimension 2EW may temporarily injure new growth of desirable plants. To reduce the possibility of injury at this time, wait to apply Dimension 2EW over the top of newly emerged vegetation until it has hardened off, unless your experience indicates that the ornamental plant will not be injured by the over the top application.
- Do not apply to plants that are under stress such as heat, drought or frost damage.

Dimension 2EW is a preemergence herbicide that controls weeds during germination. Dimension 2EW does not control emerged broadleaf or grass weeds except crabgrass up to tillering stage of growth. Apply prior to germination of target weeds to bare ground. Optimum weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves. Prior to applying, control existing vegetation by cultivation, hand weeding, or use of a postemergence herbicide. After applying Dimension 2EW, excessive soil disruption may breakdown the herbicide barrier. Minimal surface disruption such as raking should not break down the herbicide barrier but should not be done until after the herbicide has been activated with 1/2 inch of rainfall or irrigation. Following transplanting, care must be taken that soil or planting mixes have settled firmly through packing, irrigation or rainfall and that there are no cracks that would allow direct contact of this product to the plant roots. Make sequential applications at 3 to 4 month intervals for extended preemergence weed control. Do not exceed maximum use rates per year

Tolerant Ornamentals

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials.

Name	Tolerant Cultivars	Acceptable Application Method Noted by a (X)		Application to Field/Landscape or Container Grown Ornamentals Noted by a (X)	
		Over the Top	Directed	Container Grown	Field/Landscape Grown
abelia, dwarf (<i>Abelia x grandiflora</i>)	nana		X		X
abyssinian red banana (<i>Ensete ventricosum</i>)	maureli	X	X	X	X
Ajuga carpet bugle (<i>Ajuga reptans</i>) (<i>Ajuga genevensis</i>)	bronze bronze beauty		X X		X X

almond, flowering (<i>Prunus gladiosa</i>)			x		x
apple [†] (<i>Malus pumila</i>)			x		x
arborvitae (<i>Thuja occidentalis</i>)	nigra pyramidalis smaragh techny woodwardii		x x x x x		x x x x x
arborvitae, dwarf golden (<i>Thuja orientalis</i>)	aurea nana	x	x	x	x
ash, green (<i>Fraxinus pennsylvanica</i>)			x		x
ash, autumn purple (<i>Fraxinus americana</i>)	autumn purple		x		x
aster, Chinese (<i>Callistephus chinensis</i>)	dwarf queen		x		x
azalea (<i>Rhododendron</i> spp.)	brilliant buccaneer carror chimes (Belgian) Elsie Lee exbury fashion hardijzer beauty hershey red higasa hinocrimson Holland (hybrid) Marion Lee northern lights orange cup orchid lights snow southern charm		x x x x x x x x x x x x x x x x x x x		x x x x x x x x x x x x x x x x x x x
azalea, flame (<i>Rhododendron calendulaceum</i>)			x		x
azalea, kirishima (<i>Rhododendron obtusum</i>)			x		x
bamboo, heavenly (<i>Nandina domestica</i>)	compacta nana		x x		x x
barberry, Japanese (<i>Berberis thunbergii</i>)	aurea dwarf pygmy green kobold pygmy red rose glow		x x x x x x		x x x x x x
barberry, purple (<i>Berberis thunbergii</i> var <i>atropurpurea</i>)	atropurpurea		x		x
basket flower (<i>Gaillardia grandiflora</i>)			x		x

beach grass (<i>Ammophila breviligulata</i>)			X		X
bearberry (common) (<i>Arctostaphylos uva-ursi</i>)	Massachusetts		X		X
bee balm (<i>Monarda didyma</i>)			X		X
begonia (<i>Begonia spp.</i>)			X		X
birch, river (<i>Betula nigra</i>)			X		X
blackeyed Susan (<i>Rudbeckia hirta</i>)	goldstrum		X		X
blanket flower (<i>Gaillardia spp.</i>)			X		X
blue fescue (<i>Festuca ovina</i>)			X		X
blueberry† (<i>Vaccinium spp.</i>)	bluecrop blue jay Jersey north blue northland		X X X X X		X X X X X
bottlebrush (<i>Callistemon citrinus</i>)	Little John	X	X	X	X
bougainvillea (<i>Bougainvillea spectabilis</i>)	James Walker Scarlet O' Hara	X	X X	X	X X
boxwood (<i>Buxus microphylla</i>)	japonica		X		X
boxwood, green beauty (<i>Buxus microphylla japonica</i>)	green beauty	X	X	X	X
boxwood, welleri (<i>Buxus sempervirens</i>)	winter gem	X	X	X	X
broom (<i>Cytisus scoparius</i>) (<i>Genista pilosa</i>)	moonlight Vancouver gold		X X		X X
camellia (<i>Camellia japonica</i>) (<i>Camellia sasanqua</i>)	debutante mathotiana supreme chansonette setsukgekka	X X	X X X X	X X	X X X X
candytuft (<i>Iberis sempervirens</i>)	snow white		X		X
carex, variegated (<i>Carex spp.</i>)			X		X
carpet bugle (<i>Ajuga reptans</i>) (<i>Ajuga genevensis</i>)	bronze bronze beauty		X X		X X
cedar, red (<i>Juniperus virginiana</i>)			X		X
celosia (<i>Celosia spp.</i>)			X		X
centaura (<i>Centaurea montana</i>)			X		X

cockscorb, plumosa (<i>Celosia cristata</i>)	scarlet plumosa		X		X
coleus (<i>Coleus blumei</i>)	red kewpie		X		X
columbine (<i>Aquilegia spp.</i>)			X		X
coneflower, purple (<i>Echinacea purpurea</i>)	magnus purple	X	X	X	X
copper leaf (<i>Acalypha wilkesiana</i>)			X		X
coreopsis (<i>Coreopsis spp.</i>)	moonbeam		X		X
corn flower (<i>Centaurea spp.</i>)			X		X
cotoneaster (<i>Cotoneaster apiculatus</i>)			X		X
coyotebrush (<i>Baccharis pilularis</i>)			X		X
cycad (<i>Cycas revoluta</i>)			X		X
cypress, bald (<i>Taxodium distichum</i>)			X		X
cypress, Italian (<i>Cupressus sempervirens</i>)	glauca		X		X
cypress, Japanese false (<i>Chamaecyparis obtusa</i>)	gracilis		X		X
cypress, leyland (<i>Cupressocyparis leylandii</i>) hybrid			X		X
daffodil (<i>Narcissus spp.</i>)	King Alfred		X		X
daylily (<i>Heemerocallis spp.</i>)	Aztec gold bright yellow (hybrid) single gold (evergreen) wilsonís yellow		X X X X		X X X X
dianthus (sweet William) (<i>Dianthus spp.</i>)			X		X
delphinium (<i>Delphinium spp.</i>)	magic fountain		X		X
dogwood (<i>Cornus florida</i>)			X		X
dogwood, American (<i>Cornus sericea</i>)	flavarimea		X		X
Douglas fir (<i>Pseudotsuga menziesii</i>)			X		X
dusty miller (<i>Senecio cineraria</i>)		X	X	X	X
elm (<i>Ulmus parvifolia</i>)	drake		X		X
eulaliagrass (<i>Miscanthus sinensis</i>)	maiden grass	X	X	X	X

euonymus (<i>Euonymus japonicus</i>)	Argenteo-variegata		X		X
	Auereomarginata		X		X
	colorata		X		X
(<i>Euonymus fortunei</i>)	emerald gaiety		X		X
	emerald n' gold		X		X
	gold edge		X		X
	gold princess		X		X
	Manhattan	X	X	X	X
	silver king		X		X
	tricolor		X		X
	vegetus		X		X
europs, green leaved (<i>Euryops pectinatus</i>)	viridis	X	X	X	X
fan palm, European (<i>Chamaerops humilis</i>)			X		X
fan palm, Mexican (<i>Washingtonia robusta</i>)			X		X
fern (various) (<i>Asparagus spp.</i>)			X		X
fescue (<i>Festuca glauca</i>)			X		X
fetterbush (<i>Leucothoe fontanesiana</i>)	rainbow		X		X
fig (<i>Ficus retusa</i>)	nitida		X		X
fir fraser (<i>Abies fraseri</i>)			X		X
Forsythia (<i>Forsythia x 'Arnold Dwarf</i>)	arnold dwarf		X		X
(<i>Forsythia viridissima</i>)	bronxensis dwarf		X		X
(<i>Forsythia xintermedia</i>)	lynwood gold		X		X
(<i>Forsythia x 'Meadowlark'</i>)	meadowlark		X		X
(<i>Forsythia x intermedia</i>)	spring glory	X	X	X	X
(<i>Forsythia suspensa</i>)	weeping		X		X
fountain grass, purple (<i>Pennisetum setaceum</i>)	rubrum	X	X	X	X
fuchsia (<i>Fuchsia spp.</i>)			X		X
galium (<i>Galium odoratum</i>)			X		X
gardenia (<i>Gardenia jasminoides</i>)	mystery		X		X
	radicans		X		X
	white gem	X	X	X	X
geranium (<i>Pelargonium x hortorum</i>)			X		X
gum (<i>Eucalyptus citriodora</i>)			X		X

hawthorn (<i>Crataegus spp.</i>)	cockspur white crimson cloud enchantress Jack Evans Washington white		X X X X X		X X X X X
hawthorn, Indian (<i>Rhaphiolepis indica</i>)	ballerina	X	X	X	X
heather, twisted (<i>Erica cinerea</i>)	Mediterranean pink		X		X
hemlock, Canada (<i>Tsuga canadensis</i>)			X		X
hibiscus (<i>Hibiscus spp.</i>)	blue bird brilliant hula girl		X X X		X X X
holly (<i>Ilex x 'Nellie R. Stevens'</i>) (<i>Ilex x attenuata</i>)	Nellie R. Stevens fosteri Savannah	X	X X X	X	X X X
holly, blue (<i>Ilex x meserveae</i>)	blue boy blue girl China girl		X X X		X X X
holly, cassine (<i>Ilex cassine</i>)		X	X	X	X
holly, Chinese (<i>Ilex cornuta</i>)	burfordii needlepoint		X X		X X
holly, Japanese (<i>Ilex crenata</i>)	compacta hellerie Japanese northern beauty steeds	X	X X X X	X	X X X X
holly, yaupon (<i>Ilex vomitoria</i>)			X		X
honeysuckle (<i>Lonicera xylosteum</i>) (<i>Lonicera japonica</i>) (<i>Lonicera tatarica</i>)	Claveyi dwarf halliana Canadian white Zabelli		X X X X		X X X X
hosta (<i>Hosta sieboldii</i>) (<i>Hosta lancifolia</i>)	albo marginata		X X		X X
ice plant (<i>Carpobrotus edulis</i>)		X	X	X	X
ice plant, white trailing (<i>Delosperma alba</i>)		X	X	X	X
impatiens (<i>Impatiens spp.</i>)			X		X
iris (<i>Iris spp</i>)	dwarf blue wedgewood		X X		X X
ivy, English (<i>Hedera helix</i>)	Bulgaria thorndale		X X		X X
jasmine, Asian (<i>Trachelospermum asiaticum</i>)			X		X

jasmine, star (<i>Trachelospermum jasminoides</i>)			x		x
juniper (<i>Juniperus spp.</i>)	Arcadia		x		x
	Armstrong		x		x
	bar harbor		x		x
	blue chip		x		x
	blue Pacific		x		x
	blue rug		x		x
	blue star		x		x
	broadmoor		x		x
	buffalo		x		x
	calgary carpet		x		x
	emerald sea		x		x
	emerald spreader		x		x
	endora compacta		x		x
	fruitlandi		x		x
	green		x		x
	gold tip		x		x
	hetzi		x		x
	hughes		x		x
	Manhattan blue		x		x
	parsoni		x		x
	pfitzeriana		x		x
	plumosa		x		x
	Prince of Wales		x		x
	procumbens dwarf		x		x
	San Jose		x		x
	sargent blue		x		x
	sargent green		x		x
	scandia		x		x
	scopulorum moonglow		x		x
	spartan		x		x
	tamariscifolia		x		x
	weberi		x		x
	Youngstown		x		x
	Yukon belle		x		x
juniper (<i>Juniperus chinensis</i>)	gold coast	x	x	x	x
	torulosa – Hollywood twisted	x	x	x	x
juniper (<i>Juniperus procumbens nana</i>)	prostrate Japanese garden	x	x	x	x
juniper (<i>Juniperus sabina 'tamariscifolia'</i>)	tamarix	x	x	x	x
juniper (<i>Juniperus scopulorum</i>)	scopulorum skyrocket	x	x	x	x
king palm (<i>Archontophoenix cunninghamiana</i>)		x	x	x	x
laurel, Carolina cherry (<i>Prunus caroliniana</i>)	bright n' tight	x	x	x	x

laurel, mountain (<i>Kalmia latifolia</i>)			X		X
leucothoe (<i>Leucothoe fontanesiana</i>)			X		X
ligustrum, Japanese (<i>Ligustrum japonicum</i>)			X		X
lily (<i>Agapanthus spp.</i>)	streamline		X		X
Lily of the Nile (<i>Agapanthus africanus</i>)	albus Peter Pan		X X		X X
lilyturf (<i>Liriope muscari</i>)	evergreen giant lilac beauty majestic monroe white silvery sunproof variegata	X	X X X X X	X	X X X X X
lilyturf, creeping (<i>Liriope spicata</i>)			X		X
maple, amur (<i>Acer ginnala</i>)	emerald elf	X	X	X	X
maple, Japanese (<i>Acer palmatum</i>)			X		X
maple, Norway (<i>Acer platanoides</i>)			X		X
maple, red [†] (<i>Acer rubrum</i>)	red sunset	X	X	X	X
maple, silver (<i>Acer saccharinum</i>)			X		X
maple sugar [†] (<i>Acer saccharum</i>)			X		X
marigold (<i>Tagetes patula</i>)	honeycomb variegata wheeleris dwarf		X X X		X X X
mock orange [†] (<i>Philadelphus spp</i>)	golden snowflake double white		X X		X X
mondo grass (<i>Ophiopogon japonicus</i>)			X		X
moss rose (<i>Portulaca grandiflora</i>)	sunnyside		X		X
mountainash (<i>Sorbus aucuparia</i>)			X		X
myrtle, crape (<i>Lagerstroemia indica</i>)	faurei langer muskogee standard pink		X X X X		X X X X
myrtle, wax (<i>Myrica californica</i>)			X		X
narcissus (<i>Narcissus spp.</i>)			X		X
oak, laurel (<i>Quercus laurifolia</i>)			X		X
oak, pin (<i>Quercus palustris</i>)			X		X

oak, red (<i>Quercus rubra</i>)			X		X
oak, southern (<i>Quercus virginiana</i>)			X		X
oak, willow (<i>Quercus phellos</i>)			X		X
oleander (<i>Nerium oleander</i>)	hardy red petite pink sister agnes		X X X		X X X
osteospermum (<i>Osteospermum fruticosum</i>)	whirligig		X		X
pachysandra (<i>Pachysandra terminalis</i>)			X		X
palm, bangalow (<i>Archontophoenix cunninghamiana</i>)			X		X
palm, queen (<i>Syagrus romanzoffianum</i>)		X	X	X	X
pampas grass (<i>Cortaderia selloana</i>)			X		X
pansy (<i>Viola x wittrockiana</i>)			X		X
paper flower (<i>Bougainvillea glabra</i>)	Barbara Karst	X	X	X	X
peach [†] (<i>Prunus persica</i>)			X		X
periwinkle, dwarf (<i>Vinca minor</i>)			X		X
petunia (<i>Petunia x hybrida</i>)	picoti		X		X
photinia, red tip (<i>Photinia x fraseri</i>)			X		X
pieris (<i>Pieris taiwanensis</i>)			X		X
pieris, Japanese (<i>Pieris japonica</i>)	mountain fire		X		X
pine, Afghan (<i>Pinus eldarica</i>)		X	X	X	X
pine, Austrian black (<i>Pinus nigra</i>)		X	X	X	X
pine, Japanese black (<i>Pinus thunbergii</i>)			X		X
pine, loblolly (<i>Pinus taeda</i>)			X		X
pine, longleaf (<i>Pinus palustris</i>)			X		X
pine, mugo or Swiss Mt. (<i>Pinus mugo</i>)			X		X
pine, Scotch (<i>Pinus sylvestris</i>)			X		X
pine, slash (<i>Pinus elliotii</i>)			X		X

pine, Virginia (<i>Pinus virginiana</i>)			X		X
pine, white (<i>Pinus strobus</i>)		X	X	X	X
pineapple, guava [†] (<i>Feijoa sellowiana</i>)			X		X
pittosporum, tobira (<i>Pittosporum tobira</i>)	Wheeler's dwarf	X	X	X	X
plumbago, cape (<i>Plumbago auriculata</i>)	royal cape	X	X	X	X
potentilla (<i>Potentilla fruticosa</i>)) (<i>Potentilla nepalensis</i>)	abbotswood		X X		X X
privet (<i>Ligustrum x vicaryii</i>) (<i>Ligustrum japonicum</i>)	golden vicary regal texanum yellow tipped	X	X X X X	X	X X X X
privet, glossy (<i>Ligustrum lucidum</i>)			X		X
Pyracantha or firethorn (<i>Pyracantha x 'Gnome'</i>) (<i>Pyracantha coccinea</i>) (<i>Pyracantha koidzumii</i>)	gnome lalandei victory	X	X X X	X	X X X
queen palm (<i>Arecastrum rammanzoffianum</i>)			X		X
quince, Japanese [†] (<i>Chaenomeles japonica</i>)			X		X
Rhododendron (<i>Rhododendron spp</i>)	album Cunningham white PJM purple gem silvery pink		X X X X X		X X X X X
rhododendron, Carolina (<i>Rhododendron carolinianum</i>)			X		X
rhododendron, catawba (<i>Rhododendron catawbiense</i>)			X		X
rhododendron, rhodie max – rosebay (<i>Rhododendron maximum</i>)			X		X
ribbon grass (<i>Phalaris arundinacea</i>)			X		X
rockcress (<i>Arabis caucasica</i>)	snowcap		X		X
rose [†] (<i>Rosa banksiae</i>)	luta		X		X
rose, knockout shrub (<i>Rosa spp. hybrid</i>)	knockout	X	X	X	X
rosemary [†] (<i>Rosmarinus officinalis</i>)			X		X

rosemary, bog (<i>Andromeda polifolia</i>)	nana		X		X
salvia (<i>Salvia farinacea</i>)	rhea		X		X
sedum (<i>Sedum spurium</i>)	dragon blood red red carpet yellow		X X X		X X X
smoketree, royal purple (<i>Cotinus coggygia</i>)	royal purple		X		X
snapdragon (<i>Antirrhinum spp.</i>)			X		X
snowball, common (<i>Viburnum opulus</i>)	sterile	X	X	X	X
sourwood (<i>Oxydendrum arboreum</i>)			X		X
spiraea (<i>Astilbe X arendsii</i>)	fanall		X		X
spiraea (<i>Spiraea spp.</i>)	Anthony Waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white		X X X X X X		X X X X X X
spiraea, garland (<i>Spiraea x arguta</i> .)			X		X
spruce, Black Hills (<i>Picea glauca var densata</i>)			X		X
spruce, Colorado blue (<i>Picea pungens</i>)	glauca	X	X	X	X
spruce, Norway (<i>Picea abies</i>)			X		X
spruce, white (<i>Picea glauca</i>)	conica		X		X
sweetflag grass (<i>Acorus calimus</i>)			X		X
sweetgum (<i>Liquidambar styraciflua</i>)			X		X
sycamore, American (<i>Platanus occidentalis</i>)			X		X
tea tree, New Zealand (<i>Leptospermum scoparium</i>)	ruby glow	X	X	X	X
tree fern (tiki fern) (<i>Asparagus virgatus</i>)			X		X
trumpet flower or Carolina Jessamine (<i>Gelsemium sempervirens</i>)			X		X
tulip (<i>Tulip spp</i>)	apeldoorn		X		X
tufted hairgrass (<i>Deschampsia caespitosa</i>)			X		X
variegated society garlic (<i>Thulbaghia violacea</i>)	variegata	X	X	X	X

verbena, shrub (<i>Lantana sellowiana</i>)			X		X
Verbena, St. Paul's (<i>Verbena peruviana</i> .)	St. Paul		X		X
viburnum (<i>Viburnum</i> spp.)	American cranberry bush arrowwood European cranberry bush linden Mohican wright		X X X X X X		X X X X X X
vinca (periwinkle) (<i>Vinca minor</i>)			X		X
windmill palm (<i>Trachycarpus fortunei</i>)			X		X
xylosma (<i>Xylosma congestum</i>)			X		X
yarrow (<i>Achillea</i> spp.)			X		X
yaupon (<i>Ilex vomitoria</i>)	dwarf		X		X
yew (<i>Taxus cuspidata</i>) (<i>Taxus x media</i>)	capitata denisiformis	X	X X	X	X X

† Ornamental species only. Do not use on plants grown for food.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or

2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

EPA accepted: __/__/__