 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 62719-551	Date of Issuance: JUL - 6 2006
	Term of Issuance: Conditional	
	Name of Pesticide Product: GF-1406	

Name and Address of Registrant (include ZIP Code):

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268


Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit within one year of the date of this registration notice, storage stability (830.6317) and corrosion characteristics (830.6320) studies.

Signature of Approving Official: 	Date: JUL - 6 2006
Joanne I. Miller, Product Manager (23) Herbicide Branch, Registration Division (7505P)	

3. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 62719-551."
- b. *According to your Basic Confidential Statement Formula (CSF), it appears that you have intentions to apply different food use inert ingredients in the formulation. A copy of the Technical Review Branch (TRB) analysis has been enclosed and details the issue more. Please note that the Agency must be notified and an alternate CSF must be submitted if inert ingredients present in the basic formulation are replaced with others that are similar. The Agency does not allow such phrases in the CSF. Agency review and approval of an alternate CSF is required prior to marketing the alternate formulation.*

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

3/27

(Base Label):

(Logo) Dow AgroSciences

GF-1406

Concentrated - Effective - Low Volatile

Contains 2-Ethylhexyl Ester of 2,4-D

For selective control of many broadleaf weeds in certain crops, including, cereal grains (wheat, barley, millet, oats and rye), corn (field corn, popcorn and sweet corn), fallow land and crop stubble, potatoes, sorghum (grain and forage sorghum), and soybeans (preplant burndown application only), forests, rangeland and established grass pastures, including Conservation Reserve Program (CRP) acres, non-cropland, grasses grown for seed or sod, and ornamental turf.

Active Ingredient(s):

2,4-Dichlorophenoxyacetic acid,	
2-Ethylhexyl Ester [†]	87.2%
Inert Ingredients	12.8%
Total	100.0%

2,4-Dichlorophenoxyacetic Acid Equivalent: 57.5% - 5.5 lb/gal
[†] Ester Specific Analytical Method, Dow AgroSciences, 1999.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
JUL - 6 2006**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Keep Out of Reach of Children

CAUTION PRECAUCION

62719-551

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed • Causes Moderate Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Wash thoroughly with soap and water before handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selections chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves, when applying postharvest dips or sprays to citrus, applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

4/27

- Chemical resistant apron when applying postharvest dips or sprays to citrus, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

For containers of 5 gallons or more: A mechanical system (such as probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Note: Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

Environmental Hazards

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

5/27

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read "Warranty Disclaimer," "Inherent Risks of Use," and "Limitation of Remedies" at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-xxx

EPA Est. _____

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents __ gal

6/27

(Datapack cover):

(Logo) **Dow AgroSciences**

GF-1406

Concentrated - Effective - Low Volatile

Contains 2-Ethylhexyl Ester of 2,4-D

For selective control of many broadleaf weeds in certain crops, including, cereal grains (wheat, barley, millet, oats and rye), corn (field corn, popcorn and sweet corn), fallow land and crop stubble, potatoes, sorghum (grain and forage sorghum), and soybeans (preplant burndown application only), forests, rangeland and established grass pastures, including Conservation Reserve Program (CRP) acres, non-cropland, grasses grown for seed or sod, and ornamental turf.

Active Ingredient(s):

2,4-Dichlorophenoxyacetic acid,	
2-Ethylhexyl Ester [†]	87.2%
Inert Ingredients	12.8%
Total	100.0%

2,4-Dichlorophenoxyacetic Acid Equivalent: 57.5% - 5.5 lb/gal

[†] Ester Specific Analytical Method, Dow AgroSciences, 1999.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read "Warranty Disclaimer," Inherent Risks of Use," and "Limitation of Remedies" at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-XXX

EPA Est. _____

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

7/27

Herbicide

Net Contents __ gal

8/27

[Page 1 through end]:

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment	-
Engineering Controls Statements	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Non-Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
General Use Precautions and Restrictions	-
Avoiding Injury to Non-target Plants	-
Spray Drift Precautions	-
Spray Drift Management	-
Aerial Drift Information Advisory	-
Mixing Instructions	-
Application Instructions	-
Weeds Controlled	-
Crop Uses	-
Cereal Grains (Wheat, Barley, Millet, Rye, Oats)	-
Conservation Reserve Program Acres (See Rangeland and Permanent Grass Pastures)	-
Corn (Field Corn, Popcorn and Sweet Corn)	-
Fallowland and Crop Stubble	-
Field Corn (See Corn)	-
Potatoes	-
Sorghum (Grain Sorghum (Milo) and Forage Sorghum)	-
Soybeans (Preplant Only)	-
Forestry, Rangeland, Established Pastures, and Non-cropland Uses	-
Forests	-
Rangeland, and Established Grass Pastures (Including Perennial Grasslands not in Agricultural Production Such as Conservation Reserve Program Acres)	-
Non-cropland	-
Turf Uses	-
Grasses Grown for Seed or Sod	-
Ornamental Turf	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

9/27

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Swallowed • Causes Moderate Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Wash thoroughly with soap and water before handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selections chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves, when applying postharvest dips or sprays to citrus, applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- Chemical resistant apron when applying postharvest dips or sprays to citrus, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

For containers of 5 gallons or more: A mechanical system (such as probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

19/27

First Aid

If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Note: Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

Environmental Hazards

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any water-proof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: When this product is applied to rangeland and established pastures not harvested for hay or seed; non-cropland areas, ornamental turf not grown for sod or seed, and when applied by tree injection method only in forest sites, do not enter or allow people or pets to enter the treated area until sprays have dried.

Storage and Disposal
Do not contaminate water, food, or feed by storage or disposal.
Storage: Keep container tightly closed when not in use. If exposed to subfreezing temperatures, the product should be warmed to at least 40°F and mixed thoroughly before using.
Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
Plastic Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by local authorities, by burning. If burned, stay out of smoke.
General: Consult federal, state, or local disposal authorities for approved alternative procedures.

General Information

GF-1406 herbicide is intended for selective control of many broadleaf weeds in certain crops, including, cereal grains (wheat, barley, millet, oats and rye), corn (field corn, popcorn and sweet corn), fallow land and crop stubble, potatoes, sorghum (grain and forage sorghum), and soybeans (preplant burndown application only), forests, rangeland and established grass pastures including Conservation Reserve Program (CRP) acres, non-cropland, grasses grown for seed or sod, and ornamental turf.

Apply GF-1406 as a water or oil-water spray during warm weather when weeds or woody plants are actively growing. Application under drought conditions will often give poor results. Use low spray pressure to minimize drift. Generally, the lower dosages recommended on this label will be satisfactory for young, succulent growth of susceptible weed species. For less susceptible species and under conditions where control is more difficult, use higher recommended rates. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeat applications for satisfactory control. Consult your State Agricultural Experiment stations or Extension Service Weed Specialists for recommendations from this label that best fit local conditions.

General use Precautions and Restrictions

Chemigation: Do not apply this product through any type of irrigation system.

Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination and plant growth.

Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Spray drift produced during application is the responsibility of the applicator and care should be taken to minimize off-target movement of spray during application. A drift control agent suitable for agricultural use may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

Do not apply where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use consumption. Do not apply GF-1406 directly to, or otherwise permit contact with cotton, okra, flowers, fruit trees, grapes (in growing stage),

12/27

ornamentals, soybeans (vegetative stage), sunflowers, tomatoes, beans, and other vegetables, tobacco or other desirable plants that are susceptible to 2,4-D herbicides. Do not permit spray mist containing 2,4-D to contact susceptible plants since even very small quantities of the spray, that may not be visible, can cause severe injury during both active growth and dormant periods. Do not use in greenhouses. At high temperatures vapors from this product may injure susceptible plants growing nearby. Avoid product volatilization that may occur when applying 2,4-D esters during conditions of low humidity and high temperatures.

Ground Equipment: All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Spray drift can be lessened by keeping the spray boom as low as possible, with a nozzle height not greater than 4 feet above the crop canopy; by applying 10 gallons or more of spray per acre; by using no more than 20 pounds spraying pressure and large droplet producing nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 15 miles per hour. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Droplet Size: Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray. For medium or more fine spray: apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles. For coarse spray: apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

Temperature Inversions: When wind speed less than 3 mph, do not make applications if conditions of temperature inversion exist, or stable atmospheric conditions exist at or below nozzle height.

Avoid Movement of Treated Soil: Avoid conditions under which soil from treated areas may be moved or blown to areas containing susceptible plants. Wind-blown dust containing 2,4-D may produce visible symptoms when deposited on susceptible plants; however, serious plant injury is unlikely. To minimize potential movement of 2,4-D on wind-blown dust, avoid treatment of powdery dry or light sandy soils until soil is settled by rainfall or irrigation or irrigate soon after application.

Other State and Local Requirements: Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Spray Drift Management - Aerial Application

The following additional drift management requirements must be followed to avoid off-target drift movement from aerial applications. These requirements do not apply to forestry or rights of way applications, public health uses or to applications using dry formulations. The applicator should be familiar with and take into account the information covered in the following Aerial Drift Advisory Information section.

Aerial Drift Advisory Information

Importance of Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size:

Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Application Height-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

14/27

Mixing Instructions

1. Fill the spray tank about half full with water, then add the required amount of GF-1406 Herbicide, with agitation, and finally the rest of the water.
Note: GF-1406 in water forms an emulsion, which tends to separate unless agitation is maintained.
2. If oil is added, first mix the GF-1406 Herbicide and the oil and then add this mixture to the water. However, with adequate agitation, the oil can be added after the GF-1406 Herbicide is mixed in the water.
3. If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil-herbicide mixture to avoid formation of an invert emulsion.

Note: Adding oil, wetting agent, or other surfactant to the spray mixture may increase effectiveness on weeds, but also may reduce selectivity in crops resulting in crop damage.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates. Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of this product and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Mixing with Liquid Nitrogen Fertilizer

This product may be combined with liquid nitrogen fertilizer suitable for foliar application to accomplish broadleaf weed control and fertilization of corn, small grains or pastures in a single operation. Use GF-1406 in accordance with recommendations for these crops provided in this label. Use liquid fertilizer at rates recommended by the supplier or Extension Service Specialist. Test for mixing compatibility by mixing spray ingredients in correct proportions in a clear glass jar before mixing in spray tank. A compatibility aid such as Unite or Compex may be needed in some situations. Compatibility is best with liquid fertilizer solutions containing only nitrogen. Mixing with N-P-K solutions may not be satisfactory, even with the addition of a compatibility aid. Pre-mixing GF-1406 with 1 to 4 parts water may help in situations when mixing difficulty occurs.

Sprayer Clean-Out

To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned before re-use or applying other chemicals.

1. Rinse and flush application equipment thoroughly after use at least three times with water. Dispose of all rinse water by application to treatment area or apply to non-cropland area away from water supplies.
2. During the second rinse, add 1 qt of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15-20 min). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Remove nozzles and screens and clean separately.
6. If equipment is to be used to apply another pesticide or agricultural chemical to a 2,4-D susceptible crop, additional steps may be required to remove all traces of 2,4-D, including cleaning of disassembled parts and replacement of hoses or other fittings that may contain absorbed 2,4-D.

Application Instructions

Spray Volume: Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions. For broadcast application, apply the recommended rate of this product in a spray volume of 2 or more gallons per acre by air and 10 or more gallons per acre for ground equipment. Use low-pressure sprays to minimize drift. Where states have regulations, that specify minimum spray volumes, they should be observed. In general, spray volume should be increased as crop canopy, height and weed density increase in order to obtain adequate spray coverage. **Do not apply less than 2 gallons total spray volume per acre.**

Application Rates: Generally, lower rates in recommended rate ranges will be satisfactory for more sensitive weeds species, when weeds are small, and when environmental conditions are favorable for rapid growth. Use higher rates in the recommended rate range for less sensitive species and under less favorable growing conditions. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on this label. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for effective control.

Spot Treatments

To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 sq ft as indicated below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of GF-1406. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1,000 sq ft. Mix the amount of GF-1406 (fl oz or ml) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of GF-1406 required for larger areas, multiply the table value (fl oz or ml) by the thousands of sq ft to be treated. An area of 1000 sq ft is approximately 10.5 X 10.5 yards (strides) in size.

Rate Conversion Table for Spot Treatment:

Label Broadcast Rate (pt/acre)							
1/3	1/2	2/3	1	1 1/3	2	2 2/3	5 1/3
Equivalent Amount of GF-1406 per 1000 sq ft							
1/8 fl oz † (3.7 ml)	1/5 fl oz (5.9 ml)	1/4 fl oz (7.4 ml)	3/8 fl oz (11 ml)	1/2 fl oz (15 ml)	3/4 fl oz (22 ml)	1 fl oz (30 ml)	2 fl oz (60 ml)

† Conversion factors: 1 pt - 16 fl oz.; 1fl oz = 29.6 (30) ml

Band Application: GF-1406 may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

Band width in inches

----- X Broadcast rate = Band rate per
Row width in inches per acre treated acre

Band width in inches

----- X Broadcast volume Band volume
Row width in inches per acre per treated acre

Weeds Controlled

Annual or Biennial Weeds

- | | |
|---|--------------------------------|
| beggarticks ⁽¹⁾ | mousetail ⁽²⁾ |
| bittercress, smallflowered ⁽²⁾ | mustards (except blue mustard) |
| bitterweed | parsnip, wild |

- broomweed, common ⁽¹⁾
- burdock, common
- buttercup, smallflowered ^{(1) (2)}
- carpetweed
- cinquefoil, common ⁽²⁾
- cinquefoil, rough ⁽²⁾
- cocklebur, common
- coffeeweed
- copperleaf, Virginia ⁽²⁾
- croton, Texas
- croton, woolly
- flixweed
- galinsoga
- geranium, Carolina ⁽²⁾
- hemp, wild
- horseweed (marestail) ⁽²⁾
- jewelweed
- jimsonweed
- knotweed ⁽¹⁾
- kochia
- lambquarters, common
- lettuce, prickly ^{(1) (2)}
- lettuce, wild
- lupines
- mallow, little ⁽¹⁾
- mallow, Venice ⁽¹⁾
- marshelder
- morningglory, annual
- morningglory, ivy
- morningglory, woolly
- pennycress (fanweed)
- pepperweeds (*Lepidium* spp.) ^{(1) (2)}
- pigweeds (*Amaranthus* spp.) ⁽¹⁾
- poorjoe
- primrose, common
- purslane, common ⁽²⁾
- pusley, Florida
- radish, wild
- ragweed, common
- ragweed, giant
- rape, wild
- rocket, yellow
- salsify, common ⁽¹⁾
- salsify, western ⁽¹⁾
- shepherdspurse
- sicklepod
- smartweed (annual species) ^{(1) (2)}
- sneezeweed, bitter
- sowthistle, annual
- sowthistle, spiny
- spanishneedles
- sunflower
- sweetclover
- tansymustard
- thistle, bull
- thistle, musk ⁽¹⁾
- thistle, Russian (tumbleweed) ⁽¹⁾
- velvetleaf
- vetches

Perennial Weeds

- Alfalfa ^{(1), (2)}
- artichoke, Jerusalem ⁽¹⁾
- aster, many-flower ⁽¹⁾
- Austrian fieldcress ⁽¹⁾
- bindweed (hedge, field and European) ^{(1) (2)}
- blue lettuce
- blueweed, Texas
- broomweed
- bullnettle ^{(1) (2)}
- carrot, wild ⁽¹⁾
- catnip
- chicory
- clover, red ^{(1) (2)}
- coffeeweed
- cress, hoary ⁽¹⁾
- dandelion ⁽¹⁾
- docks ⁽¹⁾
- dogbanes ⁽¹⁾
- goldenrod
- eveningprimrose, cutleaf ⁽²⁾
- garlic, wild ⁽¹⁾
- hawkweed, orange ⁽¹⁾
- healal
- ironweed, western ⁽²⁾
- ivy, ground ⁽¹⁾
- Jerusalem-artichoke
- loco, bigbend
- nettles (including stinging) ⁽¹⁾
- onion, wild ⁽¹⁾
- pennywort
- plantains
- ragwort, tansy ⁽¹⁾
- sowthistle, perennial
- thistle, Canada ^{(1) (2)}
- vervains ⁽¹⁾
- wormwood

⁽¹⁾ These weeds are only partially controlled and may require repeat applications and/or use of higher recommended rates of this product even under ideal conditions of application.

⁽²⁾ This product may not be used to control this weed species in the state of California.

17/27

Crop Uses

Agricultural Use Requirements for Crops: For the following crop uses, follow PPE and Re-entry instructions in the "Agricultural Use Requirements" section of this label.

Cereal Grains (Wheat, Barley, Millet, Oats, Rye) (Not Underseeded with Legumes)

Crop/Application Timing	GF-1406 (pt/acre)	Specific Use Directions
Wheat, Barley, Millet, Rye Annual and biennial broadleaf weeds Perennial broadleaf weeds	1/3 to 1 1/3 † 2/3 to 1 1/3 †	Apply after crop is fully tillered, but before boot stage of growth (usually 4 to 8 inches tall) but not forming joints in the stem. Do not apply before tillering or from early boot through the milk stage of growth.
Oats (Spring Seeded) (Fall Seeded Southern)	1/3 1/2 to 1 †	Apply after crop is fully tillered, but before boot stage of growth (usually 4 to 8 inches tall) and weeds are small. Do not apply before tillering or from early boot through the milk stage of growth. Do not apply during or immediately following cold weather.
Preharvest application (all cereals)	2/3	Apply using air or ground equipment to control weeds that could interfere with harvest, or to suppress perennial weeds. Apply when grain is in dough stage. Do not apply from early boot through the milk stage of growth.

† Use the lower rate in the rate range if small annual or biennial weeds are the major problem. Use the higher rate if perennial weeds or annual or biennial weeds are present which are considered to be hard-to-kill as determined by local experience. Higher rates increase the risk of crop injury and should be used only where weed control justifies such risk. Do not apply GF-1406 at the crop seedling stage of growth. Consult state agricultural experiment station or extension service weed specialists for recommendations or suggestions to fit local conditions.

Restrictions:

- **Preharvest Interval:** Do not harvest within 14 days of application.
- Do not apply more than 2 1/3 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

Corn (Field Corn, Popcorn and Sweet Corn)

Application Timing/ Stage of Growth	GF-1406 (pt/acre)	Specific Use Directions
Preplant (Burndown) Preemergence (Field corn, popcorn, and sweet corn)	2/3 to 1 1/3	General: For best results, growth conditions should be favorable for active weed growth. Use high rate in rate range for less susceptible weeds, cover crops such as alfalfa, weeds in advanced stages of development, or under less favorable growth conditions. Preplant: Apply 7 to 14 days before planting corn to control emerged broadleaf weed seedlings or existing cover crops. Preemergence: Apply any time after planting, but before corn emerges to control broadleaf weed seedlings or existing cover crops. Do not use on light sandy soils.
Postemergence (Field corn, popcorn, and sweet corn) Annual broadleaf weeds Crop up to 8 inches tall Crop 8 inches tall to tasseling (directed spray only) Perennial broadleaf weeds	1/3 to 2/3 2/3 2/3	Apply when weeds are small and corn is less than 8 inches tall (to top of canopy). If corn is more than 8 inches tall, use drop nozzles to keep spray off foliage. Treat perennial weeds when they are in bud to bloom stage. Do not tank mix with atrazine, oil or other adjuvants. Do not apply from tasseling to hard dough stage. Note: Corn treated with 2,4-D may become temporarily brittle. Wind or cultivation may cause stem breakage during the period of time that corn is brittle. Sweet Corn: To minimize potential for crop injury, use only lowest rate in rate range.
Preharvest (Field corn and popcorn only)	up to 2	Apply after corn is in hard dough (or denting) stage. Do not apply to sweet corn.

Precautions:

- Preplant or preemergence applications to light sandy soils is not recommended.
- Corn hybrids vary in tolerance to 2,4-D. Some are easily injured. Apply only to varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.
- **Note:** Corn treated with 2,4-D may exhibit stem brittleness for 8 – 10 days following application. During this period, the crop is more susceptible to stem breakage from cultivation or wind.

Restrictions (Field Corn and Popcorn):

- **Preharvest interval:** Do not harvest for grain or fodder within 7 days after application.
- Do not apply more than 4.0 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

Restrictions (Sweet Corn):

- **Preharvest interval:** Do not harvest ears within 45 days after application.
- Do not make a postemergence application any less than 21 days after a prior application.
- Do not apply more than 2.0 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

19/27

Fallowland and Crop Stubble

Fallowland is idle land, postharvest to crops or between crops.

Type of Weeds	GF-1406 (pt/acre)	Specific Use Directions
Annual broadleaf weeds	2/3 to 1 1/3	Use a lower rate in the rate range when weeds are small (2 to 3 inches tall) and actively growing. Use a higher in the rate range when weeds are larger and under less favorable growth conditions.
Biennial broadleaf weeds	1 1/3 to 2 2/3	Apply when musk thistles or other biennial species are in the seedling to rosette stage and before development of flower stalks. The lower rate can be used in the spring during the rosette stage. Use the highest rate in the fall or after flower stalks have developed.
Perennial broadleaf weeds	1 1/3 to 2 2/3	Apply when perennial weeds are in bud to early bloom stage or while in good vegetative growth.
Wild garlic and onion in crop stubble	2 2/3	Apply to new regrowth of wild garlic or onion that occurs in the fall after harvest of other crops.

Precaution: For best weed control results, do not cultivate for at least 2 weeks after application or until top growth is dead.

Restrictions:

- **Preharvest Interval:** Do not cut forage or hay within 7 days of application.
- Do not apply within 30 days of a previous application.
- Do not apply more than 2 2/3 pt/acre of GF-1406 per use season.

Planting in Treated Areas

Labeled Crops: Within 29 days after an application of this product, plant only those crops listed on this or other registered 2,4-D labels. Follow more stringent limitations, if any, provided in directions for specific crops. Labeled crops may be at risk of crop injury or loss if planted soon after application, especially during the first 14 days. Degradation factors described below should be considered in weighing this risk.

Other Crops: All other crops may be planted 30 or more days after application without concern for illegal residues in the planted crop. However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

Degradation Factors: When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid breakdown of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application. Consult your local agricultural extension service or information about susceptible crops and typical conditions in your area.

Potatoes (Fresh Market Only)

Application Timing/ Stage of Growth	GF-1406	Specific Use Directions
Postemergence	1/10 pt (1.6 oz)/acre	Make first application when potatoes are in the pre-bud stage (about 7 to 10 inches high) and make a second application about 10 to 14 days later.

20/27

Restrictions:

- **Preharvest Interval:** Do not harvest within 45 days of application.
- Do not exceed two applications per crop.
- Do not apply more than 0.14 lbs a.e. (1/5 pt/acre) of GF-1406 per growing season.

Sorghum (Grain Sorghum (Milo) and Forage Sorghum)

Application Timing/ Stage of Growth	GF-1406 (pt/acre)	Specific Use Directions
Postemergence † Crop 6 - 8 inches tall Crop 8 - 15 inches tall (directed spray only)	1/3 to 2/3 1/2 to 2/3	Apply when sorghum is 6 to 15 inches tall. If sorghum more than 8 inches tall (top of canopy), use drop nozzles to keep spray off foliage. Do not use with oil or other adjuvants. Do not treat during boot, flowering or dough stage.

Precautions:

- **Note:** Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply GF-1406 under these conditions, use no more than 1/2 pint per acre.
- **Do not apply during boot, or later stages of growth.**
- Sorghum hybrids vary in tolerance to 2,4-D. Some are easily injured. Apply only to varieties known to be tolerant to 2,4-D. Consult the seed company or your agricultural experiment station or extension service weed specialist for this information.

Restrictions:

- **Preharvest Interval:** Do not harvest grain for 30 days after application.
- Do not permit meat or dairy animals to consume treated crop as fodder or forage within 30 days after application.
- Do not apply more than 2/3 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

Soybeans - For Use in Crop Residue Management Systems (Pre-plant Burndown Application Only)

Application Timing	GF-1406 (pt/acre)	Specific Use Directions
Preplant (Burndown)	1/2 to 2/3	Apply not less than 7 days before planting soybeans. See Use Precautions and Restrictions below.
	2/3 to 1 1/3	Apply not less than 15 days before planting soybeans. See Use Precautions and Restrictions below.
General Use Directions: Use GF-1406 to control emerged broadleaf weeds or existing cover crops. For best results, apply when weeds are small and actively growing. Use the higher rate in the respective rate range for larger weeds and when perennials are present. Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may be added to spray mixtures to increase the herbicidal effectiveness on certain weeds. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.		

Use Precautions, Restrictions and Limitations:

- **Important Notice:** Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of such injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

2/27

- Do not disturb treated soil through tillage between application and planting of soybeans.
- Do not use on sandy soils with less than 1.0% organic matter.
- In treated fields, plant soybean seed as deep as practical, but not less than 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- Do not make more than one application per season regardless of the application rate used.
- **Do not apply GF-1406 as a preplant application in soybeans unless you are prepared to accept the results of soybean injury, including possible stand loss and/or yield reduction.**
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with GF-1406.
- Do not apply more than 1 1/3 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

Forestry, Rangeland, Established Pasture, and Non-cropland Uses

Agricultural Use Requirements for Forest Use (Except Tree Injection Use): For use in forests, follow PPE and Reentry instructions in the "Agricultural Use Requirements" section under the "Directions for Use" heading of this label.

Agricultural Use Requirements for Rangeland, Pasture, Forest (Tree Injection Only) and Non-cropland Areas: When this product is applied to rangeland and established pastures not harvested for hay or seed; non-cropland areas, and when applied by tree injection in forest sites, follow reentry requirements given in the "Non-Agricultural Use Requirements" section under the "Directions for Use" heading of this label.

Forestry Uses

Forest site preparation, forest roadsides, brush control, established conifer release (including Christmas trees and reforestation areas)

Treatment Site Method of Application	GF-1406	Specific Use Directions
Annual Weeds	1 1/3 to 2 2/3 pt/acre	Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult to control perennial broadleaf weeds and woody species, use up to 2 2/3 qt of GF-1406 and 1 to 4 qt of Garlon® 3A herbicide per acre. For conifer release, make application in early spring before budbreak of conifers when weeds are small and actively growing.
Biennial and perennial broadleaf weeds and susceptible woody plants	2 2/3 to 5 1/3 pt/acre	
Spot Treatment to control broadleaf weeds	See Instructions for "Spot Treatment"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the recommended broadcast rate and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayers under "Application".
Conifer Release: Species such as white pine, ponderosa pine, jack pine, red pine, black spruce, white spruce, red spruce,	1 to 2 qt/acre	To control competing hardwood species such as alder, aspen, birch, hazel, and willow, apply from mid to late summer when growth of conifer trees has hardened off and woody plants are still actively growing. Apply with ground or air equipment, using sufficient spray

and balsam fir		volume to ensure complete coverage. Because this treatment may cause occasional conifer injury, do not apply if such injury cannot be tolerated.
Directed Spray: Conifer plantations including pine	2 2/3 qt/100 gal	Apply when brush or weeds are actively growing by directing the spray so as to avoid contact with conifer foliage and injurious amounts of spray. Apply in oil, oil-water, or water carrier in a spray volume of 10 to 100 gallons per acre.
Basal Spray (May also be used in rangeland, pastures, and noncropland)	5 1/3 qt/100 gal or	Thoroughly wet the base and root collar of all stems until the spray begins to accumulate around the root collar at the ground line. Wetting stems with the mixture may also aid in control.
Surface of Cut Stumps (May also be used in rangeland, pastures, and noncropland)	1.75 fl oz/gal of water	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and Girdle (May also be used in rangeland, pastures, and noncropland)		Cut frills (overlapping V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Saturate the freshly cut frills with the 2,4-D mixture.
Tree Injection Application (May also be used in rangeland, pastures, and noncropland)	(1 to 2 ml per injection site)	To control and prevent resprouting of unwanted hardwood trees such as elm, hickory, oak, and sweetgum in forests and other non-crop areas, apply by injecting at a rate of 1 ml of undiluted GF-1406 per inch of trunk diameter as measured at breast height (DBH), approximately 4 1/2 ft above the ground. Injection sites, however, should be as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Maples should not be treated during the spring sap flow. For hard to control species such as ash, maple, and dogwood use 2 ml of undiluted GF-1406 per injection site or double the number of 1 ml injections. Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Precautions and Restrictions:

- Do not allow sprays to contact conifer shoot growth (current year's new growth) or injury may occur.
- Do not apply to nursery seedbeds.
- For conifer release, do not use on plantations where pine or larch are among the desired species.
- **Grazing and Haying Restrictions:** If grazing or haying is anticipated, do not apply more than 2 2/3 pt/acre of GF-1406 per application. Do not harvest forage or hay from treated areas for 7 days after application.
- For broadcast applications, do not apply more than 5 1/3 pt/acre of GF-1406 per 12-month period.

23/27

Rangeland, Established Grass Pastures (Including Perennial Grasslands Not In Agricultural Production Such As Conservation Reserve Program Acres)

Target Weeds or Woody Plants	GF-1406 (pt/acre)	Specific Use Directions
Annual broadleaf weeds Biennial and perennial broadleaf weeds	1 1/3 1 1/3 to 2 2/3	For best results, apply when weeds are small and growing actively before the bud stage. Apply when musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks appear. Refer to the "Weeds Controlled" section for a listing of susceptible weed species and weeds that may be only partially controlled and require repeat applications and/or use of higher recommended rates, even under ideal conditions of application
Spot Treatment to control broadleaf weeds	See Instructions for "Spot Treatment"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the broadcast rate recommended for this treatment site and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayers under "Application".
Tree Injection Application		See instructions for tree injection application in "Forestry Uses" section.
Wild garlic and wild onion	2 2/3	Make three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring.
Broadleaf weed control in newly sprigged coastal bermudagrass	1 1/3 to 2 2/3	Applications may be made either preemergence or postemergence. Follow "Specific Use Directions" for annual, biennial and perennial broadleaf weed control, above.
Sand shinnery oak Sand sagebrush	1 1/3	Sand shinnery oak: Apply by aircraft between May 15 and June 15. Sand sagebrush: Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use a 1:4 oil-water emulsion as carrier and a spray volume of 3 to 5 gallons per acre.
Big sagebrush Rabbitbrush	4	Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use a 1:4 oil-water emulsion as carrier and a spray volume of 3 to 5 gallons per acre. Retreatment may be needed.
Chamise, manzanita, buckbrush, coastal sage, coyotebrush, and chaparral species.	2 2/3	Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Use water or 1:4 oil-water emulsion as carrier and a spray volume of 5 to 10 gallons per acre. Retreatment may be needed.
Southern wild rose Broadcast application Spot treatment	up to 2 2/3 2/3 gal/100 gal of spray	Broadcast: Apply in a spray volume of 5 or more gallons per acre by aircraft or 10 or more gallons per acre by ground equipment. Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Use 2/3 gallon of GF-1406 plus 4 to 8 fluid ounces of

	an agricultural surfactant per 100 gallons of water. Two or more treatments may be required. Do not exceed 2 2/3 pt per acre per application.
CRP Acres	For program lands such as CRP, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Precautions and Restrictions:

- Do not use on bentgrass, alfalfa, clover, or other legumes.
- Do not use on newly seeded areas until grass is well established.
- Do not use from early boot to milk stage where grass seed production is desired.
- Do not apply within 30 days of a previous application.
- **Grazing and Haying Restrictions:** In grazed areas, do not apply more than 2 2/3 pt/acre of GF-1406 per application. Do not harvest forage or hay from treated areas for 7 days after application.
- Do not apply more than 5 1/3 pt/acre of GF-1406 per use season.

Non-cropland Areas

Such as fencerows, hedgerows, roadsides, rights-of way, utility power lines, railroads, airports, and other non-crop areas

Treatment Site Method of Application	GF-1406 (pt/acre)	Specific Use Directions
Annual broadleaf weeds Biennial and perennial broadleaf weeds and susceptible woody plants	1 1/3 to 2 2/3 2 2/3 to 5 1/3	Apply when annual weeds are small and growing actively before the bud stage. Biennial and perennial weeds should be rosette to bud stage, but not flowering at the time of application. For difficult to control perennial broadleaf weeds and woody species, tank mix up to 2 2/3 qt of GF-1406 plus 1 to 4 qt of Garlon 3A herbicide per acre. Oil or wetting agent may be added to the spray, if needed for increased effectiveness. For ground application: (High volume) apply a total spray volume of 100 to 400 gallons per acre; (low volume) apply a total spray volume of 10 to 100 gallons per acre. For helicopter: Apply a total spray volume of 5 to 30 gallons per acre.
Spot Treatment to control broadleaf weeds	See Instructions for "Spot Treatment"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the broadcast rate recommended for this treatment site and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayers under "Application".
Tree Injection Application		See instructions for tree injection application in "Forestry Uses" section.
Southern wild rose Broadcast application Spot treatment	up to 2 2/3 2/3 gal/100 gal of spray	Broadcast: Apply in a spray volume of 5 or more gallons per acre by aircraft or 10 or more gallons per acre by ground equipment. Apply when foliage is well developed. Thorough coverage is required. Use 2 2/3 qt of GF-1406 plus 4 to 8 fluid ounces of an agricultural surfactant per 100 gallons of water. Two or more treatments

25/27

		may be required.
--	--	------------------

Precautions and Restrictions:

- Do not apply to newly seeded areas until grass is well established.
- Bentgrass, St. Augustine, clover, legumes and dichondra may be severely injured or killed by this treatment.
- When multiple applications of up to 2.0 lbs a.e. per acre are utilized to reach the maximum seasonal use rate, do not reapply to a treated area within 30 days of a previous application.
- **Grazing and Haying Restrictions:** If grazing or haying is anticipated, do not apply more than 2 2/3 pt/acre of GF-1406 per application. Do not harvest forage or hay from treated areas for 7 days after application.
- Do not apply more than 5 2/3 pt/acre of GF-1406 per use season.
- Use 2 or more gallons of spray solution per acre.

Turf Uses

Grasses Grown for Seed or Sod Farms

Agricultural Use Requirements: When used in grass grown for seed or sod farms, follow PPE and reentry instructions in the "Agricultural Use Requirements" section of this label.

Treatment Site (Application Timing)	GF-1406 (pt/acre)	Specific Use Directions
Grasses Grown for Seed (Postemergence Use) Seedling grass (five-leaf stage or later) Well-established grasses	1/2 to 2/3 2/3 to 2 2/3	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 2/3 pt/acre. Cool season grasses are tolerant of higher rates. Do not apply to grass in the early boot through milk stage if seed production is desired. When grass is well established, higher rates of up to 2 2/3 pints/acre may be applied for control of hard-to-kill annual or perennial weeds.
Sod Farms (Postemergence)	1 1/3 to 2 2/3	Deep-rooted perennials such as bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.

26/27

Precautions and Restrictions:

- Do not use on creeping grasses such as bentgrass except as a spot treatment.
- Do not use on injury-sensitive southern grasses such as St. Augustinegrass.
- Do not use on dichondra or other herbaceous ground covers. Legumes may be damaged or killed.
- **Preharvest Interval:** Do not cut forage for hay within 7 days of application.
- Do not apply within 21 days of a previous application.
- **Reseeding:** Delay reseeding at least 30 days following application. Preferably, with spring application, reseed in the fall and with fall application, reseed in the spring.
- **Grazing and Haying Restrictions:** If grazing or haying is anticipated, do not apply more than 2 2/3 pt/acre of GF-1406 per application. Do not harvest grass for hay from treated areas for 7 days after application.
- Do not apply more than 5 1/3 pt/acre of GF-1406 per use season.
- Use sufficient spray solution for thorough and uniform coverage, and no less than 2 gallons per acre.

Ornamental Turf (Excluding Grasses Grown For Seed or Sod Farms)

(Includes lawns, golf courses, cemeteries and parks, airfields, roadsides, and vacant lots)

Use Requirements for Ornamental Turf Areas: When this product is applied to ornamental turf areas, follow PPE and reentry instructions in the "Non-agricultural Use Requirements" section of this label.

Treatment Site (Application Timing)	GF-1406 (pt/acre)	Specific Use Directions
Ornamental Turf (Postemergence)		Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth.
Seedling grass (five-leaf stage or later)	1/2 to 2/3	Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeat applications.
Well-established grasses	1 1/3 to 2	Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 2/3 pt/acre. Cool season grasses are tolerant of higher rates.
Biennial and perennial broadleaf weeds	2	

Precautions, Restrictions:

- Do not use on creeping grasses such as bentgrass except as a spot treatment.
- Do not use on injury-sensitive southern grasses such as St. Augustinegrass.
- Do not use on dichondra or other herbaceous ground covers. Legumes may be damaged or killed.
- Do not reapply within 21 days of a previous application.
- **Reseeding:** Delay reseeding at least 30 days following application. Preferably, with spring application, reseed in the fall and with fall application, reseed in the spring.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- **Turf Reentry:** Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply more than 2 broadcast applications per year per treatment site (excluding spot treatments).
- Do not apply more than 3.0 lb. a.e. per year (excluding spot treatments).
- Do not apply more than 2 pt/acre of GF-1406 per application.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

27/27

Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC
EPA-Accepted: __/__/__