

62719-558

02-24-2009

1/18



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

FEB 24 2009

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Kimberly S. Gilbert  
Regulatory Leader  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: Notification(s) for Label Revisions under PRN 2007-4 and 98-10  
Added "Produced for" before registrant name and address

Dear Ms. Gilbert:

The Agency is in receipt of your Application(s) for Pesticide Notification under  
Pesticide-Registration-Notices (PRN)-98-10 and 2007-4 dated January 6, 2009 for:

**EPA Registration 62719-558**

**EF-1383 (A.I. Florasulam, 2,4-D)**

**EPA Registration 62719-563**

**GF-184 (A.I. Florasulam, Fluroxypyr)**

The Registration Division (RD) has conducted a review of the request(s) for applicability under PRN 98-10 and PRN 2007-4 and finds that the label changes requested fall within the scope of PRN-98-10 and PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identify the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

A handwritten signature in black ink that reads "Rachel C. Holloman".

Linda Arrington, *for*  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

MacF 117110



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number Dow AgroSciences LLC/62719-563	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/GF-184	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ <b>NOTIFICATION</b> _____ Product Name <b>FEB 24 2009</b>	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and PR Notice 2007-4 and EPA regulations at 40 CFR 152.46, 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and PR Notice 2007-4 and 40 CFR 152.46, 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt. 23.8 lb	No. per container 2	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2 x 2.5 gal		5. Location of Label Directions <input checked="" type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Kimberly S. Gilbert		Title Regulatory Manager		Telephone No. (Include Area Code) (317) 337-4685 (fax: 317-337-4649)	
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)  
2. Signature 		3. Title Regulatory Manager			
4. Typed Name Kimberly S. Gilbert (ksgilbert@dow.com)		5. Date January 5, 2009			

DK

308/2E  
January 5, 2009



Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
U. S. Environmental Protection Agency  
One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

GF-184 (A.I. FLORASULAM, FLUROXYPYR)  
EPA REGISTRATION NUMBER: 62719-563  
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICES 98-10 AND 2007-4

Enclosed please find labeling for the notification action of GF-184 herbicide. The following changes have been made by notification:

1. Storage and Disposal: Revised section per PR Notice 2007-4 and Chapter 13 of the Label Review Manual.
2. Added "Produced for" before registrant name and address.

This notification is consistent with the provisions of PR Notice 98-10 and PR Notice 2007-4 and EPA regulations at 40 CFR 152.46, 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and PR Notice 2007-4 and 40 CFR 152.46, 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Contents of Submission**


- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled GF-184 (S8D/GF-184/Notif/01-05-09) (15 Pages plus Registration Notes) (5 Copies)

If you require further information, please contact Cindy Loy, Regulatory Specialist at (317) 337-4655 or Paula McKinnies, Registration Assistant for this product, at (317) 337-4679.

Sincerely,

Kimberly S. Gilbert  
Regulatory Leader  
(317) 337-4685  
(317) 337-4649 (FAX)

Enclosures



# EF-1383


EPA Reg. No. 62719-558

## Registration Notes:

Source label text based on EPA-accepted text dated December 24, 2008.

### Changes by notification:

1. Storage and Disposal: Revised section per PR Notice 2007-4 and Chapter 13 of the Label Review Manual.
2. Added "Produced for" before registrant name and address.



5  
18

(Base label):

# EF-1383

## Herbicide

**For postemergent control of annual broadleaf weeds in wheat (including durum), barley, rye and triticale.**

<b>Group</b>	<b>2</b>	<b>4</b>	<b>HERBICIDES</b>
--------------	----------	----------	-------------------

Active Ingredient:

- florasulam: N-(2,6-difluorophenyl)-8-fluoro-5-methoxy (1,2,4)triazolo(1,5-c)pyrimidine-2-sulfonamide ..... 0.58%
- 2,4-dichlorophenoxyacetic Acid, 2-ethylhexyl ester ..... 42.33%
- Other Ingredients ..... 57.09%
- Total..... 100.00%

### NOTIFICATION

FEB 24 2009

Acid equivalent – 2,4-D – 28.06% - 2.5 lb/gal  
Contains 0.052 lb of florasulam per gallon

### Keep Out of Reach of Children

## CAUTION      PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Harmful If Swallowed Or Inhaled • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.**


#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber ≥14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See Engineering Controls for additional requirements.



Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

### Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### (Storage and Disposal for rigid containers 5 gal or less)

#### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Storage below 14°F will cause the product to freeze. If product freezes, bring to room temperature and agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for refillable rigid containers greater than 5 gal)**

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Storage below 14°F will cause the product to freeze. If product freezes, bring to room temperature and agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)**

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Storage below 14°F will cause the product to freeze. If product freezes, bring to room temperature and agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

8  
18

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.


Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-558

EPA Est. \_\_\_\_\_

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**Net Content**\_\_\_\_\_





2  
18

(Label booklet cover):

# EF-1383

## Herbicide

**For postemergent control of annual broadleaf weeds in wheat (including durum), barley, rye and triticale.**

<b>Group</b>	<b>2</b>	<b>4</b>	<b>HERBICIDES</b>
--------------	----------	----------	-------------------

**Active Ingredient:**

florasulam: N-(2,6-difluorophenyl)-8-fluoro-5-methoxy (1,2,4)triazolo(1,5-c)pyrimidine-2-sulfonamide .....	0.58%
2,4-dichlorophenoxyacetic Acid, 2-ethylhexyl ester .....	42.33%
Other Ingredients .....	57.09%
Total.....	100.00%

Acid equivalent – 2,4-D – 28.06% - 2.5 lb/gal  
Contains 0.052 lb of florasulam per gallon

### Keep Out of Reach of Children

## CAUTION      PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**


**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-558

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**



10  
14

(Page 1 through end):

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

## CAUTION

**Harmful If Swallowed Or Inhaled • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.**

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber  $\geq 14$  mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**All mixers, loaders, applicators, flaggers, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

#### User Safety Recommendations

Users should:


- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.



Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

### Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

2,4-dichlorophenoxyacetic acid has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.


**Pesticide Storage:** Store in original container only. Storage below 14°F will cause the product to freeze. If product freezes, bring to room temperature and agitate before use.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty



the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable containers 5 gallons or larger:**

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable containers 5 gallons or larger:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## General Information

Use EF-1383 herbicide as a postemergence herbicide for the control of annual broadleaf weeds such as chickweed, wild buckwheat, mustards and cleavers in wheat (including spring, winter and durum), barley, rye and triticale not underseeded with legumes.


EF-1383 rapidly stops growth of susceptible weeds. However, typical symptoms (discoloration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent upon weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

## Use Precautions and Restrictions

When applying this product in tank mix combination, follow all applicable use directions, precautions, and limitations on each manufacturer's label.

**Chemigation:** Do not apply this product through any type of irrigation system.

Do not apply EF-1383 directly to, or otherwise permit it to come into direct contact with, susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas,



potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, or tobacco. Do not permit spray mists containing EF-1383 to drift onto such plants.

For best resistance management stewardship, do not use more than once per season or use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Do not apply to crops underseeded with legumes.

**Crop Rotation Intervals**

The following rotational crops may be planted at the indicated interval following application of EF-1383.

Numbers in parentheses (-) refer to Specific Crop Rotation Information.

Crop	Rotation Interval (1) (Months)
barley, oats, wheat, grasses	0.5
field corn, popcorn, seed corn, sweet corn, sorghum	3
alfalfa, canola, chickpea, soybean, dry bean, field pea, flax, lentil, potato, safflower, sugar beet, sunflower	9
other crops not listed	12

**Specific Crop Rotation Information:**


1. Minimum number of months that must pass before planting other crops after application of EF-1383.

**Avoiding Injurious Spray Drift**

This product can affect broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Do not apply EF-1383 directly to, or allow spray drift to come into contact with, broadleaf crops including, but not limited to, alfalfa, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See Crop Rotation Intervals section.)

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure crops, whether dormant or actively growing. When applying EF-1383, use low pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets. Under adverse weather conditions, fine spray droplets that do not settle rapidly onto target vegetation may be carried a considerable distance from the treatment area. A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use recommendations and precautions on the product label.

**Ground Applications:** To minimize spray drift, apply EF-1383 in a total spray volume of 8 gallons or more per acre using spray equipment designed to produce large-droplet, low pressure sprays. Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Spot treatments should be applied only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.



14  
18

**Aerial Application:** To minimize spray drift, apply EF-1383 in a total spray volume of 3 gallons or more per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high potential for temperature inversion. Spray drift from aerial application can be minimized by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 3/4 the rotor or wing span of the aircraft. Spray pattern and droplet size distribution can be evaluated by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Do not apply under conditions of a low level air temperature inversion. A temperature inversion is characterized by little or no wind and lower air temperature near the ground than at higher levels. The behavior of smoke generated by an aircraft-mounted device or continuous smoke column released at or near site of application will indicate the direction and velocity of air movement. A temperature inversion is indicated by layering of smoke at some level above the ground and little or no lateral movement.

### Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

- The distance of the outer most operating nozzles on the boom must not exceed 75% of wingspan or 90% of rotor diameter.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory. [This information is advisory in nature and does not supersede mandatory label requirements.]

### Aerial Drift Reduction Advisory

**Information on Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

### Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

15  
18

**Boom Length:** For some use patterns, reducing the effective boom length to less than 75% of the wingspan or 90% of rotor length may further reduce drift without reducing swath width.

**Application Height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

## Mixing


### Mixing EF-1383 Alone

1. Fill the tank with 1/2 of the total amount of water.
2. Start agitation.
3. Add the required amount of EF-1383.
4. Continue agitation while filling the spray tank to the required volume.
5. To ensure a uniform spray mixture, continuous agitation is required during application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

### EF-1383 Applied in Tank Mix Combination

If a broader spectrum of weed control is needed, EF-1383 may be tank mixed with labeled rates of other herbicides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

### Tank Mixing Precautions:



16  
18

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates for respective products or maximum allowable application rates for any active ingredient in the tank mix.
- Do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment have been adequately cleaned. (See Equipment Clean-Out Procedures.)
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** A jar test is recommended prior to tank mixing to ensure compatibility of EF-1383 and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

**Mixing Order for Tank Mixes:**

1. Fill the spray tank to 3/4 of the total spray volume required with water.
2. Start agitation.
3. Add EF-1383 and agitate for 2 to 3 minutes
4. After adding EF-1383, add different formulation types in the following order: (1) dry flowables; (2) wettable powders; (3) aqueous suspensions, flowables and liquids. Maintain agitation and add: (4) emulsifiable concentrates; (5) solutions; and (6) adjuvants. Allow time for complete mixing and dispersion after each addition.
5. Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

**Clean-Out Procedures for Spray Equipment**

1. Drain any remaining spray mixture from the application equipment.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add household ammonia at a rate of 1 gallon per 100 gallons of water. Recirculate for 5 minutes and spray out part of this mixture for 5 minutes through the boom. Drain tank.
4. Remove all spray nozzles and screens and clean separately.
5. If spray equipment will be used for pesticide application to crops sensitive to EF-1383, steps 1-3 should be repeated. Exterior surfaces of spray equipment should also be thoroughly cleaned.


**Note:** Rinsate may be disposed of on site according to label use directions or at an approved waste disposal facility.

**Weeds Controlled or Suppressed by EF-1383**

**Weeds Controlled**

buckwheat, wild  
canola, volunteer<sup>1</sup>  
chamomile, corn  
chamomile, false (scentless)

mustard, black  
mustard, blue (purple)  
mustard, tansy  
mustard, treacle (bushy wallflower)





chamomile, mayweed (dogfennel)  
 chamomile, wild  
 chickweed, common  
 cleavers (catchweed bedstraw)  
 flixweed  
 knotweed  
 lambsquarters  
 London rocket  
 marshelder

mustard, tumble (Jim Hill)  
 mustard, wild  
 pennycress, field  
 pineappleweed  
 ragweed, common  
 radish, wild  
 shepherdspurse  
 smartweed (green, ladysthumb, Pennsylvania)  
 sunflower, annual

### Weeds Suppressed

dandelion (seedlings and overwintered  
 rosettes <6 inches in diameter  
 hempnettle<sup>3</sup>  
 pigweed, redroot<sup>3</sup>  
 prickly lettuce<sup>2</sup>

sowthistle, annual  
 sowthistle, perennial<sup>2</sup>  
 thistle, Russian<sup>3</sup>

<sup>1</sup>Including all herbicide-tolerant canola varieties.

<sup>2</sup>Up to the 4-leaf stage.

<sup>3</sup>Control can be achieved by mixing an additional 4 fl oz ae per acre of 2,4-D ester (8 fl oz per acre of a 4 lb per gallon 2,4-D ester).

## Application Directions

### Application Timing

Apply EF-1383 early postemergence to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds that have emerged at the time of application will be controlled.** If foliage is wet at the time of application, control may be decreased. Applications of EF-1383 are rainfast within 4 hours after application.

### Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Do not broadcast apply in less than 3 gallons of total spray volume per acre. For best results and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As vegetative canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under Avoiding Injurious Spray Drift.

### Adjuvants

Generally, this product does not require the use of an adjuvant to achieve satisfactory weed control. An adjuvant may be added to optimize herbicidal activity when applications are made at lower use rates or lower carrier volumes; under conditions of cool temperature, low relative humidity or drought; or when a tank mix partner recommends the use of an adjuvant.

## Wheat (Including Durum), Barley, Rye, Triticale

Apply 11 fl oz of EF-1383 per acre to actively growing wheat (including spring, winter and durum), barley, rye, and triticale from the 4-leaf crop growth stage up to the joint stage (Zadoks scale 31). Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from application made to seedling weeds. A later application when the crop is between the jointing and boot stage of growth may be used to control later emerging weeds; however, do not apply unless the risk of injury is acceptable. Do not apply after the boot stage.

Warm, moist growing conditions promote active weed growth and enhance the activity of EF-1383 by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought

18  
18

stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

**Tank Mixtures for Wheat (Including Durum):** EF-1383 may be applied in tank mix combination with labeled rates of other products registered for postemergence application in wheat. See Tank Mixing Precautions under Mixing. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

**Restrictions:**

- Do not apply more than 11 fl oz of EF-1383 per acre per growing season.
- Livestock may be grazed on treated crops 7 days following application.
- **Preharvest Interval:** Do not apply with 60 days of crop harvest.

**NOTIFICATION**

FEB 24 2009

---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

**Warranty Disclaimer**

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies ~~cannot be~~ varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

EPA-accepted 12/24/08

