


62719-565

12/11/2006

1/19

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Reg. Number: 62719-565	Date of Issuance: DEC 11 2006
	Term of Issuance: conditional	
<p>NOTICE OF PESTICIDE:</p> <p><input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	Name of Pesticide Product: Turf Fertilizer – Contains Gallery plus Team Pro	
Name and Address of Registrant (include ZIP Code): Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Submit the data listed below: <ol style="list-style-type: none"> a. Groups A and B product chemistry data on this formulation. Submit these data within eight (8) months of the date of this notice. 3. Make the following label changes: (See Page 2 of this notice) 		
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	<p>Date:</p> <p style="text-align: center;">DEC 11 2006</p>	

3a. Revise the **Hazards to Humans and Domestic Animals** section for both sub labels to read:

“Harmful if absorbed through skin, inhaled, or swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A). Remove and wash contaminated clothing before reuse. Avoid breathing dust. Wear protective eyewear. [Protective eyewear, which may include goggles, face shield and/or safety glasses, may be specified, if appropriate.]

b. Under **First Aid** (for both sub labels), add the “If on skin” and “If in eyes” treatments from PR Notice 2001-1. In addition, add a “Note to Physician” section.

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

319

[Sub Label A: Commercial Uses]

Turf Fertilizer - Contains Gallery[®] plus Team[™] Pro

Specialty Herbicide

A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.

For professional use only.

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.38%
trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.50%
benfen: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.50%
Inert Ingredients	98.62%
Total	100.00%

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

DEC 11 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-585

[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]

- Contains 0.276 lb active ingredient per 20 lb bag.
- Contains 0.345 lb active ingredient per 25 lb bag.
- Contains 0.552 lb active ingredient per 40 lb bag.
- Contains 0.69 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O).....	X.XX%

Keep Out of Reach of Children

CAUTION

See back or side panel of container for additional precautionary statements and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-XXX

EPA Est. _____

4/19

®™ Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, Indiana 46268 U.S.A.

Net Weight _____

5/19

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

6/19

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This fertilizer product contains Gallery[®] and Team[™] Pro, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliar applied postemergence herbicide.

Optimum weed control performance will be obtained when an application is followed by overhead irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Use Precautions and Restrictions:

- Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turf sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 175 lb per acre, or 12 to 16 weeks after a single application of 265 lb per acre.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. **Herbicides in this product prevent seed emergence and may injure annual bedding plants.**
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.

Treatment of Turf or Ornamental Species Not Listed on the Label for Turf Fertilizer – Contains Gallery Plus Team Pro

7/19

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury such as root stunting during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

This product is recommended for use on the following established perennial turfgrass species:

Established Cool Season Grasses

Common Name	Scientific Name
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Grasses

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum var. Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Austinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name	Scientific Name
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus spp.</i>
apricot	<i>Prunus armeniaca</i>
birch, paper	<i>Betula papyrifera</i>
cherry	<i>Prunus spp.</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
honeylocust	<i>Gleditsia triacanthos</i>
maple, flowering	<i>Abutilon hybridum</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
oak, pin	<i>Quercus palustris</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, scotch	<i>Pinus sylvestris</i>

plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>

Ornamental Shrubs

Common Name	Scientific Name
arborvitae	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cinquefoil	<i>Potentilla fruticosa</i>
cleyera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
elaeanthus, silverberry	<i>Elaeagnus pungens</i>
euonymus, winged	<i>Euonymus alata</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
forsythia	<i>Forsythia</i> spp.
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
juniper, creeping	<i>Juniperus horizontalis</i>
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
myrtle, wax	<i>Myrica cerifera</i>
privet	<i>Ligustrum</i> spp.
pyracantha	<i>Pyracantha fortuneana</i>
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spiraea, vanhouttei	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

Ornamental Ground Cover Plantings

Common Name	Scientific Name
ceanothus	<i>Ceanothus</i> spp.
coyote brush	<i>Baccharis pilularis</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
gazania	<i>Gazania</i> spp.
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lilyturf, bigblue	<i>Liriope muscari</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rosemary	<i>Rosmarinus officinalis</i>
snow-in-summer	<i>Cerastium tomentosum</i>
St. Johnswort	<i>Hypericum coris</i>

Application Directions

Application Equipment and Application Techniques

Apply this product uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granular fertilizer. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turf injury may occur. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	175	XXX	XXX	XXX	265
XXXXXXX	XX	XX	XX	XX	XX

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus</i> spp.
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
(large crabgrass)	
(smooth crabgrass)	
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria</i> spp.
(bottlegrass)	
(bristlegrass)	
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust foxtail)	
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>

10/19

woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds suppressed or partially controlled by a single application or controlled by repeat applications when applied prior to weed germination:

Common Name	Scientific Name
clover spp. ¹	<i>Trifolium</i> spp.
clover, white ¹	<i>Trifolium repens</i>
dandelion ¹	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
plantain, slender	<i>Plantago elongata</i>

¹From seed only

Uses

Established Turfgrass

Application Timing: Apply in late summer or early fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft). A minimum of 175 lb per acre (4 lb per 1000 sq ft) is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Established Ornamental Plants in Landscape Settings

This product is recommended as a preemergence application for control of certain broadleaf weeds and annual grasses in a wide variety of established ornamental trees, shrubs, and ground covers. Apply only to non-bearing fruit and nut trees and non-bearing vineyards which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing: Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft) as a spring application or 265 lb per acre (6 lb per 1000 sq ft) as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

11/19

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC
EPA-accepted ___/___/___

12/19

[Sub Label B: Residential Uses]

Turf Fertilizer - Contains Gallery[®] plus Team[™] Pro

Specialty Herbicide

A fertilizer containing preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals.

For residential use only.

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.38%
trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.50%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α - trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.50%
Inert Ingredients	98.62%
Total	100.00%

[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	X.XX%
Available Phosphoric Acid (P ₂ O ₅)	X.XX%
Soluble Potash (K ₂ O)	X.XX%

Keep Out of Reach of Children

CAUTION

See back or side panel of container for additional precautionary statements and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-280

EPA Est. _____

13/19

®™ Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, Indiana 46268 U.S.A.

Net Weight _____

14/19

Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Harmful if Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust. Wear long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes, and protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

15/19

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your solid waste agency or 1-800-992-5994 for disposal instructions.

Never place unused product down any indoor or outdoor drain.

General Information

This fertilizer product contains Gallery® and Team™ Pro, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliar applied postemergence herbicide.

Optimum weed control performance will be obtained when an application is followed by irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Use Precautions and Restrictions:

- Apply to newly seeded turf (including overseeded turf) only after seedlings are established and well rooted. Do not overseed established turf sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 4 lb per 1000 sq ft, or 12 to 16 weeks after a single application of 6 lb per 1000 sq ft.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. **Herbicides in this product prevent seed emergence and may injure annual bedding plants.**
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.

Treatment of Turf or Ornamental Species Not Listed on the Label for Turf Fertilizer – Contains Gallery Plus Team Pro

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury such as root stunting during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

This product is recommended for use on the following established perennial turfgrass species:

Established Cool Season Turfgrasses

bentgrass, colonial
bentgrass, creeping
bluegrass, Kentucky
fescue, chewing
fescue, creeping red

Established Warm Season Turfgrasses

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass

16/19

fescue, sheep	zoysiagrass
fescue, tall	
ryegrass, perennial	

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

almond,	peach
apple, drabapple	pine, Austrian
apricot	pine, eastern white
birch, paper	pine, Japanese black
cherry	pine, scotch
cottonwood	plum
dogwood, flowering	redbud, eastern
dogwood, kousa	spruce, Colorado
honeylocust	spruce, Norway
maple, flowering	sweetgum
maple, red	sycamore
maple, silver	walnut, black
oak, pin	

Ornamental Shrubs

arborvitae	honeysuckle
azalea	juniper, creeping
boxwood, littleleaf	laurel, mountain
camellia, Japanese	lilac, common
camellia, sasanqua	myrtle, wax
cinquefoil	privet
cleyera, Japanese	pyracantha
cotoneaster, cranberry	redcedar, eastern
elaeanthus, silverberry	rhododendron
euonymus, winged	spiraea, vanhoutte
euonymus,	viburnum
wintercreeper	
forsythia	yew, Japanese
holly	yewpine

Ornamental Ground Cover Plantings

ceanothus	ivy, English
coyote brush	lilyturf, bigblue
daisy trailing African	plumbago, dwarf
gazania	rosemary
ice plant, largeleaf	snow-in-summer
ivy, Algerian	St. Johnswort

Application Directions

Application Equipment and Application Techniques

Apply this product uniformly over the lawn using a properly calibrated drop or rotary-type spreader designed to apply granular fertilizer. Before each application, calibrate the spreader according to the manufacturer's directions. Check frequently to be sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. Avoid

17/19

skips and excessive overlaps as poor weed control or turf injury may occur. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	4	XX	XX	XX	6
XXXXXXX	XX	XX	XX	XX	XX

Weeds Controlled

Weeds controlled when applied prior to weed germination:

- | | |
|---------------------|----------------------|
| annual bluegrass | geranium, Carolina |
| barnyardgrass | goosegrass |
| burweed, lawn | henbit |
| buttercup | knotweed, prostrate |
| carpetweed | mallow, little |
| celery, wild | parsley - pierl |
| chickweed, common | pennywort, lawn |
| chickweed, mouseear | pineappleweed |
| crabgrass | speedwell, thymeleaf |
| (large crabgrass) | spurge, prostrate |
| (smooth crabgrass) | spurge, spotted |
| foolsparsley | spurry, corn |
| foxtails | woodsorrel, creeping |
| (bottlegrass) | woodsorrel, yellow |
| (bristlegrass) | |
| (giant foxtail) | |
| (green foxtail) | |
| (pigeongrass) | |
| (robust foxtail) | |

Weeds suppressed or partially controlled by a single application or controlled by repeat applications when applied prior to weed germination:

- clover spp.¹
- clover, white¹
- dandelion¹
- groundsel, common
- medic, black
- plantain, broadleaf
- plantain, buckhorn
- plantain, slender

¹From seed only

Uses

Established Home Lawns

Application Timing: Apply in late summer or early fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

Broadcast Application Rates

18/19

Apply this product at a rate of 4 to 6 lb per 1000 sq ft. A minimum of 4 lb per 1000 sq ft is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Established Ornamental Plants in Landscape Settings

This product is recommended as a preemergence application for control of certain broadleaf weeds and annual grasses in a wide variety of established ornamental trees, shrubs, and ground covers. Apply only to non-bearing fruit and nut trees and non-bearing vineyards which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing: Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds. Do not apply this product to plantings when the foliage is wet.

Broadcast Application Rates

Apply this product at a rate of 4 to 6 lb per 1000 sq ft as a spring application or 6 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

19/19

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC
EPA-accepted ___/___/___