

62719-565

4.30.2008

1/41



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

APR 30 2008

Ms. Kimberly Gilbert  
Regulatory Leader  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

RE: Notification of Minor Label Change per PR Notice 98-10  
EPA Registration Number: 62719-565  
Date of Submission: April 9, 2008

Dear Ms. Gilbert:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated April 9, 2008, for the product, Turf Fert-Contains Gallery Plus Team. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



United States  
**Environmental Protection Agency**  
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number Dow AgroSciences LLC/62719-565	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/Turf Fert-Contains Gallerv plus Team	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kimberly S. Gilbert	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4685 (fax: 317-337-4649)
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Kimberly S. Gilbert</i>	3. Title Regulatory Manager	
4. Typed Name Kimberly S. Gilbert (ksgilbert@dow.com)	5. Date April 9, 2008	

13/14

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro

EPA Reg. No. 62719-565

The master label for Turf Fertilizer – Contains Gallery plus Team Pro consists of separate, self-contained sub labels for professional use and residential use, respectively.


## Registration Notes:

Source label text based on EPA accepted copy dated December 11, 2006 and notification coded “Q8B / Turf Fertilizer – Contains Gallery Plus Team Pro / MSTR Notif / 09-20-07.”

### Changes by notification:

1. Updated trademark symbol throughout.
2. Added to list of optional label claims.

®Trademark of Dow AgroSciences LLC



4/41

**[Sub Label A: Commercial Use]**

(Base label for rigid containers 5 gal or less):

# **Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro**

## **Specialty Herbicide**

**A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.**

**For professional use only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] [including bentgrass]**

and/or

**For use on cool season [and warm season turfgrass]**

and/or


**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**



and/or  
**University tested [and researched]**  
 and/or  
**Labeled for use around a [wide] variety of trees [and shrubs]**  
 and/or  
**Never see crabgrass again**  
 and/or  
**Labeled for over 40 different grassy [and broadleaf] weeds**

Active Ingredients:  
 isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide  
 and isomers ..... 0.38%  
 trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine ..... 0.50%  
 benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine ..... 0.50%  
 Other Ingredients ..... 98.62%  
 Total ..... 100.00%

**[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]**

- Contains 0.276 lb active ingredient per 20 lb bag.
- Contains 0.345 lb active ingredient per 25 lb bag.
- Contains 0.552 lb active ingredient per 40 lb bag.
- Contains 0.69 lb active ingredient per 50 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O) .....	X.XX%

**Keep Out of Reach of Children**

## **CAUTION**

---

### **Precautionary Statements**

---

#### **Hazards to Humans and Domestic Animals**

**Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.**

#### **Personal Protective Equipment:**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves

*[Faint, illegible text or bleed-through from the reverse side of the page]*

6/41

- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

7/41

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-565

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC


**Produced for**

**Dow AgroSciences LLC**

**9330 Zionsville Road**

**Indianapolis, Indiana 46268**

**Net Weight** \_\_\_\_\_



(Base label for nonrigid containers any size):

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro

## Specialty Herbicide

**A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.**

**For professional use only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] [including bentgrass]**

and/or

**For use on cool season [and warm season turfgrass]**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or



Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers ..... 0.38%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine ..... 0.50%

benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine ..... 0.50%

Other Ingredients ..... 98.62%

Total..... 100.00%

[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O) .....	X.XX%

Keep Out of Reach of Children

# CAUTION

## Precautionary Statements

### Hazards to Humans and Domestic Animals


Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear



Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-565

EPA Est. \_\_\_\_\_



(Base label for refillable rigid containers greater than 5 gal):

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro

## Specialty Herbicide

A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.

For professional use only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control[/prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]


and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or



13/41

**Labeled for use around a [wide] variety of trees [and shrubs]**

and/or

**Never see crabgrass again**

and/or

**Labeled for over 40 different grassy [and broadleaf] weeds**

**Active Ingredients:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers..... 0.38%

trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine ..... 0.50%

benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro--p-toluidine ..... 0.50%

Other Ingredients ..... 98.62%

Total..... 100.00%

**[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]**

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O) .....	X.XX%

**Keep Out of Reach of Children**

# CAUTION

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals


**Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.**

### Personal Protective Equipment:

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear



14/41

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**



16  
41

(Base label for nonrefillable rigid containers larger than 5 gal):

# Turf Fertilizer - Contains Gallery® plus Team® Pro

## Specialty Herbicide

**A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.**

**For professional use only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] [including bentgrass]**

and/or

**For use on cool season [and warm season turfgrass]**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**


and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or





17/41

Labeled for use around a [wide] variety of trees [and shrubs]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different grassy [and broadleaf] weeds

Active Ingredients:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers..... 0.38%

trifluralin: α,α,α-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine ..... 0.50%

benefin: N-butyl-N-ethyl-α,α,α-trifluoro-2,6-dinitro--p-toluidine ..... 0.50%

Other Ingredients ..... 98.62%

Total ..... 100.00%

[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis

Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O).....	X.XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals


Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear



Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

19/  
41

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.


EPA Reg. No. 62719-565

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

**Produced for**  
**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, Indiana 46268**

**Net Weight** \_\_\_\_\_



(Label booklet cover):

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro

## Specialty Herbicide

A fertilizer containing Gallery and Team Pro, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established perennial turfgrasses and landscape ornamentals.

For professional use only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control[/prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] [including bentgrass]

and/or

For use on cool season [and warm season turfgrass]

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]


and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or



21/41

**Labeled for use around a [wide] variety of trees [and shrubs]**

and/or

**Never see crabgrass again**

and/or

**Labeled for over 40 different grassy [and broadleaf] weeds**

**Active Ingredients:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.38%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine .....	0.50%
benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro--p-toluidine .....	0.50%
Other Ingredients .....	98.62%
Total .....	100.00%

**[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]**

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

**Guaranteed Fertilizer Analysis**

Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O) .....	X.XX%

**Keep Out of Reach of Children**

**CAUTION**

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-565

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC


**Produced for**

**Dow AgroSciences LLC**

**9330 Zionsville Road**

**Indianapolis, Indiana 46268**

**Net Weight** \_\_\_\_\_



29/41

(Page 1 through end):

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## CAUTION

**Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.**

### Personal Protective Equipment:

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose

canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

---

### Directions for Use

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.**

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable rigid containers 5 gallons or less:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### Nonrefillable nonrigid containers:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

#### Refillable rigid containers larger than 5 gal:

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable rigid containers larger than 5 gal:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**General Information**

This fertilizer product contains Gallery® and Team® Pro, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliar applied postemergence herbicide.

Optimum weed control performance will be obtained when an application is followed by overhead irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

**Use Precautions and Restrictions:**

- Apply to newly seeded turfgrass (including overseeded turfgrass) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turfgrass sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 175 lb per acre, or 12 to 16 weeks after a single application of 265 lb per acre.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. **Herbicides in this product prevent seed emergence and may injure annual bedding plants.**
- This product may injure turfgrass that is weakened due to severe winter weather or is not well established.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turfgrass is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turfgrass injury may occur.

**Treatment of Turfgrass or Ornamental Species Not Listed on the Label for Turf Fertilizer – Contains Gallery Plus Team Pro**

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury such as root stunting during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**



25  
41

**This product is recommended for use on the following established perennial turfgrass species:**

**Established Cool Season Turfgrass**

<b>Common Name</b>	<b>Scientific Name</b>
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>


**Established Warm Season Turfgrass**

<b>Common Name</b>	<b>Scientific Name</b>
bahiagrass	<i>Paspalum notatum var. Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

**Ornamental Trees**

<b>Common Name</b>	<b>Scientific Name</b>
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus spp.</i>
apricot	<i>Prunus armeniaca</i>
birch, paper	<i>Betula papyrifera</i>
cherry	<i>Prunus spp.</i>
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
honeylocust	<i>Gleditsia triacanthos</i>
maple, flowering	<i>Abutilon hybridum</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
oak, pin	<i>Quercus palustris</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, scotch	<i>Pinus sylvestris</i>
plum	<i>Prunus spp.</i>
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>



**Ornamental Shrubs**

<b>Common Name</b>	<b>Scientific Name</b>
arborvitae	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cinquefoil	<i>Potentilla fruticosa</i>
cleyera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
elaeanthus, silverberry	<i>Elaeagnus pungens</i>
euonymus, winged	<i>Euonymus alata</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
forsythia	<i>Forsythia</i> spp.
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
juniper, creeping	<i>Juniperus horizontalis</i>
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
myrtle, wax	<i>Myrica cerifera</i>
privet	<i>Ligustrum</i> spp.
pyracantha	<i>Pyracantha fortuneana</i>
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spiraea, vanhouttei	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>

**Ornamental Ground Cover Plantings**

<b>Common Name</b>	<b>Scientific Name</b>
ceanothus	<i>Ceanothus</i> spp.
coyote brush	<i>Baccharis pilularis</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
gazania	<i>Gazania</i> spp.
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lilyturf, bigblue	<i>Liriope muscari</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rosemary	<i>Rosmarinus officinalis</i>
snow-in-summer	<i>Cerastium tomentosum</i>
St. Johnswort	<i>Hypericum coris</i>

**Application Directions****Application Equipment and Application Techniques**

Apply this product uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granular fertilizer. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**Spreader Settings as a Guide for Calibration**

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	175	XXX	XXX	XXX	265
XXXXXXXX	XX	XX	XX	XX	XX


**Weeds Controlled**

**Weeds controlled when applied prior to weed germination:**

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus</i> spp.
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
(large crabgrass)	
(smooth crabgrass)	
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria</i> spp.
(bottlegrass)	
(bristlegrass)	
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust foxtail)	
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

**Weeds suppressed or partially controlled by a single application or controlled by repeat applications when applied prior to weed germination:**

Common Name	Scientific Name
clover spp. <sup>1</sup>	<i>Trifolium</i> spp.
clover, white <sup>1</sup>	<i>Trifolium repens</i>
dandelion <sup>1</sup>	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>



plantain, slender

*Plantago elongata*

1From seed only

**Uses****Established Turfgrass**

**Application Timing:** Apply in late summer or early fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

**Broadcast Application Rates**

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft). A minimum of 175 lb per acre (4 lb per 1000 sq ft) is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

**Established Ornamental Plants in Landscape Settings**

This product is recommended as a preemergence application for control of certain broadleaf weeds and annual grasses in a wide variety of established ornamental trees, shrubs, and ground covers. Apply only to non-bearing fruit and nut trees and non-bearing vineyards which are defined as plants that will not bear fruit for at least 1 year after treatment.

**Application Timing:** Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

**Broadcast Application Rates**

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft) as a spring application or 265 lb per acre (6 lb per 1000 sq ft) as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

**Warranty Disclaimer**

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

29/41

---

### Limitation of Remedies

---


To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™Trademark of Dow AgroSciences LLC  
EPA accepted 12/11/06



30/41

**[Sub Label B: Residential Use]**

(Base label):

**Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup> Pro**

**Specialty Herbicide**

**A fertilizer containing preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals.**

**For residential use only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] [including bentgrass]**

and/or

**For use on cool season [and warm season turfgrass]**

and/or


**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**



31/41

and/or  
**University tested [and researched]**  
 and/or  
**Labeled for use around a [wide] variety of trees [and shrubs]**  
 and/or  
**Never see crabgrass again**  
 and/or  
**Labeled for over 40 different grassy [and broadleaf] weeds**

Active Ingredients:  
 isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide  
 and isomers ..... 0.38%  
 trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N*,  
*N*-dipropyl-*p*-toluidine ..... 0.50%  
 benefin: *N*-butyl-*N*-ethyl- $\alpha,\alpha,\alpha$ -  
 trifluoro-2,6-dinitro-*p*-toluidine ..... 0.50%  
 Other Ingredients ..... 98.62%  
 Total ..... 100.00%

[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]

Contains 0.276 lb active ingredient per 20 lb bag.  
 Contains 0.345 lb active ingredient per 25 lb bag.  
 Contains 0.552 lb active ingredient per 40 lb bag.  
 Contains 0.69 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N).....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	X.XX%
Soluble Potash (K <sub>2</sub> O).....	X.XX%

Keep Out of Reach of Children

# CAUTION

## Precautionary Statements

### Hazards to Humans and Domestic Animals

Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.

## User Safety Recommendations

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3/41

**First Aid**

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-565

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight \_\_\_\_\_



(Label booklet cover):

# Turf Fertilizer - Contains Gallery® plus Team® Pro

## Specialty Herbicide

**A fertilizer containing preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals.**

**For residential use only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] [including bentgrass]**

and/or

**For use on cool season [and warm season turfgrass]**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**


and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or



34/41

**Labeled for use around a [wide] variety of trees [and shrubs]**

and/or

**Never see crabgrass again**

and/or

**Labeled for over 40 different grassy [and broadleaf] weeds**

**Active Ingredients:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.38%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.50%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.50%
Other Ingredients .....	98.62%
Total.....	100.00%

**[Editor's Note: Supplemental distributor will choose one of the bag sizes listed below]**

Contains 0.276 lb active ingredient per 20 lb bag.

Contains 0.345 lb active ingredient per 25 lb bag.

Contains 0.552 lb active ingredient per 40 lb bag.

Contains 0.69 lb active ingredient per 50 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

<b>Guaranteed Fertilizer Analysis</b>	
Total Nitrogen (N) .....	X.XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	X.XX%
Soluble Potash (K <sub>2</sub> O) .....	X.XX%

**Keep Out of Reach of Children**

# CAUTION

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**


In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-565

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, Indiana 46268**



35  
41

(Page 1 through end):

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION**

**Harmful If Absorbed Through Skin, Swallowed Or Inhaled • Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, socks, shoes, chemical resistant gloves and protective eyewear. Remove and wash contaminated clothing before reuse.**

**User Safety Recommendations**

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

---

**Directions for Use**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

**Entry Restrictions:** Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering-in) until the watering-in is complete and the surface is dry.

36  
41

**Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.**

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

### General Information

This fertilizer product contains Gallery<sup>®</sup> and Team<sup>®</sup> Pro, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established home lawns and landscape ornamentals. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliar applied postemergence herbicide.

Optimum weed control performance will be obtained when an application is followed by irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

### Use Precautions and Restrictions:

- Apply to newly seeded turfgrass (including overseeded turfgrass) only after seedlings are established and well rooted. Do not overseed established turfgrass sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 4 lb per 1000 sq ft, or 12 to 16 weeks after a single application of 6 lb per 1000 sq ft.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. **Herbicides in this product prevent seed emergence and may injure annual bedding plants.**
- This product may injure turfgrass that is weakened due to severe winter weather or is not well established.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turfgrass is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turfgrass injury may occur.

### Treatment of Turfgrass or Ornamental Species Not Listed on the Label for Turf Fertilizer – Contains Gallery Plus Team Pro

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury such as root stunting during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

**This product is recommended for use on the following established perennial turfgrass species:**

37  
41

**Established Cool Season Turfgrass**

bentgrass, colonial  
bentgrass, creeping  
bluegrass, Kentucky  
fescue, chewings  
fescue, creeping red  
fescue, sheep  
fescue, tall  
ryegrass, perennial

**Established Warm Season Turfgrass**

bahiagrass  
Bermudagrass  
centipedegrass  
fescue, tall  
St. Augustinegrass  
zoysiagrass

**This product may be used for annual weed control in landscape settings on the following established ornamental plant species:**

**Ornamental Trees**


almond	peach
apple, crabapple	pine, Austrian
apricot	pine, eastern white
birch, paper	pine, Japanese black
cherry	pine, scotch
cottonwood	plum
dogwood, flowering	redbud, eastern
dogwood, kousa	spruce, Colorado
honeylocust	spruce, Norway
maple, flowering	sweetgum
maple, red	sycamore
maple, silver	walnut, black
oak, pin	

**Ornamental Shrubs**

arborvitae	honeysuckle
azalea	juniper, creeping
boxwood, littleleaf	laurel, mountain
camellia, Japanese	lilac, common
camellia, sasanqua	myrtle, wax
cinquefoil	privet
cleyera, Japanese	pyracantha
cotoneaster, cranberry	redcedar, eastern
elaeanthus, silverberry	rhododendron
euonymus, winged	spiraea, vanhoutte
euonymus, wintercreeper	viburnum
forsythia	yew, Japanese
holly	yewpine

**Ornamental Ground Cover Plantings**

ceanothus	ivy, English
coyote brush	lilyturf, bigblue
daisy trailing African	plumbago, dwarf
gazania	rosemary
ice plant, largeleaf	snow-in-summer
ivy, Algerian	St. Johnswort



**Application Directions**

**Application Equipment and Application Techniques**

3/9/11

Apply this product uniformly over the lawn using a properly calibrated drop or rotary-type spreader designed to apply granular fertilizer. Before each application, calibrate the spreader according to the manufacturer's directions. Check frequently to be sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

**Spreader Settings as a Guide for Calibration**

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	4	XX	XX	XX	6
XXXXXXXX	XX	XX	XX	XX	XX

**Weeds Controlled**

**Weeds controlled when applied prior to weed germination:**

- annual bluegrass
- barnyardgrass
- burweed, lawn
- buttercup
- carpetweed
- celery, wild
- chickweed, common
- chickweed, mouseear
- crabgrass
- (large crabgrass)
- (smooth crabgrass)
- foolsparsley
- foxtails
- (bottlegrass)
- (bristlegrass)
- (giant foxtail)
- (green foxtail)
- (pigeongrass)
- (robust foxtail)
- geranium, Carolina
- goosegrass
- henbit
- knotweed, prostrate
- mallow, little
- parsley - pier
- pennywort, lawn
- pineappleweed
- speedwell, thymeleaf
- spurge, prostrate
- spurge, spotted
- spurry, corn
- woodsorrel, creeping
- woodsorrel, yellow


**Weeds suppressed or partially controlled by a single application or controlled by repeat applications when applied prior to weed germination:**

- clover spp.<sup>1</sup>
- clover, white<sup>1</sup>
- dandelion<sup>1</sup>
- groundsel, common
- medic, black
- plantain, broadleaf
- plantain, buckhorn
- plantain, slender

<sup>1</sup>From seed only

**Uses**

**Established Home Lawns**



**Application Timing:** Apply in late summer or early fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

#### **Broadcast Application Rates**

Apply this product at a rate of 4 to 6 lb per 1000 sq ft. A minimum of 4 lb per 1000 sq ft is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

#### **Established Ornamental Plants in Landscape Settings**

This product is recommended as a preemergence application for control of certain broadleaf weeds and annual grasses in a wide variety of established ornamental trees, shrubs, and ground covers. Apply only to non-bearing fruit and nut trees and non-bearing vineyards which are defined as plants that will not bear fruit for at least 1 year after treatment.

**Application Timing:** Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds. Do not apply this product to plantings when the foliage is wet.

#### **Broadcast Application Rates**

Apply this product at a rate of 4 to 6 lb per 1000 sq ft as a spring application or 6 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

---

#### **Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

#### **Warranty Disclaimer**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

#### **Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

---

#### **Limitation of Remedies**

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.





4/14

308/2E  
April 9, 2008



Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
U. S. Environmental Protection Agency  
One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

TURF FERTILIZER – CONTAINS GALLERY PLUS TEAM PRO (A.I. BENEFIN, TRIFLURALIN, ISOXABEN)  
EPA REGISTRATION NUMBER: 62719-565  
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICE 98-10

Enclosed please find labeling for the notification action of Turf Fertilizer – Contains Gallery® plus Team® Pro specialty herbicide. The following changes have been made by notification:

1. Updated trademark symbol throughout.
2. Added to list of optional label claims.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Contents of Submission**

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled Turf Fertilizer-Contains Gallery plus Team Pro (Q8B/Turf Fertilizer - Contains Gallery Plus Team Pro/MSTR Notif/04-09-08) (37 Pages plus Registration Notes) (5 Copies)

If you require further information, please contact Cindy Loy, Regulatory Specialist at (317) 337-4655 or Paula McKinnies, Registration Assistant for this product, at (317) 337-4679.

Sincerely,

Kimberly S. Gilbert  
Regulatory Leader  
(317) 337-4685  
(317) 337-4649 (FAX)

Enclosures

