

62719-577

01/12/2012

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Niamh McMahon
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

JAN 12 2012

Subject: Supplemental Label for Post Emergence broadleaf Weed Control
in Triticale
Starane® Ultra
EPA Reg. No: 62719-577
Application Dated: October 18th 2011

Dear Ms. McMahon

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

The supplemental labeling must be incorporated into the main product labeling within eighteen (18) months of the date of this letter.

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Grant Rowland at rowland.grant@epa.gov or 703-347-0254.

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Starane[®] Ultra EPA Reg. No. 62719-577

Postemergence Broadleaf Weed Control in Triticale

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane[®] Ultra herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Starane Ultra according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Starane Ultra.

This supplemental label expires on January 17, 2015, and must not be used or distributed after this date.

Directions for Use


Refer to product label for Application Precautions and Restrictions, Mixing and Application directions.

Apply as a broadcast postemergence treatment to actively growing triticale, from the 2 leaf crop growth stage up to and including flag leaf emergence (Zadoks scale 39), for control of broadleaf weeds. Apply when weeds are actively growing, but before weeds are 8 inches tall or vining. For control of volunteer potatoes, apply before potato plants are 8 inches tall. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions, such as, drought or near freezing temperatures, prior to, and following time of application, may reduce weed control and increase the risk of crop injury at all stages of growth. **Do not use if cereal crop is underseeded with a legume.**

Spot Application: Spot applications may be made; however, to prevent over application, apply spot treatments at rates and spray volumes equivalent to broadcast application. See instructions for Spot Treatment in Application Directions section on the label affixed to the container.

Broadcast Application Rates:

Weed Size or Species ¹	Application Rate (pint/acre)
susceptible broadleaf weed seedlings less than 4 inches tall ²	0.3
susceptible broadleaf weed seedlings less than 8 inches tall or vining	0.4
volunteer potatoes	0.7



¹ See Weeds Controlled or Suppressed section on the label affixed to the container for a complete listing of weeds controlled or suppressed.

² The 0.3 pint per acre rate will generally provide satisfactory control of kochia seedlings less than 4 inches tall (including ALS resistant biotypes). However, when conditions for control are less favorable, such as under drought or cool temperatures, the 0.4 pint per acre rate will provide more consistent control of kochia seedlings 1 to 4 inches tall. Control of small kochia with reduced rates will be more consistent if kochia is at least 1 inch tall. Use the 0.4 pint per acre rate for optimal control of dicamba

tolerant kochia populations (see Management of Kochia Biotypes in the General Information section of the label affixed to the container).

Restrictions:

- **Preharvest Interval:** Do not apply within 14 days of cutting hay or within 40 days before grain and straw harvest.
- Do not allow livestock to graze treated areas or harvest treated forage within 7 days of application.
- Do not apply more than 0.7 pints of Starane Ultra per acre per growing season.
- Do not apply Starane Ultra directly to, or otherwise permit it to come in direct contact with, susceptible crops or desirable plants including, alfalfa, canola, cotton, lettuce, edible beans, grapes, lentils, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, or tobacco.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Plantback Restriction:** If replanting is required, plant only those crops listed on this label or the label affixed to the container within 120 days following application.

®Trademark of Dow AgroSciences LLC

R357-019

EPA accepted: ___/___/___

Initial printing