



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 27, 2024

Elaine Bauer  
Regulatory Specialist  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Subject: Label Amendment - Registration Review Mitigation for Aminopyralid  
Product Name: CHAPARRAL  
EPA Registration Number: 62719-597  
Application Date: December 17, 2021  
Decision Number: 580807

Dear Elaine Bauer:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Aminopyralid Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at [carr.caleb@epa.gov](mailto:carr.caleb@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a stylized flourish at the end.

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

**[Sub-Label A for Range and Pasture]**

(Base label):


AMINOPYRALID	GROUP	4	HERBICIDE
METSULFURON-METHYL	GROUP	2	HERBICIDE

**Chaparral™****SPECIALTY HERBICIDE**

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), most warm-season grasses grown for hay, Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites.

**IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS**

- Carefully read the section Restrictions in Hay or Manure Use.
- It is mandatory to follow the Use Restrictions section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied.
- Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.
- For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>

**Forage and Manure Management**

**Warning:** Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.

Not For Sale, Distribution, or Use in New York State.

**ACCEPTED**

Sep 27, 2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 62719-597

**Active Ingredients:**

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro- .....	62.13%
Metsulfuron methyl (Methyl 2-[[[(4-methoxy-6- methyl-1,3,5- triazin-2-yl)-amino]carbonyl] amino]sulfonyl]benzoate) .....	9.45%
Other Ingredients .....	<u>28.42%</u>
Total .....	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

**Keep Out of Reach of Children****WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

---

**Precautionary Statements**

---

**Hazard to Humans and Domestic Animals**

**Causes Substantial but Temporary Eye Injury • Harmful if Swallowed**

**Do not get in eyes or on clothing. Avoid contact with skin.**

**Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate or Viton ≥14 mils
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

## Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**Groundwater Advisory:** This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Users are advised not to apply this product where soils have a rapid to very rapid permeability (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching both surface water and aquatic sediment via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron-methyl and aminopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**Irrigation Water Statement:** Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

## Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

## Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the "Directions for Use" section for information about this standard.

## Nonrefillable rigid containers 5 gallons or less:

### Storage and Disposal

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for

later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Nonrefillable nonrigid containers:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable rigid containers larger than 5 gal:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable rigid containers larger than 5 gal:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water.

Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times.

Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-597

EPA Est. \_\_\_\_\_  
[CD02-xxx-xxx]

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET WEIGHT \_\_\_\_\_**



(Booklet cover / base label for small containers):

<b>AMINOPYRALID</b>	<b>GROUP</b>	<b>4</b>	<b>HERBICIDE</b>
<b>METSULFURON-METHYL</b>	<b>GROUP</b>	<b>2</b>	<b>HERBICIDE</b>

# Chaparral™


## SPECIALTY HERBICIDE

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), most warm-season grasses grown for hay, Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites.

### IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- Carefully read the section Restrictions in Hay or Manure Use.
- It is mandatory to follow the Use Restrictions section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied.
- Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.
- For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>

### Forage and Manure Management



**Warning:** Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.

Not For Sale, Distribution, or Use in New York State.



## Active Ingredients:

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro- .....	62.13%
Metsulfuron methyl (Methyl 2-[[[(4-methoxy-6- methyl-1,3,5- triazin-2-yl)-amino]carbonyl] amino]sulfonyl]benzoate) .....	9.45%
Other Ingredients .....	<u>28.42%</u>
Total .....	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

**Keep Out of Reach of Children****WARNING      AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the "Directions for Use" section for information about this standard.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-597

EPA Est. \_\_\_\_\_  
[CD02-xxx-xxx]

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET WEIGHT** \_\_\_\_\_

(Booklet page 1 through end):

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## WARNING

**Causes Substantial but Temporary Eye Injury • Harmful if Swallowed**

**Do not get in eyes or on clothing. Avoid contact with skin.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate or Viton  $\geq 14$  mils
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**Groundwater Advisory:** This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Users are advised not to apply this product where soils have a rapid to very rapid permeability (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching both surface water and aquatic sediment via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron-methyl and aminopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**Irrigation Water Statement:** Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

#### **Windblown Soil Particles Advisory**

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

---

### **Directions for Use**


---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Not For Sale, Distribution, or Use in New York State.**

**Not for use on pastures in Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont. All other labeled uses are permitted in these states including grazed areas in and around approved use sites.**



Light grey = states where use in pastures is not permitted  
Dark grey = NY where the product is not registered

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate or Viton  $\geq 14$  mils
- Shoes plus socks

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Do not enter or allow people or pets to enter the treated area until sprays have dried.

**Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Nonrefillable rigid containers 5 gallons or less:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows:

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Nonrefillable nonrigid containers:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable rigid containers larger than 5 gal:**

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty

the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable rigid containers larger than 5 gal:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Chaparral™ may be applied by aerial or ground equipment to control susceptible broadleaf weeds and certain woody plants, including invasive and noxious weeds on rangeland, permanent grass pastures (including grasses grown for hay), most warm-season grasses grown for hay, CRP acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites without injury to most grasses.

It is permissible to treat non-irrigation ditch banks, seasonally dry wetlands (such as flood plains, deltas, marshes, swamps, or bogs), and transitional areas between upland and lowland sites only when dry. Chaparral can be used up to the water's edge. Do not apply directly to water and take precautions to minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent, or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize overspray to open water. Note: Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

## Weed Resistance Management

This product contains aminopyralid, a Group 4 synthetic auxin, and metsulfuron-methyl, a Group 2 acetolactate synthase (ALS). Appropriate resistance management strategies should be followed.

Development of plant populations resistant to the mode of action of aminopyralid is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications. There may be resistant weed biotypes to metsulfuron and adequate control of these species cannot be expected.

To delay herbicide resistance, take one or more of the following steps:

- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to a herbicide. Application of a herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where identified, spreading of resistant weeds to other fields may be prevented by cleaning harvesting and tillage equipment before moving to other areas and by planting weed-free seed.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants

mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as mowing.

- Use tank mixtures with herbicides from a different group if such use is permitted. Where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- For further information or to report suspected resistance, contact your extension specialist or contact a Corteva Agriscience customer service representative at 800-992-5994.

## **Mandatory Spray Drift Management**

### **Aerial Applications**

- Do not release spray at a height greater than 10 feet above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to select a coarse or coarser droplet size (ANSI/ASABE S641).
- For all other applications, applicators are required to select a medium or coarser droplet size (ANSI/ASABE S641).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply during temperature inversions.

### **Ground Boom Applications**

- For applications on pastures and rangeland, do not release spray at a height greater than 4 feet above the ground. For all other uses, do not release spray at a height greater than 3 feet above the ground or vegetative canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ANSI/ASAE S572.3).
- For all other applications, applicators are required to use a medium or coarser droplet size (ANSI/ASAE S572.3).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

### **Boom-less Ground Sprayer Applications**

- Applicators are required to select a nozzle and pressure combination that delivers a medium or coarser droplet size (ANSI/ASAE S572.3) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

## **Spray Drift Advisories**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

**Boom-less Ground Applications:** Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.



**Handheld Technology Applications:** Take precautions to minimize spray drift.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### **Controlling Droplet Size – Ground Boom**

- **Volume:** Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure:** Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle:** Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size – Aircraft**

- **Adjust Nozzles:** Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift

## Use Restrictions

**Consult with a Corteva Agriscience representative if you do not understand the Use Restrictions. Call 800-258-3033 for more information.**

<p style="text-align: center;"><b>IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS</b></p> <ul style="list-style-type: none"> <li>Carefully read the section Restrictions in Hay or Manure Use.</li> <li>It is mandatory to follow the Use Restrictions section of this label.</li> <li>Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.</li> <li>Hay can only be used on the farm or ranch where product is applied.</li> <li>Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.</li> <li>For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <a href="https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides">https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides</a></li> </ul>	<p style="text-align: center;"><b>Forage and Manure Management</b></p> <p style="text-align: center;"><b>Warning: Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.</b></p>
--	--

- Do not use grasses treated with Chaparral in the preceding 18 months for hay intended for export outside the United States.**
- Do not use hay or straw from areas treated with Chaparral within the preceding 18 months or manure from animals feeding on hay treated with Chaparral in compost.**
- Do not use grasses treated with Chaparral in the preceding 18 months for seed production.**

**Maximum Application Rate:** On all labeled use sites do not broadcast apply more than 3.3 ounce/acre of Chaparral per year. The total amount of Chaparral applied broadcast, as a re-treatment, and/or spot treatment cannot exceed 3.3 oz of product per acre per year. Spot treatments may be applied at an equivalent broadcast rate of up to 6.6 oz product of Chaparral per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 3.3 ounce/acre of Chaparral per annual growing season as a result of broadcast, spot, or repeat applications.

- Do not use on Timothy hay or other cool-season grasses grown for hay.
- Do not apply this product on lawns, turf, ornamental plantings, urban walkways, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
- Do not overseed ryegrass for 4 months after treatment.

- **Chaparral is highly active against many broadleaf plant species.** Do not use this product on areas where loss of broadleaf plants, including legumes, cannot be tolerated.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Do not contaminate water intended for irrigation or domestic purposes.** Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not apply to irrigated land where the tailwater will be used to irrigate crops.
- **Do not use this product for impregnation on dry fertilizer unless specified in Corteva Agriscience state-specific product bulletin.**
- Do not use Chaparral in the following counties of Colorado: Alamosa, Conejos, Costilla, Rio Grande, and Saguache.
- **Trees** adjacent to or in a treated site can occasionally be affected by root uptake of Chaparral. Do not apply Chaparral within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.
  - Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots extend, or in locations where the product may be washed or moved into contact with their roots, as injury or loss of desirable trees or other plants may result.
- Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of Chaparral. Injury to crops may result if treated soil and/or runoff water containing Chaparral is washed, or moved onto land used to produce crops. Exposure to Chaparral may injure or kill susceptible crops and other plants, such as grapes, soybeans, tobacco, sensitive ornamentals. Do not treat frozen soil where runoff could damage sensitive plants.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if aminopyralid or metsulfuron concentration remaining in the soil will adversely affect the legume establishment.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after Chaparral application, temporary discoloration, and/or grass injury may occur. Chaparral should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Do not apply to frozen ground as surface runoff may occur.
- Do not apply to snow-covered ground.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or grass hay harvest intervals following application of Chaparral at labeled rates. However, cutting hay too soon after spraying weeds will reduce weed control. Wait 14 days after herbicide application to cut grass hay to allow herbicide to work.
- **Grazing Poisonous Plants:** Application of this product may increase palatability of certain poisonous plants. Do not allow livestock to graze areas treated with Chaparral until poisonous plants are dry and no longer palatable to livestock.
- **Restrictions in Hay or Manure Use:**
  - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18 months, in compost, mulch, or mushroom spawn.
  - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch, or mushroom spawn.
  - Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields treated with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.

- To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- **Off-site Compost and Animal Feed/Bedding Restrictions**
  - This product is persistent and may be present in treated plant materials for months to years after application. Do not sell or transport treated plant materials or manure from animals that have grazed on treated plant materials off-site for compost distribution or for use as animal bedding/feed for 18 months after application. Treated plant materials can be recycled onsite or left in the field to decompose.
  - Manure from animals that have grazed or eaten forage or hay harvested from treated areas within the previous 3 days may only be applied to the fields where the following crops will be grown: pasture grasses, grass grown for seed, wheat, and corn.
  - Animals that have been fed aminopyralid-treated forage must be fed forage free of aminopyralid for at least 3 days before movement to an area where manure may be collected or sensitive crops are grown.
- **Crop Rotation:** Do not rotate to any crop from rangeland, permanent pasture, or CRP acres within one year following treatment. Cereals and corn can be planted one year after treatment. Most broadleaf crops are more sensitive and can require **at least 2 years** depending on the crop and environmental conditions. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid or metsulfuron present in the soil will not adversely affect that broadleaf crop.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern, or drainage. The field bioassay can be one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, perennial forage grasses, native grasses or grasses grown for hay.
- **Avoiding Injury to Non-Target Plants:** Do not aerially apply Chaparral within 50 feet of a border downwind (in the direction of wind movement), or allow spray drift to come in contact with, any broadleaf crop or other desirable broadleaf plants, including, but not limited to, alfalfa, cotton, dry beans, flowers, grapes, lettuce, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes or other broadleaf or vegetable crop, fruit trees, ornamental plants, or soil where sensitive crops are growing or will be planted. Avoid application under conditions that may allow spray drift because very small quantities of spray may seriously injure susceptible crops. Read the Spray Drift Management section of this label to help minimize the potential for spray drift.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, plant residue mulch, reduced tillage, or other cultural practices. Injury to immediately adjacent crops may occur when treated soil is blown onto land used to produce crops other than pasture, rangeland, or CRP.

#### **Record Keeping For Applications to Pastures**

- The applicator must document that they have notified property owners/operators or customers, in writing, of the compost and animal feed/bedding prohibitions within 14 days of the application. Applicators must keep the records of notification for two years. This record must include date of application, the name of the applicator, the EPA registration number of the product applied, area(s) treated, and a copy of the written notification provided to the property owner/operator. Notification may be made via email, via mail, via paper handout, or by any other written communication method. Records must be made available to State Pesticide Regulatory Official(s) and to EPA upon request. If this information is already being retained, duplicate records are not needed.
- It is recommended that applicators also transmit at the time of notification relevant educational materials for managing treated plant matter, as available. Additional educational materials for aminopyralid will be posted at: <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>
- Applications to pastures by property owners/operators on their own property are exempt from this notification and recordkeeping requirement.

- Application to pastures on public land (i.e., lands managed directly by state, tribal, or local authorities) are exempt from this notification requirement.

### **Sprayer Clean-Out Instructions**

**It is recommended to use separate spray equipment on highly sensitive crops such as tobacco, soybeans, peanuts, and tomatoes.**

Do not use spray equipment used to apply Chaparral for other applications to land planted to, or to be planted to, broadleaf plants unless it has been determined that all residues of this herbicide has been removed by thorough cleaning of equipment.

Equipment used to apply Chaparral should be thoroughly cleaned before reusing to apply any other chemicals as follows:

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
2. Rinse a second time, adding 1 quart of household ammonia or tank cleaning agent for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Spray nozzles and screens should be removed and cleaned separately.

Do not apply this product with mist blower systems that deliver very fine spray droplets. Use of mist blower equipment can reduce control achieved with the herbicide and increase spray drift potential.

### **Application Methods**

Apply the specified rate of Chaparral as a coarse low-pressure spray. Do not apply this product with mist blower systems that deliver very fine spray droplets. Spray volume should be sufficient to uniformly cover foliage. Increase the spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. To enhance foliage wetting and coverage, an approved non-ionic agricultural surfactant may be added to the spray mixture as specified by the surfactant label.

**Ground Broadcast Application:** Higher spray volumes (greater than 10 gallons per acre) generally provide better coverage and better control, particularly in dense and/or tall foliage.

**Aerial Broadcast Application:** Do not apply less than 2 gallons per acre total spray volume. Five gallons per acre or greater will generally provide better coverage and better control, particularly in dense and/or tall foliage.

**High-Volume Foliar Application:** High volume foliar treatments may be applied at rates equivalent to a maximum of 3.3 ounces per acre annual growing season. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems.

**Spot Application:** Spot treatments may be applied at an equivalent broadcast rate of up to 6.6 oz of product per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 3.3 ounce/acre of Chaparral per annual growing season as a result of broadcast, spot, or repeat applications. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage, but not to the point of runoff. Repeat treatments may be made, but the total amount of Chaparral applied must not exceed 3.3 ounce/acre per year. Take precautions to minimize spray drift. To prevent misapplication, spot treatments should be applied with a calibrated sprayer.

In general, for spot treatments mix 2.5 oz for weeds and 3.3 oz for brush of Chaparral per 100 gallons of water (assuming an application volume of 100 gallons per acre).

### **Product Measurement**

Chaparral is measured using the Chaparral volumetric measuring cylinder. Scales calibrated in ounces may also be used.



**Mixing Instructions**

1. Fill the tank 1/4 to 1/3 full with water. (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures section for additional details.)
2. While agitating, add the required amount of Chaparral.
3. Continue agitation until the Chaparral is fully dispersed, at least 5 minutes.
4. Once the Chaparral is fully dispersed, maintain agitation and continue filling tank with water. Chaparral should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of spray adjuvants. Always add spray adjuvants last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply Chaparral spray mixture within 24 hours of mixing to avoid product degradation.
8. If Chaparral and a tank mix partner are to be applied in multiple loads, pre-slurry the Chaparral in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Chaparral.

**Soil pH Limitations**

Chaparral should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Chaparral could remain in the soil for 34 months or more injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of Chaparral.

**Checking Soil pH**

Before using Chaparral, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil-sampling procedures.

**Spray Adjuvants**

Unless otherwise directed, applications of Chaparral must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer can be used unless specifically prohibited by tank mix partner labeling. If another herbicide is tank mixed with Chaparral, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

**Petroleum Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO)**

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- MSO adjuvants may be used at 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

**Nonionic Surfactant (NIS)**

- Apply at 0.25% v/v (1 quart per 100 gallons spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

**Ammonium Nitrogen Fertilizer**

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 pounds/acre of a spray grade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.
- Pensacola bahiagrass control can be reduced when Chaparral is applied in liquid fertilizer solutions.

**Mixing With Liquid Nitrogen Solution Fertilizer**

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing Chaparral in fertilizer solution.



Chaparral must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the Chaparral is added. Use of this mixture may result in temporary grass yellowing and stunting.

If using low rates of liquid nitrogen fertilizer (less than 50% of the spray solution volume) in the spray solution, the addition of a non-ionic surfactant is necessary. Add surfactant at 1/4 pt per 100 gal of spray solution (0.03% v/v).

Do not use a spray adjuvant other than non-ionic surfactant. When using high rates of liquid nitrogen fertilizer (greater than or equal to 50% of the spray solution volume) in the spray solution, adding spray adjuvant(s) increases the risk of grass injury. Consult your agricultural dealer, consultant, or Corteva Agriscience representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with Chaparral and the fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Do not add spray adjuvants when using Chaparral in tank mix with 2,4-D ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid fertilizer as a substitute for spray adjuvants.  
Do not use with liquid fertilizer solutions with a pH less than 3.0.

### Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO, and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions. Exception: On tall fescue pastures use a reduced rate of 1/2 to 1 pint non-ionic surfactant per 100 gallons.
- Antifoaming agents may be used if needed.
- Do not use Chaparral with spray additives that reduce the pH of the spray solution to below 3.0.

**Tank Mixing with Other Herbicides:** Chaparral at rates of up to 3.3 ounce/acre may be mixed with labeled rates of other herbicides registered for application on all labeled use sites. Chaparral may be applied in tank-mix combination with labeled rates of other herbicides provided: (1) the tank-mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products, and (3) that the tank-mix combination is physically compatible (see tank-mix compatibility testing below). When tank mixing, use only in accordance with the restrictions, precautions, and limitations on the respective product labels.

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not exceed specified application rates. If products containing the same active ingredient are mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** Perform a jar test prior to mixing in a spray tank to ensure compatibility of Chaparral and other pesticides or carriers. Use a clear glass jar with lid and mix ingredients in the same order and proportions as will be used in the spray tank. The mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily remix if agitated. An incompatible mixture is indicated by separation into distinct layers that do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film in the jar. Use of an appropriate compatibility aid may resolve mix incompatibility. If the mixture is incompatible do not use that tank mix partner in tank mixtures.

**Note:** Foliar-applied liquid fertilizers themselves can cause yellowing of the foliage of forage grasses and other vegetation.

**Seeding grasses:**

**Preemergence:** In general, Chaparral may be applied in the spring or early summer, depending on the target weed species, as a broadcast application over grass that was planted at least 4 months prior to the application and that has been growing under favorable conditions for grass establishment.

With fall applications, do not plant grasses the following spring.

Do not overseed ryegrass for 4 months after treatment.

**Application Timing – Established Pastures, Rangeland, and CRP**

Chaparral may be applied to established native grasses such as wheatgrasses, bluestems, and grama, and to other established pasture grasses such as bermudagrass, bluegrass, orchardgrass, brome grass, and tall fescue that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

- Chaparral may suppress certain established grasses such as smooth brome grass (*Bromus inermis*), especially when plants are stressed by adverse environmental conditions. Plants should recover from this transient suppression with the onset of environmental conditions favorable to grass growth and upon release from weed competition.
- Varieties and species of forage grasses differ in their tolerance to herbicides. When using Chaparral on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated.
- Application of Chaparral to Pensacola bahiagrass, ryegrass (Italian or perennial), and Garrison's creeping foxtail may cause severe injury to and/or loss of pastures.

**Tall Fescue Precautions:**

Chaparral may stunt tall fescue, cause it to turn yellow, or cause seed head suppression. To minimize these symptoms, take the following precautions.

- Do not use on tall fescue grown for seed.
- Do not use more than 2 ounce/acre of Chaparral.
- Tank-mix Chaparral with 2,4-D.
- Use a reduced rate of non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution (1/16 to 1/8 % v/v).
- Make application later in the spring after the new growth is 5 to 6 inches tall (until after reproductive culm has started to elongate), or in the fall.
- Do not use surfactant when liquid nitrogen is used as a carrier.
- Do not use a spray adjuvant other than non-ionic surfactant.

Initial grass yields may be reduced due to fescue seed head suppression resulting from treatment with Chaparral at labeled rates. However, this could be beneficial because in tall fescue infected with the fungal endophyte (*Neotyphodium* spp.), the endophyte is concentrated in the seed, and cattle grazing plants with the seed head will get the maximum exposure to the endophyte. Increased levels of ingestions of the fungal endophyte can reduce weight gain and conception rates in cattle. Since the first grazing is often delayed in the spring until long after seed head development, Chaparral could potentially be used to reduce development of the seed head, thereby reducing the amount of the endophyte that would be consumed by livestock when grazing (see below).

**Tall Fescue Seed Head Suppression and Broadleaf Weed Control:** Chaparral herbicide can be used to reduce the number of seed heads of tall fescue when applied prior to flower emergence. For best results apply 2.0 to 2.5 ounce/acre Chaparral after initial greenup when grass height is approximately 6 inches. Later applications may still be effective, however, the seed head suppression will be less effective, and the number of seed heads could be noticeable higher. Many weed species can be controlled with this application timing in addition to the suppression of seed head development.

## Use Rates and Timing

Chaparral may be applied post emergence as a broadcast spray or as a spot application to control weeds and brush including, but not limited to, those listed on this label. When a rate range is given use the higher rate to control weeds at advanced growth stages, or under less than favorable growing conditions, or for longer residual control. Best results are obtained when spray volume is sufficient to provide uniform coverage of treated weeds. For optimum uptake and translocation of Chaparral, avoid mowing, haying, shredding, burning, or soil disturbance in treated areas for at least 14 days following application.

Chaparral also provides preemergence control of emerging seedlings of susceptible weeds, and re-growth of certain perennial weeds following application. Preventing establishment of weeds will depend upon application rate, season of application, and environmental conditions after application.

Chaparral can provide long-term control of susceptible weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Long-term weed control is most effective where grass vegetation is allowed to recover from overgrazing, drought, etc., and compete with weeds.

Chaparral can be an important component of integrated vegetation management programs designed to renovate or restore desired plant communities. To maximize and extend the benefits of weed control provided by Chaparral, it is important that other vegetation management practices, including proper grazing management, biological control agents, replanting, fertilization, prescribed fire, etc., be used in appropriate sequences and combinations to further alleviate the adverse effects of weeds on desirable plant species and to promote development of desired plant communities. Agricultural and natural resources specialists with federal and state government agencies can provide guidance on best management practices and development of integrated vegetation management programs.

## Species Controlled

**Table 1: Grass Weeds Controlled or Suppressed by Chaparral**

**Note:** Weeds marked with a \* indicate more information is included in the specific weed problems section after the table.

**Grass Weeds Controlled by Chaparral**

Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
arrowgrass, seaside‡	<i>Trifglochin maritima</i>	perennial	Juncaginaceae	3.0-3.3
bahiagrass, Pensacola*	<i>Paspalum notatum</i> <i>Flugge</i>	perennial	Poaceae	2.0-2.5
ryegrass	<i>Lolium multiflorum</i>	annual	Poaceae	2.0-2.5

‡ This symbol denotes weed suppression which is a reduction in weed competition compared to untreated areas. A second treatment may be necessary.

### Pensacola bahiagrass control in established Bermudagrass pasture:

Apply Chaparral at 2-2.5 ounce/acre after green-up in the spring, but before bahiagrass seed head formation. Application should be made when environmental conditions favor grass growth.

Bahiagrass suppression could take up to 30 days before the desired level of control is achieved. Application of 2,4-D mixed with Chaparral could decrease bahiagrass control.

In pastures severely infested with bahiagrass, a positive response in forage yield may be slowed until desired forage grasses like bermudagrass grow into areas previously infested with bahiagrass. To reduce this effect, consider treating different portions of heavily infested pastures with Chaparral over a period of several years. Do not apply Chaparral to an entire farm or ranch in one year. Fertilization and/or replanting may accelerate bermudagrass recovery following bahiagrass control with Chaparral.

Bahiagrass regrowth may occur in pastures heavily infested with bahiagrass, intense grazing pressure, or when adverse environmental conditions (heat and drought), slows the recovery of desired grass forages.

Chaparral will not control common or Argentine bahiagrass.

Pensacola bahiagrass control can be reduced when Chaparral is applied in liquid fertilizer solutions.

**General Mix of Broadleaf Weeds:** Chaparral at 2.0 ounce/acre is the standard rate to provide control of most pasture weeds when applied early in the season. If a certain weed is dominant in the pasture, use the rate in Table 2 for that species. The addition of 0.5 lbs ae/acre (1 pint/acre of 4 lb ae/gallon 2,4-D) 2,4-D amine may broaden the weed spectrum.

For rates for specific weeds, see Table 2. The life cycle is included for each weed species. The general timing of application for each life cycle is as follows:

**Annuals:** Use lower rates when weeds are less than 6 inches and actively growing. Increase rate as season progresses and plants become more mature.

**Biennials:** Apply in the spring and early summer to rosette or bolting plants or in the fall to seedlings and rosettes before ground is frozen. Use higher rates after bolting through early flower.

**Perennials:** Apply to vegetative stage prior to bloom. Use higher rate when weeds are larger.

For best results, most weeds should be treated when they are actively growing and under conditions favorable for growth. Use a higher rate in the rate range indicated when growing conditions are less than favorable (drought conditions), weeds are large and mature, weed density and foliage cover is high and canopy height is tall, or when residual control is desired. Chaparral also provides preemergence control of germinating seeds or seedlings of susceptible weeds following application.

**Table 2: Broadleaf Species Controlled with Chaparral**

**Note:** Weeds marked with \* indicate more information is included in the specific weed problems section after the table.

**Broadleaf Weeds Controlled by Chaparral**

Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
actinomeris, wingstem	<i>Verbesina alternifolia</i>	perennial	Asteraceae	3.0
alyssum, hoary	<i>Berteroa incana</i>	biennial	Brassicaceae	2.0-2.5
amaranth, spiny	<i>Amaranthus spinosus</i>	summer annual	Amaranthaceae	1.5-2.0
aster	<i>Aster spp.</i>	perennial	Asteraceae	1.5-2.0
babysbreath	<i>Gypsophila paniculata</i>	perennial	Caryophyllaceae	2.5-3.0
bedstraw	<i>Galium spp.</i>	perennial	Rubiaceae	2.0-2.5
beggarticks	<i>Bidens spp.</i>	annual	Asteraceae	2.0-3.0
bittercress	<i>Cardamine spp.</i>	perennial	Brassicaceae	2.0-2.5
blackeyed-susan	<i>Rudbeckia hirta</i>	annual	Asteraceae	1.5-2.0
brackenfern	<i>Pteridium spp.</i>	perennial	Dennstaedtiaceae	2.5-3.3
broomweed, annual	<i>Amphichayris dracunculoides</i>	annual	Asteraceae	1.0-1.5
bur buttercup (testiculate)	<i>Ranunculus testiculatus</i>	annual	Ranunculaceae	1.0-1.5
burclover	<i>Medicago spp.</i>	annual	Fabaceae	1.5-2.0
burdock, Common	<i>Arctium minus</i>	biennial	Asteraceae	2.0-2.5
buttercup, hairy	<i>Ranunculus sardous</i>	perennial	Ranunculaceae	1.0-1.5
buttercup, tall	<i>Ranunculus acris</i>	perennial	Ranunculaceae	2.0-2.5
camelthorn	<i>Alhagi pseudalhagi</i>	perennial	Fabaceae	2.0-3.0
cat's ear, common	<i>Hypochaeris radicata</i>	perennial	Asteraceae	2.5-3.3

Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
camphorweed	<i>Heterotheca subaxillaris</i>	summer annual	Asteraceae	2.0-3.0
campion, bladder†	<i>Silene vulgaris</i>	perennial	Caryophyllaceae	2.0-2.5
caraway, wild	<i>Carum carvi</i>	biennial	Apiaceae	2.5-3.0
carrot, wild	<i>Daucus carota</i>	biennial	Apiaceae	2.0-2.5
catchfly, conical	<i>Silene conoidea</i>	annual	Caryophyllaceae	1.0-1.5
chamomile	<i>Matricaria spp</i>	annual	Asteraceae	2.5-3.0
chickweed, common	<i>Stellaria media</i>	winter annual	Caryophyllaceae	3.0
chicory	<i>Cichorium intybus</i>	perennial	Asteraceae	1.5-2.0
cinquefoil*	<i>Potentilla spp</i>	perennial	Rosaceae	2.0-2.5
clover, sweet	<i>Melilotus officinalis</i>	biennial	Fabaceae	2.5-3.0
clover, white	<i>Trifolium repens</i>	perennial	Fabaceae	1.5-2.0
cockle, corn	<i>Agrostemma githago</i>	annual	Caryophyllaceae	2.0-3.0
cocklebur	<i>Xanthium strumarium</i>	annual	Asteraceae	1.5-2.0
coreopsis, plains	<i>Coreopsis tinctoria</i>	annual	Asteraceae	2.0-3.0
cowcockle	<i>Vaccaria pyramidata</i>	annual	Caryophyllaceae	1.5-2.0
crazyweed, silky	<i>Oxytropis Lambertii</i>	perennial	Fabaceae	2.0-2.5
croton, woolly	<i>Croton capitatus</i>	annual	Euphorbiaceae	1.5-2.0
crownvetch	<i>Securigera varia</i>	perennial	Fabaceae	1.5-2.0
crupina, common	<i>Crupina vulgaris</i>	perennial	Asteraceae	3.0-3.3
cudweed, purple	<i>Gnaphalium purpureum</i>	annual	Asteraceae	2.0-2.5
daisy, oxeye*	<i>Leucanthemum vulgare</i>	perennial	Asteraceae	2.5-3.3
dandelion, common	<i>Taraxacum officinale</i>	perennial	Asteraceae	1.5-2.0
dock	<i>Rumex spp</i>	perennial	Polygonaceae	2.0-2.5
dyer's woad†	<i>Istis tinctoria</i>	perennial	Brassicaceae	3.3
evening primrose, cutleaf	<i>Oenothera laciniata</i>	annual	Asteraceae	1.5-2.0
false dandelion, Carolina	<i>Tragopogon dubius</i>	biennial	Asteraceae	1.5-2.0
falseflax, smallseed	<i>Camelina microcarpa</i>	annual/ biennial	Brassicaceae	1.5-2.0
fiddleneck, common	<i>Amsinckia intermedia</i>	annual	Boraginaceae	1.5-2.0
filaree, redstem	<i>Erodium cicutarium</i>	annual/ biennial	Geraniaceae	3.0-3.3
fireweed	<i>Epilobium angustifolium</i>	perennial	Onagraceae	2.5-3.0
fleabane, annual	<i>Erigeron annuus</i>	annual	Asteraceae	1.5-2.0
fleabane, hairy	<i>Conyza bonariensis</i>	annual/ biennial	Asteraceae	2.5-3.3
garlic. wild	<i>Allium vineale</i>	perennial	Liliaceae	1.5-2.0
geranium, Carolina	<i>Geranium carolinianum</i>	winter annual	Geraniaceae	1.5-2.0
goldenrod spp	<i>Solidago canadensis</i>	perennial	Asteraceae	2.0-2.5
gumweed, curlycup	<i>Grindelia squarrosa</i>	biennial	Asteraceae	2.0-2.5
halogeton	<i>Halogeton glomeratus</i>	annual	Chenopodiaceae	3.0-3.3
hawkweed, orange*	<i>Hieracium aurantiacum</i>	perennial	Asteraceae	2.5-3.3
hawkweed, yellow*	<i>Hieracium pratense</i>	perennial	Asteraceae	2.5-3.3



Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
hemlock, poison‡	<i>Conium maculatum</i>	perennial	Apiaceae	2.5-3.3
henbane, black	<i>Hyoscyamus niger</i>	annual/ biennial	Solanaceae	2.5-3.0
henbit	<i>Lamium amplexicaule</i>	annual/ biennial	Lamiaceae	2.0-2.5
hogweed, giant	<i>Heracleum mantegazzianum</i>	perennial	Apiaceae	3.3
horsemint (beebalm)	<i>Monarda spp</i>	annual	Lamiaceae	1.5-2.0
horsenettle, Carolina	<i>Solanum carolinense</i>	perennial	Solanaceae	2.0-2.5
horseweed (maretail)	<i>Conyza canadensis</i>	annual	Asteraceae	1.5-2.0
houndstongue*	<i>Cynoglossum officinale</i>	biennial	Boraginaceae	2.5-3.3
ironweed, tall	<i>Vernonia gigantea</i>	perennial	Asteraceae	2.0-3.0
ironweed, western	<i>Vernonia baldwinii</i>	perennial	Asteraceae	2.0-3.0
knapweed	<i>Centaurea sp.</i>	biennial	Asteraceae	2.5-3.3
knapweed, brown	<i>Centaurea jacea</i>	perennial	Asteraceae	2.5-3.3
knapweed, diffuse*	<i>Centaurea diffusa</i>	biennial	Asteraceae	2.5-3.3
knapweed, Russian*	<i>Acroptilon repens</i>	perennial	Asteraceae	2.5-3.3
knapweed, spotted*	<i>Centaurea stoebe</i>	biennial	Asteraceae	2.5-3.3
knotweed, prostrate	<i>Polygonum aviculare</i>	annual	Polygonaceae	3.0
kochia*	<i>Kochia scoparia</i>	annual	Chenopodiaceae	1.5-2.0
lady's thumb	<i>Polygonum persicaria</i>	annual	Polygonaceae	1.5-2.0
lambquarters, common	<i>Chenopodium album</i>	annual	Chenopodiaceae	2.0-2.5
lespedeza, annual	<i>Lespedeza striata</i>	annual	Fabaceae	2.0-2.5
lespedeza, sericea*	<i>Lespedeza cuneata</i>	perennial	Fabaceae	2.5-3.0
lettuce, Miner's	<i>Montia perfoliata</i>	annual	Portulacaceae	1.5-2.0
lettuce, prickly*	<i>Lactuca serriola</i>	annual	Asteraceae	1.5-2.0
licorice, wild	<i>Glycyrrhiza lepidota</i>	perennial	Fabaceae	2.5-3.0
locoweed	<i>Astragalus spp.</i>	perennial	Fabaceae	2.0-2.5
loosestrife, purple	<i>Lythrum salicaria</i>	perennial	Lythraceae	3.0-3.3
marshelder, annual‡	<i>Iva annua</i>	annual	Asteraceae	2.0-2.5
mayweed, scentless	<i>Tripleurospermum perforata</i>	annual	Asteraceae	1.5-2.0
mayweed, stinking	<i>Anthemis cotula</i>	annual	Asteraceae	3.0-3.3
medic, black	<i>Medicago lupulina</i>	perennial	Fabaceae	2.0-2.5
Mexican-tea	<i>Dysphania ambrosioides</i>	annual/ perennial	Chenopodiaceae	2.0-2.5
mullein*	<i>Verbascum spp.</i>	biennial	Scrophulariaceae	2.0-3.3
mustard, blue*	<i>Chorispora tenella</i>	annual	Brassicaceae	1.5-2.0
mustard, garlic	<i>Alliaria petiolata</i>	biennial	Brassicaceae	3.3
mustard, tumble/Jim Hill	<i>Sisymbrium altissimum</i>	winter annual	Brassicaceae	1.5-2.0
mustard, wild	<i>Brassica kaber</i>	annual	Brassicaceae	1.5-2.0
needles, Spanish needles	<i>Bidens bipinnata</i>	annual	Asteraceae	2.0-2.5
nightshade, silverleaf	<i>Solanum elaeagnifolium</i>	perennial	Solanaceae	2.0-3.0
oxtongue, bristly	<i>Picris echioides</i>	biennial	Asteraceae	2.5-3.0



Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
parsnip, Wild	<i>Pastinaca sativa</i>	biennial	Apiaceae	2.0-3.0
partridgepea	<i>Chamaecrista fasciculata</i>	annual	Fabaceae	2.5-3.0
pea, Swainson	<i>Sphaerophysa salsula</i>	perennial	Fabaceae	2.0-3.0
pepperweed, perennial†	<i>Lepidium latifolium</i>	perennial	Brassicaceae	3.3
pigweeds	<i>Amaranthus spp</i>	annual	Amaranthaceae	1.5-2.0
plantain, broadleaf	<i>Plantago major</i>	perennial	Plantaginaceae	2.0-2.5
plantain, buckhorn	<i>Plantago lanceolata</i>	perennial	Plantaginaceae	2.0-2.5
povertyweed	<i>Iva axillaris</i>	perennial	Asteraceae	2.0-3.0
purslane, common	<i>Portulaca oleracea</i>	annual	Portulacaceae	1.5-2.0
ragweed, common	<i>Ambrosia artemisiifolia</i>	annual	Asteraceae	2.0-2.5
ragweed, western*	<i>Ambrosia psilostachya</i>	perennial	Asteraceae	2.0-2.5
ragwort, tansy	<i>Senecio jacobaea</i>	perennial	Asteraceae	2.5-3.0
rush skeletonweed	<i>Chondrilla juncea</i>	perennial	Asteraceae	2.5-3.0
salsify, Western†	<i>Tragopogon dubius</i>	biennial	Asteraceae	3.0-3.3
shephardspurse	<i>Capsella bursa-pastoris</i>	winter annual	Brassicaceae	1.5-2.0
sicklepod	<i>Senna obtusifolia</i>	annual	Fabaceae	2.5-3.0
sida, arrowleaf	<i>Sida rhombifolia</i>	annual	Malvaceae	2-2.5
smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>	annual	Polygonaceae	1.5-2.0
snakeweed, broom*	<i>Gutierrezia sarothrae</i>	perennial	Asteraceae	3.0
sneezeweed, bitter	<i>Helenium amarum</i>	annual	Asteraceae	1.0-1.5
snow-on-the-mountain	<i>Euphorbia marginata</i>	annual	Euphorbiaceae	2.0-2.5
soda apple, tropical*	<i>Solanum viarum</i>	perennial	Solanaceae	2.5-3.0
sorrel, red	<i>Rumex acetosella</i>	perennial	Polygonaceae	2.0-2.5
sowthistle, annual	<i>Sonchus oleraceae</i>	annual	Asteraceae	2.5-3.0
sowthistle, perennial	<i>Sonchus arvensis</i>	perennial	Asteraceae	2.0-2.5
sowthistle, prickly	<i>Sonchus asper</i>	annual	Asteraceae	1.5-2.0
spurge, spotted	<i>Euphorbia maculata</i>	Perennial	Euphorbiaceae	1.5-2.5
St. Johnswort, common	<i>Hypericum perforatum</i>	perennial	Clusiaceae	2.5-3.0
starthistle, purple*	<i>Centaurea calcitrapa</i>	biennial	Asteraceae	1.5-2.0
star-thistle, Malta*	<i>Centaurea melitensis</i>	annual	Asteraceae	1.5-2.0
starthistle, yellow*	<i>Centaurea solstitialis</i>	annual	Asteraceae	1.5-2.0
sunflower, common	<i>Helianthus annua</i>	annual	Asteraceae	1.5-2.0
tansy, common	<i>Tanacetum vulgare</i>	perennial	Asteraceae	2.5-3.3
teasel	<i>Dipsacus spp.</i>	biennial	Dipsacaceae	2.0-3.0
thistle, Russian*	<i>Salsola iberica</i>	annual	Chenopodiaceae	1.5-2.0
thistle, artichoke	<i>Cynara cardunculus</i>	perennial	Asteracea	2.0-3.0
thistle, blessed milk	<i>Silybum marianum</i>	Biennial	Asteraceae	2.0-3.0
thistle, bull*	<i>Cirsium vulgare</i>	biennial	Asteraceae	1.0-2.5
thistle, Canada*	<i>Cirsium arvense</i>	perennial	Asteraceae	2.0-3.3
thistle, Italian	<i>Carduus pycnocephalus</i>	annual	Asteraceae	2.0-3.0
thistle, musk*	<i>Carduus nutans</i>	biennial	Asteraceae	1.0-2.5
thistle, plumeless*	<i>Carduus acanthoides</i>	biennial	Asteraceae	1.0-2.5

Common Name	Scientific Name	Life Cycle	Plant Family	Chaparral rate oz/a
thistle, scotch	<i>Onopordum acanthium</i>	biennial	Asteraceae	1.5-2.5
thistle, woolly distaff	<i>Carthamus lanatus</i>	annual	Asteraceae	1.5-2.0
vervain‡	<i>Verbena spp.</i>	perennial	Asteraceae	2.0-2.5
vetch, common*	<i>Vicia sativa</i>	annual	Fabaceae	1.5-2.0
wallflower, bushy	<i>Erysimum repandum</i>	annual	Brassicaceae	1.5-2.0
waterpod	<i>Ellisia nyctelea</i>	annual	Brassicaceae	1.5-2.0
whitetop (hoary cress)*	<i>Cardaria draba</i>	perennial	Brassicaceae	3.3
woodsorrel, yellow	<i>Oxalis stricta</i>	perennial	Oxalidaceae	3.0-3.3
wormwood, absinth*	<i>Artemisia absinthium</i>	perennial	Asteraceae	3.0-3.3
yankeeweed	<i>Eupatorium compositifolium</i>	perennial	Asteraceae	3.0-3.3
yarrow, common	<i>Achillea millefolium</i>	perennial	Asteraceae	1.5-2.0

‡ This symbol denotes weed suppression which is a reduction in weed competition compared to untreated areas. A second treatment may be necessary. The addition of 0.5 lbs ae/acre of 2,4-D may improve initial control.

**Hawkweed, orange or yellow:** Apply Chaparral at 2.5 to 3.3 ounce/acre to plants in the bolting stage of development.

**Houndstongue:** Apply 2.5 ounce/acre to rosettes. As plant bolts, increase the rate to 3.0 to 3.3 ounce/acre up to early bud stage. Add 1 quart of 2,4-D/acre after the bud stage.

**Ironweed, Western:** Apply Chaparral at 2.5 to 3.0 ounce/acre when plants are in the vegetative growth stage. The addition of 0.5 to 1 lb ae/acre (1 to 2 pints/acre of 4 lb ae/gallon 2,4-D) of 2,4-D/acre will improve control in dense stands or when ironweed is greater than 6 inches.

**Knapweeds, diffuse and spotted:** Apply Chaparral at 2.5 to 3.3 ounce/acre when plants are actively growing with the optimum time of application occurring from rosette to the bolting stages of development or in the fall. Plants will be controlled by mid-summer and fall applications even though plants may not show any changes in form or stature the year of application.

**Knapweed, Russian:** Apply Chaparral at 2.5 to 3.3 ounce/acre to plants in the spring and summer to plants from early bud to flowering stage and to dormant plants in the fall.

**Lespedeza, Sericea:** Apply 2.5 to 3.0 ounce/acre beginning at flower bud initiation through the full bloom stage of growth.

**Mullein:** Apply 2.0 ounce/acre in the rosette stage in spring or fall. Use rates from 2.5 to 3.3 ounce/acre for bolting plants less than 12 inches tall.

**Oxeye daisy:** Apply Chaparral at 2.5 to 3.3 ounce/acre to plants in the prebud stage of development.

**Pepperweed, perennial:** Apply Chaparral at 3.3 ounce/acre plus 2 lb ae/a 2, 4-D when plants are at early flowering through bloom for optimum control.

**Ragweed, Western:** Apply Chaparral at 2.0 to 2.5 ounce/acre when plants are in the vegetative growth stage. The addition of 0.5 to 1 lb ae/acre (1 to 2 pints/acre of 4 lb ae/gallon 2,4-D) of 2,4-D/acre will improve control in dense stands or when ragweed is greater than 6 inches.

**Russian thistle, kochia, and prickly lettuce:** Naturally occurring resistant biotypes of these weeds to metsulfuron are known to occur. For best results, use Chaparral at 1.5 to 2.0 ounces/acre in tank-mix with 2,4-D. Applications to these weeds should be made early to weeds less than 6 inches in height.

**Snakeweed, broom:** Applications should be made in the fall at 3.0 ounces/acre. Spring applications will provide suppression only.

**Soda apple, tropical:** Apply Chaparral at 2.5 to 3.0 ounce/acre at any growth stage, but application by flowering will reduce seed production potential.

**Starthistle, malta, purple, and yellow:** Apply Chaparral at 1.5 to 2.0 ounce/acre to plants at the rosette through bolting growth stages.

**Sulfur cinquefoil:** Apply Chaparral at 2.0 to 2.5 ounce/acre to plants in the prebud stage of development.

**Thistle, Canada:** Apply Chaparral at 2.0 to 3.3 ounce/acre either in the spring after all plants have fully emerged (some may be budding) or until the oldest plants are in full flower stage. Use the higher rate when

applying to the flower stage. Applications are also effective in the fall before a killing frost. Use higher rates for older/dense stands or for longer residual control.

**Thistles, Bull, musk, and plumeless:** Apply Chaparral at 1.0 to 2.0 ounce/acre in the spring and early summer to rosette or bolting plants or in the fall to seedlings and rosettes. Apply at 2.0 to 2.5 ounce/acre plus 0.5 lb ae/acre 2,4-D when plants are at the late bolt through early flowering growth stages.

**Vervain: Chaparral alone will provide suppression.** Apply 1.5 to 2.0 oz/acre of Chaparral with 0.5 lb ae/acre (1 pint/acre of 4 lb ae/gallon 2,4-D) of 2,4-D.

**Whitetop:** Apply 3.3 ounce/acre early in the spring to actively growing rosettes or to regrowth before the bud stage. Treatment after bloom is generally less effective and the addition of 2,4-D at 1 lb ae/acre (2 pint/acre of 4 lb ae/gallon 2,4-D) is recommended.

**Wormwood, absinth:** Apply 3.0 to 3.3 ounce/acre before wormwood is 12 inches tall. When applying by air on CRP, coverage is important and a minimum of 3 GPA is specified. Remove old duff and litter by fire or mowing for best results. Fall applications are also effective if green regrowth is present.

### Woody Plant Control:

#### FOLIAR APPLICATIONS:

For broad spectrum brush control using a foliar application, Chaparral may be added to tank mixes with Remedy Ultra or other herbicides labeled for rangeland to improve brush control on species such as aspen, conifers (pine), elm, maple, cherry, poplar, oak, Scotch broom, boxelder, hackberry, Russian olive, and salt cedar.

For mixed stands of woody species or shrubs such as buckbrush, absinth wormwood, fringe sage, silverberry, shrubby cinquefoil, wild rose, wild strawberry, and pasture sage Apply 2.5-3 oz of Chaparral with 1 lb ae/A or 2,4-D ester with 0.25% v/v surfactant. Use higher rate for dense stands or later in the application window

**Table 3: Woody Plant Control: Apply Chaparral at 3.3 ounce/acre at the timing described below**

Common Name	Scientific Name	Plant Family	Application Details
blackberry*	<i>Rubus spp</i>	Rosaceae	Apply when leaves are fully expanded and the plant has stopped rapid spring and early summer growth. Application after bloom and before frost is optimal. It is recommended that after mowing, shredding, or burning, applications should wait until the next season and enough re-growth has occurred for good uptake and translocation.
buckbrush (coralberry) or Western snowberry	<i>Symphoricarpos orbiculus</i> or <i>Symphoricarpos occidentalis</i>	Caprifoliaceae	Apply 3 oz/acre of Chaparral alone or 2.0 to 3.0 oz/acre with 1 lb ae/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) in the spring when leaves are fully expanded and foliage is mature. Apply 3 oz/acre with 1 lb ae/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) from full leaf expansion up to the flowering stage.
dewberry*	<i>Rubus flagellaris</i>	Rosaceae	Apply when leaves are fully expanded and the foliage is dark green, either before first flower or after fruit drop. Application after fruit drop is preferred until frost. It is recommended that after mowing, shredding, or burning applications should wait until the next season and enough re-growth has occurred for good uptake and translocation.
honey locust	<i>Gleditsia triacanthos</i>	Fabaceae	Apply in spring when leaves are fully expanded and foliage is mature.

Common Name	Scientific Name	Plant Family	Application Details
honeysuckle	<i>Lonicera japonica</i>	Caprifoliaceae	Apply in spring when leaves are fully expanded and foliage is mature.
kudzu	<i>Pueraria montana</i>	Fabaceae	Apply at or after bloom (July) in the summer until fall when the foliage begins to senesce. Kudzu should be actively growing; avoid treating when drought stressed.
locust, black	<i>Robinia pseudoacacia</i>	Fabaceae	Apply in spring when leaves are fully expanded and foliage is mature.
mimosa	<i>Albizia julibrissin</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
plum, American and sand hill	<i>Prunus Americana</i> and <i>Prunus angustifolia</i>	Rosaceae	Apply in spring when leaves are fully expanded and foliage is mature.
redbud	<i>Cercis canadensis</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
rose, Cherokee	<i>Rosa laevigata</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
rose, multiflora	<i>Rosa multiflora</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
rose, prairie wild	<i>Rosa arkansana</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
Tree of heaven	<i>Ailanthus altissima</i>	Simaroubaceae	Apply after full leaf out and before fall foliage color change
wisteria	<i>Wisteria brachybotrys</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
yucca‡	<i>Yucca glauca</i>	Agavaceae	Add 1 lb ai/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) to Chaparral at 3.3 ounce/acre. Another option for additional woody plant control is Chaparral plus 1 pint/acre Remedy® Ultra. Make applications from flower stalk elongation through seed pod development. Crop oil concentrate (COC), Methylated Seed Oil (MSO), or Methylated Seed Oil/ Organosilicone (MSO/OS) are the preferred adjuvants. Aerial application is recommended with a minimum of 4 gallons per acre volume for dense yucca populations.

‡ This symbol denotes weed suppression which is a reduction in weed competition compared to untreated areas. A second treatment may be necessary.

\* This recommendation is for blackberry and dewberry control in bermudagrass or other non-sensitive grasses only. For control in tall fescue pastures, only apply Chaparral as a spot treatment. For broadcast blackberry control in tall fescue pastures, use 1 pint/acre of Remedy Ultra + 2 pts/acre of ForeFront™ R&P.

## For Use on Grass Harvested for Hay Intended for Distribution or Sale Off the Farm or Ranch

## For Use on Grass Harvested for Silage, Haylage, Baleage, and Green Chop Intended for Use On the Farm or Ranch

For Use Only in the Following States: AL, AR, AZ, CO, FL, GA, ID, KS, KY, LA, MO, MS, MT, ND, NE, NV, NM, OK, SD, TN, TX, UT, WY

## Use Precautions and Restrictions

Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Use Restrictions. Call 800-258-3033 for more information.

<p style="text-align: center;"><b>IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS</b></p> <ul style="list-style-type: none"> <li>Carefully read the section Restrictions in Hay or Manure Use.</li> <li>It is mandatory to follow the Use Restrictions section of this label.</li> <li>Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.</li> <li>Hay can only be used on the farm or ranch where product is applied.</li> <li>Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.</li> <li>For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <a href="https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides">https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides</a></li> </ul>	<p style="text-align: center;"><b>Forage and Manure Management</b></p> <p style="text-align: center;">Warning: Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.</p>
--	---

- Do not use grasses treated with Chaparral in the preceding 18 months for hay intended for export outside the United States.
- Do not use hay or straw from areas treated with Chaparral within the preceding 18 months, or manure from animals feeding on hay treated with Chaparral, in compost.
- Do not use grasses treated within the preceding 18 months for seed production.
- Do not use on Timothy hay or other cool-season grasses grown for hay.
- Do not overseed ryegrass for 4 months after treatment.



- **Chaparral is highly active against many broadleaf plant species.** Do not use this product on areas where loss of broadleaf plants, including legumes, cannot be tolerated.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if aminopyralid or metsulfuron concentration remaining in the soil will adversely affect the legume establishment.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or grass hay harvest intervals following application of Chaparral at labeled rates. However, cutting hay too soon after spraying weeds will reduce weed control. Wait 14 days after herbicide application to cut grass hay to allow herbicide to work. Do not transfer grazing animals from areas treated with Chaparral to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- **Grazing Poisonous Plants:** Application of this product may increase palatability of certain poisonous plants. Do not graze areas treated with Chaparral until poisonous plants are dry and no longer palatable to livestock.
- **Restrictions in Hay or Manure Use:**
  - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18 months, in compost, mulch or mushroom spawn.
  - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch, or mushroom spawn.
  - Do not spread manure from animals that have grazed or consumed forage or eaten hay from treated areas within the previous 3 days on land used for growing susceptible broadleaf crops.
  - Manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days may only be used on pasture grasses, grass grown for seed, wheat, and corn.
  - Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields treated with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- **Crop Rotation:** Do not rotate to any crop from rangeland, permanent pasture, or CRP acres within one year following treatment. Cereals and corn can be planted one year after treatment. Most broadleaf crops are more sensitive and can require **at least 2 years** depending on the crop and environmental conditions. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid or metsulfuron present in the soil will not adversely affect that broadleaf crop.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern, or drainage. The field bioassay can be initiated one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, perennial forage grasses, native grasses, or grasses grown for hay.



---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

**Warranty Disclaimer**

---

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™®Trademarks of Corteva Agriscience and its affiliated companies

EPA accepted \_\_/\_\_/\_\_\_\_

**[Sub-Label B for IVM - Alternate Brand Name – Opensight Specialty Herbicide]**

(Base label):


AMINOPYRALID	GROUP	4	HERBICIDE
METSULFURON-METHYL	GROUP	2	HERBICIDE

**Opensight®****SPECIALTY HERBICIDE**

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), most warm-season grasses grown for hay, Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites.

**IMPORTANT USE PRECAUTIONS AND  
RESTRICTIONS TO PREVENT INJURY TO  
DESIRABLE PLANTS**

- Carefully read the section Restrictions in Hay or Manure Use.
- It is mandatory to follow the Use Restrictions section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied.
- Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.
- For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>

**Forage and Manure Management**

**Warning:** Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.

**Not For Sale, Distribution, or Use in New York State.**

**Active Ingredients:**

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro- .....	62.13%
Metsulfuron methyl (Methyl 2-[[[(4-methoxy-6- methyl-1,3,5- triazin-2-yl)-amino]carbonyl] amino]sulfonyl]benzoate) .....	9.45%
Other Ingredients .....	<u>28.42%</u>
Total .....	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

**Keep Out of Reach of Children****WARNING      AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

---

**Precautionary Statements**

---

**Hazard to Humans and Domestic Animals**

**Causes Substantial but Temporary Eye Injury • Harmful if Swallowed**

**Do not get in eyes or on clothing. Avoid contact with skin.**

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate or Viton ≥14 mils
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

## Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**Groundwater Advisory:** This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Users are advised not to apply this product where soils have a rapid to very rapid permeability (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching both surface water and aquatic sediment via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron-methyl and aminopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**Irrigation Water Statement:** Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

## Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the "Directions for Use" section for information about this standard.

## Nonrefillable rigid containers 5 gallons or less:

### Storage and Disposal

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Nonrefillable nonrigid containers:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable rigid containers larger than 5 gal:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable rigid containers larger than 5 gal:****Storage and Disposal**

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water.

Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times.

Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-597

EPA Est. \_\_\_\_\_  
[CD02-xxx-xxx]

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for**  
**Corteva Agriscience LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET WEIGHT**\_\_\_\_\_



(Booklet cover / base label for small containers):

<b>AMINOPYRALID</b>	<b>GROUP</b>	<b>4</b>	<b>HERBICIDE</b>
<b>METSULFURON-METHYL</b>	<b>GROUP</b>	<b>2</b>	<b>HERBICIDE</b>

# Opensight®


## SPECIALTY HERBICIDE

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), most warm-season grasses grown for hay, Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites.

### IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- Carefully read the section Restrictions in Hay or Manure Use.
- It is mandatory to follow the Use Restrictions section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- Hay can only be used on the farm or ranch where product is applied.
- Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.
- For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>

### Forage and Manure Management



**Warning:** Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.

**Not For Sale, Distribution, or Use in New York State.**

## Active Ingredients:

Potassium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro- .....	62.13%
Metsulfuron methyl (Methyl 2-[[[(4-methoxy-6- methyl-1,3,5- triazin-2-yl)-amino]carbonyl] amino]sulfonyl]benzoate) .....	9.45%
Other Ingredients .....	<u>28.42%</u>
Total .....	100.0%

Acid Equivalent: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) - 52.5%

Contains 0.62 pound potassium salt of aminopyralid active ingredient (0.525 pound acid equivalent) and 0.0945 pound metsulfuron methyl per pound of product

**Keep Out of Reach of Children****WARNING      AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the "Directions for Use" section for information about this standard.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-597

EPA Est. \_\_\_\_\_  
[CD02-xxx-xxx]

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET WEIGHT** \_\_\_\_\_

(Booklet page 1 through end):

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## WARNING

**Causes Substantial but Temporary Eye Injury • Harmful if Swallowed**

**Do not get in eyes or on clothing. Avoid contact with skin.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate or Viton  $\geq 14$  mils
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**Groundwater Advisory:** This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Users are advised not to apply this product where soils have a rapid to very rapid permeability (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching both surface water and aquatic sediment via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron-methyl and aminopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**Irrigation Water Statement:** Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

#### **Windblown Soil Particles Advisory**

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

---

### **Directions for Use**


---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **Not For Sale, Distribution, or Use in New York State.**

**Not for use on pastures in Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont. All other labeled uses are permitted in these states including grazed areas in and around approved use sites.**



Light grey = states where use in pastures is not permitted  
Dark grey = NY where the product is not registered

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and

exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate or Viton  $\geq 14$  mils
- Shoes plus socks

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Do not enter or allow people or pets to enter the treated area until sprays have dried.

### Storage and Disposal

Do not contaminate water, food, feed, or fertilizer by storage or disposal.

**Pesticide Storage:** Store in original container only. In case of spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable rigid containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows:

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### Nonrefillable nonrigid containers:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### Refillable rigid containers larger than 5 gal:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.



**Nonrefillable rigid containers larger than 5 gal:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Opensight® may be applied by aerial or ground equipment to control susceptible broadleaf weeds and certain woody plants, including invasive and noxious weeds on rangeland, permanent grass pastures (including grasses grown for hay\*), most warm-season grasses grown for hay, CRP acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads), non-irrigation ditch banks, natural areas (such as wildlife management areas, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites without injury to most grasses.

It is permissible to treat non-irrigation ditch banks, seasonally dry wetlands (such as flood plains, deltas, marshes, swamps, or bogs), and transitional areas between upland and lowland sites only when dry. Opensight can be used up to the water's edge. Do not apply directly to water and take precautions to minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent, or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize overspray to open water. Note: Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

## Weed Resistance Management

This product contains aminopyralid, a Group 4 synthetic auxin, and metsulfuron-methyl, a Group 2 acetolactate synthase (ALS). Appropriate resistance management strategies should be followed.

Development of plant populations resistant to the mode of action of aminopyralid is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications. There may be resistant weed biotypes to metsulfuron and adequate control of these species cannot be expected.

To delay herbicide resistance, take one or more of the following steps:

- Development of plant populations resistant to the mode of action of aminopyralid is usually not a problem on rangeland, permanent grass pastures, Conservation Reserve Program (CRP), or non-cropland sites since these sites receive infrequent pesticide applications. There may be resistant weed biotypes to metsulfuron and adequate control of these species cannot be expected.
- Similar looking biotypes of a given weed species occurring in a treated area may vary in their susceptibility to a herbicide. Application of a herbicide below its labeled rate may allow more tolerant weeds to survive and a shift to more tolerant biotypes within the treated area.
- Where identified, spreading of resistant weeds to other fields may be prevented by cleaning harvesting and tillage equipment before moving to other areas and by planting weed-free seed.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed

production in the affected area by an alternative herbicide from a different group or by a mechanical method such as mowing.

- Use tank mixtures with herbicides from a different group if such use is permitted. Where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- For further information or to report suspected resistance, contact your extension specialist or contact a Corteva Agriscience customer service representative at 800-992-5994.

## **Mandatory Spray Drift Management**

### **Aerial Applications**

- Do not release spray at a height greater than 10 feet above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to select a coarse or coarser droplet size (ANSI/ASABE S641).
- For all other applications, applicators are required to select a medium or coarser droplet size (ANSI/ASABE S641).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply during temperature inversions.

### **Ground Boom Applications**

- For applications on pastures and rangeland, do not release spray at a height greater than 4 feet above the ground. For all other uses, do not release spray at a height greater than 3 feet above the ground or crop canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ANSI/ASAE S572.3).
- For all other applications, applicators are required to use a medium or coarser droplet size (ANSI/ASAE S572.3).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

### **Boom-less Ground Sprayer Applications**

- Applicators are required to select a medium or coarser droplet size (ANSI/ASAE S572.3) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

## **Spray Drift Advisories**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

**Boom-less Ground Applications:** Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

**Handheld Technology Applications:** Take precautions to minimize spray drift.

**IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

**Controlling Droplet Size – Ground Boom**

- **Volume:** Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure:** Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle:** Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

**Controlling Droplet Size – Aircraft**

- **Adjust Nozzles:** Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

**BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

**RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

**WIND**

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift

## Use Restrictions

**Consult with a Corteva Agriscience representative if you do not understand the Use Restrictions. Call 800-258-3033 for more information.**

<p style="text-align: center;"><b>IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS</b></p> <ul style="list-style-type: none"> <li>Carefully read the section Restrictions in Hay or Manure Use.</li> <li>It is mandatory to follow the Use Restrictions section of this label.</li> <li>Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broadleaf plants.</li> <li>Hay can only be used on the farm or ranch where product is applied.</li> <li>Consult with a Corteva Agriscience representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 Customer Information Group.</li> <li>For more information on how to manage materials treated with aminopyralid and to prevent this product from contaminating compost please visit <a href="https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides">https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides</a></li> </ul>	<p style="text-align: center;"><b>Forage and Manure Management</b></p> <p style="text-align: center;"><b>Warning: Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.</b></p>
--	--

- Do not use grasses treated with Opensight in the preceding 18 months for hay intended for export outside the United States.**
- Do not use hay or straw from areas treated with Opensight within the preceding 18 months or manure from animals feeding on hay treated with Opensight in compost.**
- Do not use grasses treated with Opensight in the preceding 18 months for seed production.**

**Maximum Application Rate:** On all labeled use sites do not broadcast apply more than 3.3 ounce/acre of Opensight per year. The total amount of Opensight applied broadcast, as a re-treatment, and/or spot treatment cannot exceed 3.3 oz of product per acre per year. Spot treatments may be applied at an equivalent broadcast rate of up to 6.6 oz product of Opensight per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 3.3 ounce/acre of Opensight per annual growing season as a result of broadcast, spot, or repeat applications.

- Do not use on Timothy hay or other cool-season grasses grown for hay.
- Do not apply this product on lawns, turf, ornamental plantings, urban walkways, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
- Do not overseed ryegrass for 4 months after treatment.

- **Opensight is highly active against many broadleaf plant species.** Do not use this product on areas where loss of broadleaf plants, including legumes, cannot be tolerated.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Do not contaminate water intended for irrigation or domestic purposes.** Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not apply to irrigated land where the tailwater will be used to irrigate crops.
- Do not use this product for impregnation on dry fertilizer, unless specified in Corteva Agriscience state-specific product bulletin.
- Do not use Opensight in the following counties of Colorado: Alamosa, Conejos, Costilla, Rio Grande, and Saguache.
- **Trees** adjacent to or in a treated site can occasionally be affected by root uptake of Opensight. Do not apply Opensight within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.
  - Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots extend, or in locations where the product may be washed or moved into contact with their roots, as injury or loss of desirable trees or other plants may result.
- Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of Opensight. Injury to crops may result if treated soil and/or runoff water containing Opensight is washed, or moved onto land used to produce crops. Exposure to Opensight may injure or kill susceptible crops and other plants, such as grapes, soybeans, tobacco, sensitive ornamentals. Do not treat frozen soil where runoff could damage sensitive plants.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if aminopyralid or metsulfuron concentration remaining in the soil will adversely affect the legume establishment.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after Opensight application, temporary discoloration, and/or grass injury may occur. Opensight should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Do not apply to frozen ground as surface runoff may occur.
- Do not apply to snow-covered ground.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or grass hay harvest intervals following application of Opensight at labeled rates. However, cutting hay too soon after spraying weeds will reduce weed control. Wait 14 days after herbicide application to cut grass hay to allow herbicide to work.
- **Grazing Poisonous Plants:** Application of this product may increase palatability of certain poisonous plants. Do not allow livestock to graze areas treated with Opensight until poisonous plants are dry and no longer palatable to livestock.
- **Restrictions in Hay or Manure Use:**
  - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18 months, in compost, mulch, or mushroom spawn.
  - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch, or mushroom spawn.
  - Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields treated with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
  - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.



- To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid in plant residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.
- **Off-site Compost and Animal Feed/Bedding Restrictions**
  - This product is persistent and may be present in treated plant materials for months to years after application. Do not sell or transport treated plant materials or manure from animals that have grazed on treated plant materials off-site for compost distribution or for use as animal bedding/feed for 18 months after application. Treated plant materials can be recycled onsite or left in the field to decompose.
  - Manure from animals that have grazed or eaten forage or hay harvested from treated areas within the previous 3 days may only be applied to the fields where the following crops will be grown: pasture grasses, grass grown for seed, wheat, and corn.
  - Animals that have been fed aminopyralid-treated forage must be fed forage free of aminopyralid for at least 3 days before movement to an area where manure may be collected or sensitive crops are grown.
- **Crop Rotation:** Do not rotate to any crop from rangeland, permanent pasture, or CRP acres within one year following treatment. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid or metsulfuron present in the soil will not adversely affect that broadleaf crop.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern, or drainage. The field bioassay can be initiated one year after the last application of aminopyralid in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, perennial forage grasses, native grasses or grasses grown for hay.
- **Avoiding Injury to Non-Target Plants:** Do not aerially apply Opensight within 50 feet of a border downwind (in the direction of wind movement), or allow spray drift to come in contact with, any broadleaf crop or other desirable broadleaf plants, including, but not limited to, alfalfa, cotton, dry beans, flowers, grapes, lettuce, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes or other broadleaf or vegetable crop, fruit trees, ornamental plants, or soil where sensitive crops are growing or will be planted. Avoid application under conditions that may allow spray drift because very small quantities of spray may seriously injure susceptible crops. Read the Spray Drift Management section of this label to help minimize the potential for spray drift.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery dry or light sandy soils until they have been stabilized by rainfall, plant residue mulch, reduced tillage, or other cultural practices. Injury to immediately adjacent crops may occur when treated soil is blown onto land used to produce crops other than pasture, rangeland, or CRP.

#### **Record Keeping For Applications to Pastures**

- The applicator must document that they have notified property owners/operators or customers, in writing, of the compost and animal feed/bedding prohibitions within 14 days of the application. Applicators must keep the records of notification for two years. This record must include date of application, the name of the applicator, the EPA registration number of the product applied, area(s) treated, and a copy of the written notification provided to the property owner/operator. Notification may be made via email, via mail, via paper handout, or by any other written communication method. Records must be made available to State Pesticide Regulatory Official(s) and to EPA upon request. If this information is already being retained, duplicate records are not needed.
- It is recommended that applicators also transmit at the time of notification relevant educational materials for managing treated plant matter, as available. Additional educational materials for aminopyralid will be posted at: <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>
- Applications to pastures by property owners/operators on their own property are exempt from this notification and recordkeeping requirement.

- Application to pastures on public land (i.e., lands managed directly by state, tribal, or local authorities) are exempt from this notification requirement.

### **Sprayer Clean-Out Instructions**

**It is recommended to use separate spray equipment on highly sensitive crops such as tobacco, soybeans, peanuts, and tomatoes.**

Do not use spray equipment used to apply Opensight for other applications to land planted to, or to be planted to, broadleaf plants unless it has been determined that all residues of this herbicide has been removed by thorough cleaning of equipment.

Equipment used to apply Opensight should be thoroughly cleaned before reusing to apply any other chemicals as follows:

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
2. Rinse a second time, adding 1 quart of household ammonia or tank cleaning agent for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Spray nozzles and screens should be removed and cleaned separately.

Do not apply this product with mist blower systems that deliver very fine spray droplets. Use of mist blower equipment can reduce control achieved with the herbicide and increase spray drift potential.

## **Application Methods**

Apply the specified rate of Opensight as a coarse low-pressure spray. Do not apply this product with mist blower systems that deliver very fine spray droplets. Spray volume should be sufficient to uniformly cover foliage. Increase the spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. To enhance foliage wetting and coverage, an approved non-ionic agricultural surfactant may be added to the spray mixture as specified by the surfactant label.

**Ground Broadcast Application:** Higher spray volumes (greater than 10 gallons per acre) generally provide better coverage and better control, particularly in dense and/or tall foliage.

**Aerial Broadcast Application:** Do not apply less than 2 gallons per acre total spray volume. Five gallons per acre or greater will generally provide better coverage and better control, particularly in dense and/or tall foliage.

**High-Volume Foliar Application:** High volume foliar treatments may be applied at rates equivalent to a maximum of 3.3 ounces per acre annual growing season. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems.

**Spot Application:** Spot treatments may be applied at an equivalent broadcast rate of up to 6.6 oz of product per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate. Do not apply more than a total of 3.3 ounce/acre of Opensight per annual growing season as a result of broadcast, spot, or repeat applications. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage, but not to the point of runoff. Repeat treatments may be made, but the total amount of Opensight applied must not exceed 3.3 ounce/acre per year. Take precautions to minimize spray drift. To prevent misapplication, spot treatments should be applied with a calibrated sprayer.

In general for spot treatments, mix 2.5 oz for weeds and 3.3 oz for brush of Opensight per 100 gallons of water (assuming an application volume of 100 gallons per acre).

### **Product Measurement**

Opensight is measured using the Opensight volumetric measuring cylinder. Scales calibrated in ounces may also be used.

**Mixing Instructions**

1. Fill the tank 1/4 to 1/3 full with water. (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures sections for additional details.)
2. While agitating, add the required amount of Opensight.
3. Continue agitation until the Opensight is fully dispersed, at least 5 minutes.
4. Once the Opensight is fully dispersed, maintain agitation and continue filling tank with water. Opensight should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of spray adjuvants. Always add spray adjuvants last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply Opensight spray mixture within 24 hours of mixing to avoid product degradation.
8. If Opensight and a tank mix partner are to be applied in multiple loads, pre-slurry the Opensight in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Opensight.

**Soil pH Limitations**

Opensight should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Opensight could remain in the soil for 34 months or more injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of Opensight.

**Checking Soil pH**

Before using Opensight, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil-sampling procedures.

**Spray Adjuvants**

Unless otherwise directed, applications of Opensight must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer can be used unless specifically prohibited by tank mix partner labeling. If another herbicide is tank mixed with Opensight, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

**Petroleum Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO)**

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- MSO adjuvants may be used at 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

**Nonionic Surfactant (NIS)**

- Apply at 0.25% v/v (1 quart per 100 gallons spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

**Ammonium Nitrogen Fertilizer**

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 pounds/acre of a spray grade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.

**Special Adjuvant Types**

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO, and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions. Exception: On tall fescue pastures use a reduced rate of 1/2 to 1 pint non-ionic surfactant per 100 gallons.

- Antifoaming agents may be used if needed.
- Do not use Opensight with spray additives that reduce the pH of the spray solution to below 3.0.

**Tank Mixing with Other Herbicides:** Opensight at rates of up to 3.3 ounce/acre may be mixed with labeled rates of other herbicides registered for application on all labeled use sites. Opensight may be applied in tank-mix combination with labeled rates of other herbicides provided: (1) the tank-mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products, and (3) that the tank-mix combination is physically compatible (see tank-mix compatibility testing below). When tank mixing, use only in accordance with the restrictions, precautions, and limitations on the respective product labels.

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not exceed specified application rates. If products containing the same active ingredient are mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

**Tank Mix Compatibility Testing:** Perform a jar test prior to mixing in a spray tank to ensure compatibility of Opensight and other pesticides or carriers. Use a clear glass jar with lid and mix ingredients in the same order and proportions as will be used in the spray tank. The mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily remix if agitated. An incompatible mixture is indicated by separation into distinct layers that do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film in the jar. Use of an appropriate compatibility aid may resolve mix incompatibility. If the mixture is incompatible do not use that tank mix partner in tank mixtures.

**Note:** Foliar-applied liquid fertilizers themselves can cause yellowing of the foliage of forage grasses and other vegetation.

## Guidelines for Grass Management

Opensight may be applied to established native grasses such as wheatgrasses, bluestems and grama, and on other established pasture grasses such as bermudagrass, bluegrass, orchardgrass, brome grass, and tall fescue that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

- Opensight may suppress certain established grasses, such as smooth brome grass (*Bromus inermis*), especially when plants are stressed by adverse environmental conditions. Plants should recover from this transient suppression with the onset of environmental conditions favorable to grass growth and upon release from weed competition.
- Varieties and species of forage grasses differ in their tolerance to herbicides. When using Opensight on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated.
- Application of Opensight to Pensacola bahiagrass, ryegrass (Italian or perennial), and Garrison's creeping foxtail may cause severe injury to and/or loss of pastures.

## Seeding grasses:

**Preemergence:** In general, Opensight may be applied in the spring or early summer, depending on the target weed species, as a broadcast application over grass that was planted at least 4 months prior to the application and that has been growing under favorable conditions for grass establishment.

With fall applications, do not plant grasses the following spring.

Do not overseed ryegrass for 4 months after treatment.

**Tall Fescue:**

Opensight may stunt tall fescue, cause it to turn yellow, or cause seed head suppression. To minimize these symptoms, take the following precautions.

- Do not use on tall fescue grown for seed.
- Do not use more than 2 ounce/acre of Opensight.
- Tank-mix Opensight with 2,4-D.
- Use a reduced rate of non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution (1/16 to 1/8% v/v).
- Make application later in the spring after the new growth is 5 to 6 inches tall (until after reproductive culm has started to elongate), or in the fall.
- Do not use surfactant when liquid nitrogen is used as a carrier.
- Do not use a spray adjuvant other than non-ionic surfactant.

Initial grass yields may be reduced due to fescue seed head suppression resulting from treatment with Opensight at labeled rates. However, this could be beneficial because in tall fescue infected with the fungal endophyte (*Neotyphodium* spp.), the endophyte is concentrated in the seed, and cattle grazing plants with the seed head will get the maximum exposure to the endophyte. Increased levels of ingestions of the fungal endophyte can reduce weight gain and conception rates in cattle. Since the first grazing is often delayed in the spring until long after seed head development, Opensight could potentially be used to reduce development of the seed head, thereby reducing the amount of the endophyte that would be consumed by livestock when grazing.

**Seed Head Suppression:** If the intent is to control weeds and reduce tall fescue seed heads, apply Opensight at 2.0 to 2.5 ounce/acre early to fescue that is less than 6 inches tall.

**Pensacola bahiagrass control in established Bermudagrass pasture:**

Apply Opensight at 2-2.5 ounce/acre after green-up in the spring, but before bahiagrass seed head formation. Application should be made when environmental conditions favor grass growth.

Bahiagrass suppression could take up to 30 days before the desired level of control is achieved. Application of 2,4-D mixed with Opensight could decrease bahiagrass control. In pastures severely infested with bahiagrass, a positive response in forage yield may be slowed until desired forage grasses, like bermudagrass, grow into areas previously infested with bahiagrass. To reduce this effect, consider treating different portions of heavily infested pastures with Opensight over a period of several years. Do not apply Opensight to an entire farm or ranch in one year. Fertilization and/or replanting may accelerate bermudagrass recovery following bahiagrass control with Opensight.

Bahiagrass regrowth may occur in pastures heavily infested with bahiagrass, intense grazing pressure, or when adverse environmental conditions (heat and drought), slows the recovery of desired grass forages.

Opensight will not control common or Argentine bahiagrass.

Pensacola bahiagrass control can be reduced when Opensight is applied in liquid fertilizer solutions.

**Use Rates and Timing**

Opensight may be applied post emergence as a broadcast spray or as a spot application to control weeds and brush including, but not limited to, those listed on this label. When a rate range is given use the higher rate to control weeds at advanced growth stages, or under less than favorable growing conditions, or for longer residual control. Best results are obtained when spray volume is sufficient to provide uniform coverage of treated weeds. For optimum uptake and translocation of Opensight, avoid mowing, haying, shredding, burning, or soil disturbance in treated areas for at least 14 days following application.

Opensight also provides preemergence control of emerging seedlings of susceptible weeds, and re-growth of certain perennial weeds following application. Preventing establishment of weeds will depend upon application rate, season of application, and environmental conditions after application.



Opensight can provide long-term control of susceptible weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Long-term weed control is most effective where grass vegetation is allowed to recover from overgrazing, drought, etc., and compete with weeds.

Opensight can be an important component of integrated vegetation management programs designed to renovate or restore desired plant communities. To maximize and extend the benefits of weed control provided by Opensight, it is important that other vegetation management practices, including proper grazing management, biological control agents, replanting, fertilization, prescribed fire, etc., be used in appropriate sequences and combinations to further alleviate the adverse effects of weeds on desirable plant species and to promote development of desired plant communities. Agricultural and natural resources specialists with federal and state government agencies can provide guidance on best management practices and development of integrated vegetation management programs.

## Species Controlled

**General Mix of Broadleaf Weeds:** Opensight at 2.0 ounce/acre is the standard rate to provide control of many problem weeds when applied early in the season. If certain weeds are key targets, use the rate in Table 1 for that species. The addition of Garlon herbicides, DMA 4 IVM, or other herbicides allowed for use on the site to be sprayed can be tank mixed to broaden the weed spectrum.

Opensight controls weeds and woody plants primarily by postemergent activity. Although Opensight has preemergence activity, best results are generally obtained when Opensight is applied to foliage after emergence or dormancy break. Generally, for the control of annual weeds, Opensight provides the best results when applied to young, actively growing weeds. For the control of perennial weeds, applications made at the bud/bloom stage or while the target weeds are in the fall rosette stage typically provide the best results. The use rate depends upon the weed species and size of the weed at the time of application.

The degree and duration of control depends on weed spectrum and infestation intensity, weed size at application, environmental conditions at and following treatment, soil pH, soil moisture, and soil organic matter, and other factors.

For best results, most weeds should be treated when they are actively growing and under conditions favorable for growth. Use a higher rate in the rate range indicated when growing conditions are less than favorable (drought conditions), weeds are large and mature, weed density and foliage cover is high and canopy height is tall, or when residual control is desired. Opensight also provides preemergence control of germinating seeds or seedlings of susceptible weeds following application.

For rates for specific weeds, see Table 1. The life cycle is included for each weed species. The general timing of application for each life cycle is as follows:

**Annuals:** Use lower rates when weeds are less than 6 inches and actively growing. Increase rate as season progresses and plants become more mature.

**Biennials:** Apply in the spring and early summer to rosette or bolting plants or in the fall to seedlings and rosettes before ground is frozen. Use higher rates after bolting through early flower.

**Perennials:** Apply to vegetative stage prior to bloom. Use higher rate when weeds are larger.

**Table 1: Species Controlled with Opensight**

**Note:** Weeds marked with \* indicate more information is included in the specific weed problems section after the table.

**Broadleaf Weeds Controlled by Opensight**

Common Name	Scientific Name	Life Cycle	Plant Family	Opensight rate oz/a
actinomeris, wingstem	<i>Verbesina alternifolia</i>	perennial	Asteraceae	3.0
alyssum, hoary	<i>Berteroa incana</i>	biennial	Brassicaceae	2.0-2.5
amaranth, spiny	<i>Amaranthus spinosus</i>	summer annual	Amaranthaceae	1.5-2.0
arrowgrass, seaside ‡	<i>Trifglochin maritima</i>	perennial	Juncaginaceae	3.0-3.3
aster	<i>Aster spp.</i>	perennial	Asteraceae	1.5-2.0
bahiagrass, Pensacola*	<i>Paspalum notatum</i> <i>Flugge</i>	perennial	Poaceae	2.0-2.5
babysbreath	<i>Gypsophila paniculata</i>	perennial	Caryophyllaceae	2.5-3.0
bedstraw	<i>Galium spp.</i>	perennial	Rubiaceae	2.0-2.5
bittercress	<i>Cardimane spp</i>	perennial	Brassicaceae	2.0-2.5
blackeyed-susan	<i>Rudbeckia hirta</i>	annual	Asteraceae	1.5-2.0
brackenfern	<i>Pteridiums spp.</i>	perennial	Dennstaedtiaceae	2.5-3.3
broomweed, annual	<i>Amphiachyris dracunculoides</i>	annual	Asteraceae	1.0-1.5
bur buttercup (testiculate)	<i>Ranunculus testiculatus</i>	annual	Ranunculaceae	1.0-1.5
burclover	<i>Medicago spp</i>	annual	Fabaceae	1.5-2.0
burdock, common	<i>Arctium minus</i>	biennial	Asteraceae	2.0-2.5
buttercup, hairy	<i>Ranunculus sardous</i>	perennial	Ranunculaceae	1.0-1.5
buttercup, tall	<i>Ranunculus acris</i>	perennial	Ranunculaceae	2.0-2.5
camelthorn	<i>Alhagi pseudalhagi</i>	perennial	Fabaceae	2.0-3.0
camphorweed	<i>Heterotheca subaxillaris</i>	summer annual	Asteraceae	2.0-3.0
campion, bladder ‡	<i>Silene vulgaris</i>	perennial	Caryophyllaceae	2.0-2.5
caraway, wild	<i>Carum carvi</i>	biennial	Apiaceae	2.5-3.0
carrot, wild	<i>Daucus carota</i>	biennial	Apiaceae	2.0-2.5
catchfly, conical	<i>Silene conoidea</i>	annual	Caryophyllaceae	1.0-1.5
chamomile	<i>Matricaria spp</i>	annual	Asteraceae	2.5-3.0
chickweed, common	<i>Stellaria media</i>	winter annual	Caryophyllaceae	3.0
chicory	<i>Cichorium intybus</i>	perennial	Asteraceae	1.5-2.0
cinquefoil*	<i>Potentilla spp</i>	perennial	Rosaceae	2.0-2.5
clover, sweet	<i>Melilotus officinalis</i>	biennial	Fabaceae	2.5-3.0
clover, white	<i>Trifolium repens</i>	perennial	Fabaceae	1.5-2.0
cockle, corn	<i>Agrostemma githago</i>	annual	Caryophyllaceae	2.0-3.0
cocklebur	<i>Xanthium strumarium</i>	annual	Asteraceae	1.5-2.0
coreopsis, plains	<i>Coreopsis tinctoria</i>	annual	Asteraceae	2.0-3.0
cowcockle	<i>Vaccaria pyramidata</i>	annual	Caryophyllaceae	1.5-2.0
crazyweed, silky	<i>Oxytropis Lambertii</i>	perennial	Fabaceae	2.0-2.5
croton, woolly	<i>Croton capitatus</i>	annual	Euphorbiaceae	1.5-2.0
crownvetch	<i>Securigera varia</i>	perennial	Fabaceae	1.5-2.0

Common Name	Scientific Name	Life Cycle	Plant Family	Opensight rate oz/a
crupina, common	<i>Crupina vulgaris</i>	perennial	Asteraceae	3.0-3.3
cudweed, purple	<i>Gnaphalium purpureum</i>	annual	Asteraceae	2.0-2.5
daisy, oxeye*	<i>Leucanthemum vulgare</i>	perennial	Asteraceae	2.5-3.3
dandelion, common	<i>Taraxacum officinale</i>	perennial	Asteraceae	1.5-2.0
dock	<i>Rumex spp</i>	perennial	Polygonaceae	2.0-2.5
dyer's woad ‡	<i>Istis tinctoria</i>	perennial	Brassicaceae	3.3
evening primrose, cutleaf	<i>Oenothera laciniata</i>	annual	Asteraceae	1.5-2.0
false dandelion, Carolina	<i>Tragopogon dubius</i>	biennial	Asteraceae	1.5-2.0
falseflax, smallseed	<i>Camelina microcarpa</i>	annual/ biennial	Brassicaceae	1.5-2.0
fiddleneck, common	<i>Amsinckia intermedia</i>	annual	Boraginaceae	1.5-2.0
filaree, redstem	<i>Erodium cicutarium</i>	annual/ biennial	Geraniaceae	3.0-3.3
fireweed	<i>Epilobium angustifolium</i>	perennial	Onagraceae	2.5-3.0
fleabane, annual	<i>Erigeron annuus</i>	annual	Asteraceae	1.5-2.0
garlic, wild	<i>Allium vineale</i>	perennial	Liliaceae	1.5-2.0
geranium, Carolina	<i>Geranium carolinianum</i>	winter annual	Geraniaceae	1.5-2.0
goldenrod spp	<i>Solidago canadensis</i>	perennial	Asteraceae	2.0-2.5
gumweed, curlycup	<i>Grindelia squarrosa</i>	biennial	Asteraceae	2.0-2.5
halogeton	<i>Halogeton glomeratus</i>	annual	Chenopodiaceae	3.0-3.3
hawkweed, orange*	<i>Hieracium aurantiacum</i>	perennial	Asteraceae	2.5-3.3
hawkweed, yellow*	<i>Hieracium pratense</i>	perennial	Asteraceae	2.5-3.3
hemlock, poison ‡	<i>Conium maculatum</i>	perennial	Apiaceae	2.5-3.3
henbane, black	<i>Hyoscyamus niger</i>	annual/ biennial	Solanaceae	2.5-3.0
henbit	<i>Lamium amplexicaule</i>	annual/ biennial	Lamiaceae	2.0-2.5
horsemint (beebalm)	<i>Monarda spp</i>	annual	Lamiaceae	1.5-2.0
horsenettle, Carolina	<i>Solanum carolinense</i>	perennial	Solanaceae	2.0-2.5
horseweed (maretail)	<i>Conyza canadensis</i>	annual	Asteraceae	1.5-2.0
houndstongue*	<i>Cynoglossum officinale</i>	biennial	Boraginaceae	2.5-3.3
ironweed, tall	<i>Vernonia gigantea</i>	perennial	Asteraceae	2.0-3.0
ironweed, western	<i>Vernonia baldwinii</i>	perennial	Asteraceae	2.0-3.0
knapweed	<i>Centaurea sp.</i>	biennial	Asteraceae	2.5-3.3
knapweed, brown	<i>Centaurea jacea</i>	perennial	Asteraceae	2.5-3.3
knapweed, diffuse*	<i>Centaurea diffusa</i>	biennial	Asteraceae	2.5-3.3
knapweed, Russian*	<i>Acroptilon repens</i>	perennial	Asteraceae	2.5-3.3
knapweed, spotted*	<i>Centaurea stoebe</i>	biennial	Asteraceae	2.5-3.3
knotweed, prostrate	<i>Polygonum aviculare</i>	annual	Polygonaceae	3.0
kochia*	<i>Kochia scoparia</i>	annual	Chenopodiaceae	1.5-2.0
lady's thumb	<i>Polygonum persicaria</i>	annual	Polygonaceae	1.5-2.0
lambsquarters, common	<i>Chenopodium album</i>	annual	Chenopodiaceae	2.0-2.5
lespedeza, annual	<i>Lespedeza striata</i>	annual	Fabaceae	2.0-2.5

Common Name	Scientific Name	Life Cycle	Plant Family	Opensight rate oz/a
lespedeza, sericea*	<i>Lespedeza cuneata</i>	perennial	Fabaceae	2.5-3.0
lettuce, Miner's	<i>Montia perfoliata</i>	annual	Portulacaceae	1.5-2.0
lettuce, prickly*	<i>Lactuca serriola</i>	annual	Asteraceae	1.5-2.0
locoweed	<i>Astragalus spp.</i>	perennial	Fabaceae	2.0-2.5
loosestrife, purple	<i>Lythrum salicaria</i>	perennial	Lythraceae	3.0-3.3
marshelder, annual ‡	<i>Iva annua</i>	annual	Asteraceae	2.0-2.5
mayweed, scentless	<i>Tripleurospermum perforata</i>	annual	Asteraceae	1.5-2.0
mayweed, stinking	<i>Anthemis cotula</i>	annual	Asteraceae	3.0-3.3
medic, black	<i>Medicago lupulina</i>	perennial	Fabaceae	2.0-2.5
Mexican-tea	<i>Dysphania ambrosioides</i>	annual/ perennial	Chenopodiaceae	2.0-2.5
mullein*	<i>Verbascum spp.</i>	biennial	Scrophulariaceae	2.0-3.3
mustard, blue*	<i>Chorispora tenella</i>	annual	Brassicaceae	1.5-2.0
mustard, tumble/Jim Hill	<i>Sisymbrium altissimum</i>	winter annual	Brassicaceae	1.5-2.0
mustard, wild	<i>Brassica kaber</i>	annual	Brassicaceae	1.5-2.0
needles, Spanish needles	<i>Bidens bipinnata</i>	annual	Asteraceae	2.0-2.5
oxtongue, bristly	<i>Picris echioides</i>	biennial	Asteraceae	2.5-3.0
parsnip, Wild	<i>Pastinaca sativa</i>	biennial	Apiaceae	2.0-3.0
partridgepea	<i>Chamaecrista fasciculata</i>	annual	Fabaceae	2.5-3.0
pepperweed, perennial‡*	<i>Lepidium latifolium</i>	perennial	Brassicaceae	3.3
pigweeds	<i>Amaranthus spp</i>	annual	Amaranthaceae	1.5-2.0
plantain, broadleaf	<i>Plantago major</i>	perennial	Plantaginaceae	2.0-2.5
plantain, buckhorn	<i>Plantago lanceolata</i>	perennial	Plantaginaceae	2.0-2.5
purslane, common	<i>Portulaca oleracea</i>	annual	Portulacaceae	1.5-2.0
ragweed, common	<i>Ambrosia artemisiifolia</i>	annual	Asteraceae	2.0-2.5
ragweed, western*	<i>Ambrosia psilostachya</i>	perennial	Asteraceae	2.0-2.5
ragwort, tansy	<i>Senecio jacobaea</i>	perennial	Asteraceae	2.5-3.0
rush skeletonweed	<i>Chondrilla juncea</i>	perennial	Asteraceae	2.5-3.0
salsify, Western ‡	<i>Tragopogon dubius</i>	biennial	Asteraceae	3.0-3.3
scouringrush ‡	<i>Equisetum hyemale</i>	grass	Equisetaceae	3.3
shephardspurse	<i>Capsella bursa-pastoris</i>	winter annual	Brassicaceae	1.5-2.0
sicklepod	<i>Senna obtusifolia</i>	annual	Fabaceae	2.5-3.0
sida, arrowleaf	<i>Sida rhombifolia</i>	annual	Malvaceae	2-2.5
smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	annual	Polygonaceae	1.5-2.0
snakeweed, broom*	<i>Gutierrezia sarothrae</i>	perennial	Asteraceae	3.0
sneezeweed, bitter	<i>Helenium amarum</i>	annual	Asteraceae	1.0-1.5
snow-on-the-mountain	<i>Euphorbia marginata</i>	annual	Euphorbiaceae	2.0-2.5
soda apple, tropical*	<i>Solanum viarum</i>	perennial	Solanaceae	2.5-3.0
sorrel, red	<i>Rumex acetosella</i>	perennial	Polygonaceae	2.0-2.5
sowthistle, perennial	<i>Sonchus arvensis</i>	perennial	Asteraceae	2.0-2.5
sowthistle, prickly	<i>Sonchus asper</i>	annual	Asteraceae	1.5-2.0

Common Name	Scientific Name	Life Cycle	Plant Family	Opensight rate oz/a
St. Johnswort, common	<i>Hypericum perforatum</i>	perennial	Clusiaceae	2.5-3.0
starthistle, purple*	<i>Centaurea calcitrapa</i>	biennial	Asteraceae	1.5-2.0
star-thistle, Malta*	<i>Centaurea melitensis</i>	annual	Asteraceae	1.5-2.0
starthistle, yellow*	<i>Centaurea solstitialis</i>	annual	Asteraceae	1.5-2.0
sunflower, common	<i>Helianthus annua</i>	annual	Asteraceae	1.5-2.0
tansy, common	<i>Tanacetum vulgare</i>	perennial	Asteraceae	2.5-3.3
teasel	<i>Dipsacus spp.</i>	biennial	Dipsacaceae	2.0-3.0
thistle, Russian*	<i>Salsola iberica</i>	annual	Chenopodiaceae	1.5-2.0
thistle, artichoke	<i>Cynara cardunculus</i>	perennial	Asteracea	2.0-3.0
thistle, bull*	<i>Cirsium vulgare</i>	biennial	Asteraceae	1.0-2.5
thistle, Canada*	<i>Cirsium arvense</i>	perennial	Asteraceae	2.0-3.3
thistle, Italian	<i>Carduus pycnocephalus</i>	annual	Asteraceae	2.0-3.0
thistle, musk*	<i>Carduus nutans</i>	biennial	Asteraceae	1.0-2.5
thistle, plumeless*	<i>Carduus acanthoides</i>	biennial	Asteraceae	1.0-2.5
thistle, Scotch	<i>Onopordum acanthium</i>	biennial	Asteraceae	1.5-2.5
thistle, woolly distaff	<i>Carthamus lanatus</i>	annual	Asteraceae	1.5-2.0
vervains ‡	<i>Verbena spp.</i>	perennial	Asteraceae	2.0-2.5
vetch, common*	<i>Vicia sativa</i>	annual	Fabaceae	1.5-2.0
wallflower, bushy	<i>Erysimum repandum</i>	annual	Brassicaceae	1.5-2.0
waterpod	<i>Ellisia nyctelea</i>	annual	Brassicaceae	1.5-2.0
whitetop (hoary cress)*	<i>Cardaria draba</i>	perennial	Brassicaceae	3.3
woodsorrel, yellow	<i>Oxalis stricta</i>	perennial	Oxalidaceae	3.0-3.3
wormwood, absinth*	<i>Artemisia absinthium</i>	perennial	Asteraceae	3.0-3.3
yankeeweed	<i>Eupatorium compositifolium</i>	perennial	Asteraceae	3.0-3.3
yarrow, common	<i>Achillea millefolium</i>	perennial	Asteraceae	1.5-2.0

‡ This symbol denotes weed suppression which is a reduction in weed competition compared to untreated areas. A second treatment may be necessary. The addition of 0.5 lbs ae/acre of 2,4-D may improve initial control.

**Hawkweed, orange or yellow:** Apply Opensight at 2.5 to 3.3 ounce/acre to plants in the bolting stage of development.

**Houndstongue:** Apply 2.5 ounce/acre to rosettes. As plant bolts, increase the rate to 3.0 to 3.3 ounce/acre up to early bud stage. Add 1 quart of 2,4-D/acre after the bud stage.

**Knapweeds, diffuse and spotted:** Apply Opensight at 2.5 to 3.3 ounce/acre when plants are actively growing with the optimum time of application occurring from rosette to the bolting stages of development or in the fall. Plants will be controlled by mid-summer and fall applications even though plants may not show any changes in form or stature the year of application.

**Knapweed, Russian:** Apply Opensight at 2.5 to 3.3 ounce/acre to plants in the spring and summer to plants from early bud to flowering stage and to dormant plants in the fall.

**Lespedeza, Sericea:** Apply 2.5 to 3.0 ounce/acre beginning at flower bud initiation through the full bloom stage of growth.

**Mullein:** Apply 2.0 ounce/acre in the rosette stage in spring or fall. Use rates from 2.5 to 3.3 ounce/acre for bolting plants less than 12 inches tall.

**Oxeye daisy:** Apply Opensight at 2.5 to 3.3 ounce/acre to plants in the prebud stage of development.

**Pepperweed, perennial:** Apply Opensight at 3.3 ounce/acre plus 2 lb ae/a 2, 4-D when plants are at early flowering through bloom for optimum control.



**Ragweed, Western:** Apply Opensight at 2.0 to 2.5 ounce/acre when plants are in the vegetative growth stage. The addition of 0.5 to 1 lb ae/acre (1 to 2 pints/acre of 4 lb ae/gallon 2,4-D) of 2,4-D/acre will improve control in dense stands or when ragweed is greater than 6 inches.

**Russian thistle, kochia, and prickly lettuce:** Naturally occurring resistant biotypes of these weeds to metsulfuron are known to occur. For best results, use Opensight at 1.5 to 2.0 ounces/acre in tank-mix with 2,4-D. Applications to these weeds should be made early to weeds less than 6 inches in height.

**Snakeweed, broom:** Applications should be made in the fall at 3.0 ounces/acre. Spring applications will provide suppression only.

**Soda apple, tropical:** Apply Opensight at 2.5 to 3.0 ounce/acre at any growth stage, but application by flowering will reduce seed production potential.

**Starthistle, malta, purple, and yellow:** Apply Opensight at 1.5 to 2.0 ounce/acre to plants at the rosette through bolting growth stages.

**Sulfur cinquefoil:** Apply Opensight at 2.0 to 2.5 ounce/acre to plants in the prebud stage of development.

**Thistle, Canada:** Apply Opensight at 2.0 to 3.3 ounce/acre either in the spring or summer to fully emerged Canada thistle. The goal is to insure all plants have emerged and many of the thistles will be in the bud to early flower stage at this time. Applications are also effective in the fall before a killing frost. Use higher rates for older/dense stands or for longer residual control.

**Thistles, Bull, musk, and plumeless:** Apply Opensight at 1.0 to 2.0 ounce/acre in the spring and early summer to rosette or bolting plants or in the fall to seedlings and rosettes. Apply at 2.0 to 2.5 ounce/acre plus 0.5 lb ae/acre 2,4-D when plants are at the late bolt through early flowering growth stages.

**Vervain:** Apply 1.5 to 2.0 oz/acre of Opensight with 0.5 lb ae/acre (1 pint/acre of 4 lb ae/gallon 2,4-D) of 2,4-D.

**Whitetop:** Apply 3.3 ounce/acre early in the spring to actively growing rosettes or to regrowth before the bud stage. Treatment after bloom is generally less effective and the addition of 2,4-D at 1 lb ae/acre (2 pint/acre of 4 lb ae/gallon 2,4-D) is recommended. Treatments can also be made to fall regrowth before the first killing frost.

**Wormwood, absinth:** Apply 3.0 to 3.3 ounce/acre before wormwood is 12 inches tall. When applying by air on CRP, coverage is important and a minimum of 3 GPA is specified. Remove old duff and litter by fire or mowing for best results. Fall applications are also effective if green regrowth is present.

#### Woody Plant Control:

Apply Opensight at 3.3 ounce/acre at the timing described below in Table 2.

**Table 2: Woody Plant Control with Opensight**

Common Name	Scientific Name	Plant Family	Application Details
blackberry*	<i>Rubus spp</i>	Rosaceae	Apply when leaves are fully expanded and the plant has stopped rapid spring and early summer growth. Application after bloom and before frost is optimal. It is recommended that after mowing, shredding, or burning, applications should wait until the next season and enough re-growth has occurred for good uptake and translocation.
buckbrush	<i>Symphoricarpos orbiculus</i>	Caprifoliaceae	Apply 2.0 to 3.0 oz/acre in spring or early summer when new growth is 6-12 inches tall. Add 0.5 to 1 lb ae/acre of 2,4-D (1 to 2 pints/acre of 4 lb ae/gallon 2,4-D) to the lower rate.

Common Name	Scientific Name	Plant Family	Application Details
dewberry*	<i>Rubus flagellaris</i>	Rosaceae	Apply when leaves are fully expanded and the foliage is dark green, either before first flower or after fruit drop. Application after fruit drop is preferred until frost. It is recommended that after mowing, shredding, or burning, applications should wait until the next season and enough re-growth has occurred for good uptake and translocation.
honey locust	<i>Gleditsia triacanthos</i>	Fabaceae	Apply in spring when leaves are fully expanded and foliage is mature.
honeysuckle	<i>Lonicera japonica</i>	Caprifoliaceae	Apply in spring when leaves are fully expanded and foliage is mature.
kudzu	<i>Pueraria montana</i>	Fabaceae	Apply at or after bloom (July) in the summer until fall when the foliage begins to senesce. Kudzu should be actively growing; avoid treating when drought stressed.
locust, black	<i>Robinia pseudoacacia</i>	Fabaceae	Apply in spring when leaves are fully expanded and foliage is mature.
mimosa	<i>Albizia julibrissin</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
redbud	<i>Cercis canadensis</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
rose, Cherokee	<i>Rosa laevigata</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
rose, multiflora	<i>Rosa multiflora</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
rose, prairie wild	<i>Rosa arkansana</i>	Rosaceae	Apply from full leaf through flowering. For best results, delay treatment for 9-12 months after mowing.
snowberry, Western	<i>Symphoricarpos occidentalis</i>	Caprifoliaceae	Apply 3 oz/acre of Opensight alone or 2.0 to 3.0 oz/acre with 1 lb ae/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) in the spring when leaves are fully expanded and foliage is mature. Apply 3 oz/acre with 1 lb ae/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) from full leaf expansion up to the flowering stage.
wisteria	<i>Wisteria brachybotrys</i>	Fabaceae	Apply after full leaf emergence in the spring until fall foliage color change.
yucca ‡	<i>Yucca glauca</i>	Agavaceae	Add 1 lb ai/acre of 2,4-D ester (2 pints/acre of 4 lb ae/gallon 2,4-D) to Opensight at 3.3 ounce/acre. Another option for additional woody plant control is Opensight plus 1 pint/acre Remedy® Ultra. Make applications from flower stalk elongation through seed pod development. Crop oil concentrate (COC), Methylated Seed Oil (MSO) or Methylated Seed Oil/ Organosilicone (MSO/OS) are the preferred adjuvants. Aerial application is recommended with a minimum of 4 gallons per acre volume for dense yucca populations.

‡ This symbol denotes weed suppression which is a reduction in weed competition compared to untreated areas. A second treatment may be necessary.

\* This recommendation is for blackberry and dewberry control in bermudagrass or other non-sensitive grasses only. For control in tall fescue pastures, only apply Opensight as a spot treatment. For broadcast blackberry control in tall fescue pastures, use 1 pint/acre of Remedy Ultra + 2 pts/acre of ForeFront™ R&P.

Opensight alone provides brush control for a number of woody/perennial species. In most situations, Opensight is added to brush control tank mixtures to improve control of the species listed below.

#### Control of Woody Species with Opensight Alone

Common Name	Scientific Name	Life Cycle	Plant Family
ash	<i>Fraxinus spp.</i>	perennial	Oleaceae
aspen	<i>Populus tremuloides</i>	perennial	Salicaceae
camelthorn	<i>Alhagi pseudalhagi</i>	perennial	Fabaceae
cherry	<i>Prunus spp.</i>	perennial	Rosaceae
cottonwood	<i>Populus spp.</i>	perennial	Salicaceae
Eastern red cedar	<i>Juniperus virginiana</i>	perennial	Cupressaceae
elder	<i>Sambucus spp.</i>	perennial	Caprifoliaceae
elm	<i>Ulmus spp.</i>	perennial	Ulmaceae
firs	<i>Abies spp.</i>	perennial	Pinaceae
hawthorn	<i>Crataegus spp.</i>	perennial	Rosaceae
mulberry	<i>Morus spp.</i>	perennial	Moraceae
muscadine (wild grape)	<i>Muscadinia rotundifolia</i>	perennial	Vitaceae
oaks	<i>Quercus spp.</i>	perennial	Fagaceae
ocean spray	<i>Holodiscus discolor</i>	perennial	Rosaceae
Osage orange	<i>Maclura pomifera</i>	perennial	Moraceae
maple, red	<i>Acer rubrum</i>	perennial	Aceraceae
salmonberry	<i>Rubus spectabilis</i>	perennial	Rosaceae
spruce, black	<i>Picea mariana</i>	perennial	Pinaceae
spruce, white	<i>Picea glauca</i>	perennial	Pinaceae
thimbleberry	<i>Rubus parviflorus</i>	perennial	Rosaceae
tree of heaven	<i>Ailanthus altissima</i>	perennial	Simaroubaceae
willow	<i>Salix spp.</i>	perennial	Salicaceae
poplar, yellow	<i>Liriodendron tulipifera</i>	perennial	Magnoliaceae

#### Selective Weed Control with tank mixes

Opensight is tank mix compatible with other selective herbicides such as Garlon 3A. Spot treatments using a tank mixture of Garlon 3A at 3% to 5 % v/v + Opensight at 20 oz product per 100 gallons of water (0.2 oz product/gallon water) + non-ionic surfactant, will control the following species, **in addition to species listed above**, without harming the grasses.

**Control of Woody Species with Opensight in Tank Mixes**

Common Name	Scientific Name	Life Cycle	Plant Family
alder	<i>Alnus rubra</i>	perennial	Betulaceae
arrowweed	<i>Pluchea sericea</i>	perennial	Asteraceae
Australian pine	<i>Pinus nigra</i>	perennial	Pinaceae
bear clover (bearmat)	<i>Chamaebatia foliolosa</i>	perennial	Rosaceae
beech	<i>Fagus spp.</i>	perennial	Fagaceae
birch	<i>Betula spp.</i>	perennial	Betulaceae
blackgum	<i>Nyssa sylvatica</i>	perennial	Cornaceae
Brazilian pepper-tree	<i>Schinus terebinthifolius</i>	perennial	Anacardiaceae
cascara	<i>Rhamnus purshiana</i>	perennial	Rhamnaceae
ceanothus	<i>Ceanothus spp.</i>	perennial	Rhamnaceae
chinquapin	<i>Chrysolepis chrysophylla</i>	perennial	Fagaceae
choke cherry	<i>Prunus virginiana</i>	perennial	Rosaceae
dogwood	<i>Cornus spp.</i>	perennial	Cornaceae
Douglas-fir	<i>Pseudotsuga menziesii</i>	perennial	Pinaceae
elderberry	<i>Sambucus spp.</i>	perennial	Adoxaceae
gallberry	<i>Ilex coriacea</i>	perennial	Aquifoliaceae
hazel	<i>Corylus spp.</i>	perennial	Betulaceae
hornbean	<i>Corylus spp.</i>	perennial	Betulaceae
madrone	<i>Arbutus menziesii</i>	perennial	Ericaceae
maple	<i>Acer spp.</i>	perennial	Aceraceae
mulberry	<i>Morus spp.</i>	perennial	Moraceae
persimmon	<i>Diospyros spp.</i>	perennial	Ebenaceae
pine	<i>Pinus spp.</i>	perennial	Pinaceae
poison ivy	<i>Toxicodendron radicans</i>	perennial	Anacardiaceae
poison oak	<i>Toxicodendron pubescens</i>	perennial	Anacardiaceae
poplar	<i>Populus spp.</i>	perennial	Salicaceae
coyote bush	<i>Baccharis pilularis</i>	perennial	Asteraceae
sassafras	<i>Sassafras spp.</i>	perennial	Lauraceae
Scotch broom	<i>Cytisus scoparius</i>	perennial	Fabaceae
sumac	<i>Rhus coriaria</i>	perennial	Anacardiaceae
sweetbay magnolia	<i>Magnolia virginiana</i>	perennial	Magnoliaceae
sweetgum	<i>Liquidambar spp.</i>	perennial	Altingiaceae
sycamore	<i>Platanus spp.</i>	perennial	Platanaceae
tanoak	<i>Lithocarpus densiflorus</i>	perennial	Fagaceae
wax myrtle	<i>Myrica spp.</i>	perennial	Myricaceae
Western hemlock	<i>Tsuga heterophylla</i>	perennial	Pinaceae
winged elm	<i>Ulmus alata</i>	perennial	Ulmaceae

Apply either with a low volume backpack or handgun (hose reel 7 hydraulic spraygun). In all cases, use the amount specified to provide uniform and complete coverage of the plants to be controlled. Total spray volume should not exceed 16 gallons of spray mix per acre.

**Non-selective weed control with tank mixes**

Opensight is tank mix compatible with non-selective herbicides such as Accord XRT II or Rodeo at 3 to 4 quarts per acre + a non-ionic surfactant at 0.25% v/v for control of grasses and many broadleaf woody species such as red oak, white oak, cherry, sweetgum, loblolly pine, red maple, and yellow poplar.

---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

**Warranty Disclaimer**

---

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

**Limitation of Remedies**

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™®Trademarks of Corteva Agriscience and its affiliated companies

EPA accepted \_\_/\_\_/\_\_\_\_