

62719 615

7/26/2012

10238

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

July 26 2012

Darin W Lickfeldt
Dow AgroSciences
9330 Zionsville Road
Indianapolis IN 46268

Dear Dr Lickfeldt

Subject Amended label to revise the directions for use in nurseries
Product Name Cobalt® Advanced
EPA Registration No 62719 615
EPA Decision No 465682

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act is acceptable A stamped copy of the label is enclosed for your records Please submit one copy of your final printed labeling before you release the product for shipment Your release for shipment of the product constitutes acceptance of these conditions If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA section 6(e) If you have any questions please contact Julie Chao by phone at (703) 308 8735 or by email at chao.julie@epa.gov

Regards

A handwritten signature in black ink, appearing to read "Venus Eagle".

for Venus Eagle Product Manager (01)
Insecticide Rodenticide Branch
Registration Division (7505P)

Enclosure

(Base label)

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification

Cobalt[®] Advanced**Insecticide**

For control of listed insects infesting listed field, fruit, nut, and vegetable crops

Active Ingredients

chlorpyrifos O O diethyl O (3 5 6 trichloro 2 pyridinyl) phosphorothioate	28 12%
<i>Lambda</i> cyhalothrin [1 α (S) 3 α (Z)] (±) cyano (3 phenoxyphenyl)methyl 3 (2 chloro 3 3 3 trifluoro 1 propenyl) 2 2 dimethylcyclopropanecarboxylate	1 44%
Other Ingredients	70 44%
Total	100 00%

ACCEPTED
JUL 26 2012

**Under the Federal Insecticide Fungicide,
and Rodenticide Act, as amended for the
pesticide registered under**

EPA Reg No 62719-615

Contains 2 503 lb chlorpyrifos and 0 129 lb *lambda* cyhalothrin per gallon

Contains petroleum distillate

Keep Out Of Reach Of Children

WARNING AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

Precautionary Statements**Hazard to Humans and Domestic Animals**

**May Be Fatal If Swallowed Causes Skin Irritation • Causes Moderate Eye Irritation Harmful If
Absorbed Through Skin Harmful If Inhaled • Prolonged Or Frequently Repeated Skin Contact
May Cause Allergic Reactions In Some Individuals**

Do not get on skin or clothing Avoid breathing spray mist Avoid contact with eyes

Personal Protective Equipment (PPE)

Materials that are chemical resistant to this product are barrier laminate and viton ≥ 14 mils

Mixers and loaders using a mechanical transfer loading system and applicators using aerial application equipment must wear

Long sleeved shirt and long pants
Shoes and socks
Protective eyewear

In addition to the above **mixers and loaders** using a mechanical transfer loading system must wear

Chemical resistant gloves
Chemical resistant apron

A NIOSH approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC 21C or a NIOSH approved respirator with any R P or HE filter

See Engineering Controls for additional requirements

All **other mixers loaders applicators and handlers** must wear

Coveralls over long sleeved shirt and long pants

Chemical resistant gloves

Chemical resistant apron when mixing or loading or exposed to the concentrate

Chemical resistant footwear plus socks

Chemical resistant headgear for overhead exposure

A NIOSH approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC 21C or a NIOSH approved respirator with any R P or HE filter

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

Mixers and loaders supporting aerial applications must use a mechanical transfer system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for dermal protection and must

Wear the personal protective equipment required above for mixers/loaders

Wear protective eyewear

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]

Use of human flaggers is prohibited. Mechanical flagging equipment must be used.

When handlers use closed cab motorized ground application equipment in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product and wash contaminated clothing before reuse.

Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

Organophosphate

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

If in eyes Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Note to physician Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early, however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

Note to physician Contains petroleum distillate – vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, small mammals, and birds. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under Agricultural Use Requirements in the Directions for Use section for information about this standard.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage Store in original container in secured dry storage area. Prevent cross contamination with other pesticides and fertilizers.

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water food or feed by storage and disposal

Pesticide Storage Store in original container in secured dry storage area Prevent cross contamination with other pesticides and fertilizers

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Container Handling Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Cleaning the container before final disposal is the responsibility of the person disposing of the container

Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10% full with water and if possible spray all sides while adding water If practical agitate vigorously or recirculate water with the pump for two minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)**Storage and Disposal**

Do not contaminate water food or feed by storage and disposal

Pesticide Storage Store in original container in secured dry storage area Prevent cross contamination with other pesticides and fertilizers

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Refer to label booklet for Directions for Use

Notice Read the entire label Use only according to label directions **Before using this product read Warranty Disclaimer and Limitation of Remedies at end of label booklet If terms are unacceptable return at once unopened**

In case of emergency endangering health or the environment involving this product call 1 800 992 5994

Agricultural Chemical Do not ship or store with food feeds drugs or clothing

Shake well before use

EPA Reg No 62719 615

EPA Est _____

®Trademark of Dow AgroSciences LLC

600f 38

Q4J / Cobalt Advanced / Amend / 07 25 12

Page 5

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268

Net Contents __

(Cover shipping container)

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification

Cobalt[®] Advanced

Insecticide

For control of listed insects infesting listed field, fruit, nut, and vegetable crops

Active Ingredients

chlorpyrifos O O diethyl O	
(3 5 6 trichloro 2 pyridinyl)	
phosphorothioate	28 12%
Lambda cyhalothrin [1α(S) 3α(Z)] (±)	
cyano (3 phenoxyphenyl)methyl 3	
(2 chloro 3 3 3 trifluoro 1 propenyl) 2	
2 dimethylcyclopropanecarboxylate	1 44%

Other Ingredients	70 44%
-------------------	--------

Total	100 00%
-------	---------

Contains 2 503 lb chlorpyrifos and 0 129 lb lambda cyhalothrin per gallon

Contains petroleum distillate

Keep Out Of Reach Of Children

WARNING AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 Refer to label booklet under Agricultural Use Requirements in the Directions for Use section for information about this standard

Refer to inside of label booklet for additional precautionary information including Directions for Use

Notice Read the entire label Use only according to label directions **Before using this product read Warranty Disclaimer and Limitation of Remedies at end of label booklet If terms are unacceptable return at once unopened**

In case of emergency endangering health or the environment involving this product call 1 800 992 5994

Agricultural Chemical Do not ship or store with food feeds drugs or clothing

Shake well before use

EPA Reg No 62719 615

EPA Est _____

®Trademark of Dow AgroSciences LLC

Produced for

80 of 38

Q4J / Cobalt Advanced / Amend / 07 25 12

Page 7

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268

Net Contents __

(Page 1 through end)

Table of Contents	Page
Precautionary Statements	
Hazard to Humans and Domestic Animals	
Personal Protective Equipment (PPE)	
Engineering Controls	
User Safety Recommendations	
First Aid	
Environmental Hazards	
Physical and Chemical Hazards	
Directions for Use	
Agricultural Use Requirements	
Storage and Disposal	
General Information	
Use Precautions	
Spray Drift Management	
Application Directions	
Mixing Directions	
Uses	
Alfalfa	
Apple Tree Trunk	
Brussels Sprout	
Conifer and Deciduous Trees (Plantations Nurseries and Seed Orchards)	
Corn (Field Sweet Seed)	
Cotton	
Sorghum Grain Sorghum (Milo)	
Soybean	
Sunflower	
Tree Fruits and Almond (Dormant/Delayed Dormant Sprays)	
Tree Fruits and Almond (Trunk Spray or Preplant Dip)	
Tree Nuts (Foliar Sprays)	
Wheat	
Terms and Conditions of Use	
Warranty Disclaimer	
Inherent Risks of Use	
Limitation of Remedies	

Precautionary Statements

Hazard to Humans and Domestic Animals

WARNING

May Be Fatal If Swallowed • Causes Skin Irritation • Causes Moderate Eye Irritation • Harmful If Absorbed Through Skin • Harmful If Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get on skin or clothing Avoid breathing spray mist Avoid contact with eyes

Personal Protective Equipment (PPE)

Materials that are chemical resistant to this product are barrier laminate and viton ≥ 14 mils

Mixers and loaders using a mechanical transfer loading system and applicators using aerial application equipment must wear

- Long sleeved shirt and long pants
- Shoes and socks
- Protective eyewear

In addition to the above **mixers and loaders** using a mechanical transfer loading system must wear

- Chemical resistant gloves
- Chemical resistant apron
- A NIOSH approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC 21C or a NIOSH approved respirator with any R P or HE filter

See Engineering Controls for additional requirements

All other mixers loaders applicators and handlers must wear

- Coveralls over long sleeved shirt and long pants
- Chemical resistant gloves
- Chemical resistant apron when mixing or loading or exposed to the concentrate
- Chemical resistant footwear plus socks
- Chemical resistant headgear for overhead exposure
- A NIOSH approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC 21C or a NIOSH approved respirator with any R P or HE filter

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

Mixers and loaders supporting aerial applications must use a mechanical transfer system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for dermal protection and must

- Wear the personal protective equipment required above for mixers/loaders
- Wear protective eyewear

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]

Use of human flaggers is prohibited Mechanical flagging equipment must be used

When handlers use closed cab motorized ground application equipment in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Recommendations

Users should

Wash hands thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet

Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

Remove PPE immediately after handling this product and wash contaminated clothing before reuse

Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

First Aid

Organophosphate

If swallowed Immediately call a poison control center or doctor Do not induce vomiting unless told to do so by a poison control center or doctor Do not give any liquid to the person Do not give anything by mouth to an unconscious person

If on skin or clothing Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice

If inhaled Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a poison control center or doctor for further treatment advice

If in eyes Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice

Note to physician Chlorpyrifos is a cholinesterase inhibitor Treat symptomatically If exposed plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful) Atropine only by injection is the preferable antidote Oximes such as 2 PAM/protopam may be therapeutic if used early however use only in conjunction with atropine In case of severe acute poisoning use antidote immediately after establishing an open airway and respiration

Note to physician Contains petroleum distillate – vomiting may cause aspiration pneumonia

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 800 992 5994 for emergency medical treatment information

Environmental Hazards

This product is toxic to fish aquatic invertebrates small mammals and birds Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when disposing of equipment washwaters or rinsate This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area

Physical or Chemical Hazards

Combustible Do not use or store near heat or open flame

Directions for Use

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Read all Directions for Use carefully before applying

This product cannot be reformulated or repackaged into other end use products

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Certified crop advisors or persons entering under their direct supervision under certain circumstances may be exempt from the early reentry requirements pursuant to 40 CFR Part 170.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short sleeved shirt and short pants
- Chemical resistant gloves made out of any waterproof material
- Chemical resistant footwear plus socks
- Chemical resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross contamination with other pesticides and fertilizers.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable containers 5 gallons or less

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable containers 5 gallons or larger

Container Handling Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10% full with water and if possible spray all sides while adding water If practical agitate vigorously or recirculate water with the pump for two minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Nonrefillable containers 5 gallons or larger

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Product Information

Cobalt[®] Advanced insecticide is an emulsion in water for use in listed crops This product resists washoff once it is dry Target pests and application rates are provided in the accompanying tables

Use Precautions

Insect control may be reduced at low spray volumes under high temperature and wind conditions

Some reduction in insect control may occur under unusually cool conditions

Flood irrigation To avoid contamination of irrigation tail waters do not flood irrigate within 24 hours following a soil surface or foliar application of Cobalt Advanced

Spray Drift Management**Buffer Zones**

In New York State a 25 ft vegetated non cropped buffer strip not traversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh for both aerial or ground application For aerial applications the 25 ft vegetated non cropped buffer strip for runoff protection would be part of the larger 50 ft buffer strip (or 450 ft buffer strip for ULV application) required for spray drift

Vegetative Buffer Strip Construct and maintain a minimum 10 foot wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as lakes reservoirs rivers permanent streams marshes or natural ponds estuaries and commercial fish farm ponds)

Only apply products containing lambda cyhalothrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat

For guidance refer to the following publication for information on constructing and maintaining effective buffers *Conservation Buffers to Reduce Pesticide Losses* Natural Resources Conservation Services USDA NRCS 2000 Fort Worth Texas 21 pp <http://www.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>

Buffer Zone for Ground Application (Groundboom or Overhead Chemigation) Do not apply within 25 feet of aquatic habitats (such as lakes reservoirs rivers streams marshes ponds estuaries and commercial fish ponds)

Buffer Zone for Ground Application (Airblast) Do not apply within 50 feet of aquatic habitats (such as lakes reservoirs rivers streams marshes ponds estuaries and commercial fish ponds)

Buffer Zone for ULV Aerial Application Do not apply within 450 feet of aquatic habitats (such as lakes reservoirs rivers streams marshes ponds estuaries and commercial fish ponds)

Buffer Zone for Non ULV Aerial Application Do not apply within 150 feet of aquatic habitats (such as lakes reservoirs rivers streams marshes ponds estuaries and commercial fish ponds)

Spray Drift Requirements

Wind Direction and Speed Only apply this product if the wind direction favors on target deposition Do not apply when the wind velocity exceeds 15 mph

Temperature Inversion Do not make aerial or ground applications during a temperature inversions Inversions are characterized by stable air and increasing temperatures with height above the ground Mist or fog may indicate the presence of an inversion in humid areas The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface

Droplet Size Use only medium or coarser spray nozzles (for ground and non ULV aerial application) according to ASABE (S572.1) definition for standard nozzles In conditions of low humidity and high temperatures applicators should use a coarser droplet size

Additional Requirements for Ground Applications

- Wind speed must be measured adjacent to the application site on the upwind side immediately prior to application
For groundboom applications apply using a nozzle height of no more than 4 feet above the ground or crop canopy
- For airblast applications turn off outward pointing nozzles at row ends and when spraying the outer two rows To minimize spray loss over the top in orchard applications spray must be directed into the canopy

Additional Requirements for Aerial Applications

- Mount the spray boom on the aircraft so as to minimize drift caused by wingtip or rotor vortices Use the minimum practical boom length and do not exceed 75% of the wingspan or 80% of the rotor diameter
- Flight speed and nozzle orientation must be considered in determining droplet size
- Release spray at the lowest height consistent with pest control and flight safety Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety
- When applications are made with a crosswind the swath will be displaced downwind The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind

Application Directions

Broadcast Foliar Application

Apply with conventional power operated spray equipment using nozzles and spray pressures specified for insecticides. Apply Cobalt Advanced in a spray volume of not less than 2 gallons per acre (gpa) for aerial application equipment (fixed wing or helicopter) or not less than 10 gpa for ground equipment unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy.

Ground Application Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's specifications for insecticide nozzles with respect to nozzle type, pressure, and spacing.

Broadcast Soil Application

Apply with conventional power operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets. Unless otherwise indicated, apply in a spray volume of 10 gpa or more. For band application, use proportionally less spray volume.

Aerial Application

Use a minimum spray volume of 2 gpa. Mark swaths by mechanical flagging, permanent markers, or GPS equipment.

Chemigation Application

Apply Cobalt Advanced through properly equipped chemigation systems for insect control in alfalfa, corn, cotton, sorghum, soybean, and wheat or other crops as specified in Dow AgroSciences supplemental labeling. Do not apply this product by chemigation unless specified in crop specific directions in this label or Dow AgroSciences supplemental labeling. Do not apply to labeled crops through any other type of irrigation system.

Note Unless otherwise indicated in specific use directions, the application rates for chemigation are the same as those for broadcast application.

General Directions for Sprinkler Chemigation Cobalt Advanced may be applied through overhead sprinkler irrigation systems that will apply water uniformly, including center pivot, lateral move, end tow side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system. Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

For continuously moving systems, the mixture containing Cobalt Advanced must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. If continuously moving irrigation equipment is used, apply in no more than 0.25 inch of water. For irrigation systems that do not move during operation, apply in no more than 0.25 inch of irrigation immediately before the end of the irrigation cycle.

Chemigation Preparation The following use directions must be followed when Cobalt Advanced is applied through sprinkler irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of Cobalt Advanced needed to cover the desired acreage. Mix according to instructions in the Mixing Directions section and bring mixture to desired volume. Continually agitate the mixture during mixing and application.

Chemigation Equipment Calibration In order to calibrate the irrigation system and injector to apply the mixture containing Cobalt Advanced, determine the following: 1) Calculate the number of acres irrigated by the system. Calculate the amount of product required and premix. 3) Determine the irrigation rate and determine the number of minutes for the system to cover the intended treatment area. 4) Calculate the

total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes (minus time to flush out) to cover the treatment area. This value equals the gallons per minute output that the injector or eductor must deliver. Convert the gallons per minute to milliliters or ounces per minute if needed. Calibrate the injector system with the system in operation at the desired irrigation rate. It is suggested that the injection pump/system be checked at least twice before operation and the system monitored during operation.

Chemigation Equipment Requirements

- The system must contain an air gap, an approved back flow prevention device, a functional check valve, vacuum relief valve (including inspection port) and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineers Engineering Practice 409 for more information or state specific regulations.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection when the water pressure drops too low or water flow stops.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (or flow meter or eductor), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- To ensure uniform mixing of the insecticide into the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all back flow prevention devices on the water line.
- The tank holding the insecticide mixture must be free of rust, fertilizer, sediment, and foreign material and equipped with an in-line strainer situated between the tank and the injection point.

Chemigation Operation Start the water pump and system and let the system achieve the desired pressure and speed before starting the injector. Check for leaks and uniformity and make repairs before any chemigation takes place. Start the injector system and calibrate according to manufacturer's specifications. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, flush and clean the entire irrigation and injector system prior to shutting down the system.

Chemigation Precautions

- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non uniform distribution of treated water.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers, or other experts.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.

Chemigation Restrictions

Do not add crop oil when Cobalt is applied by chemigation.

Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection.

The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate non target areas

- Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops
- Do not enter treated area during the reentry interval unless required PPE is worn
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units

Maximum Application Rate

Do not exceed the allowed maximum application rate of ai per acre per year by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
alfalfa	0.12	0.06
Brussels sprout	0.24	0.12
conifer	0.24	0.12
corn	0.12	0.06
cotton	0.2	0.1
sorghum	0.08	0.04
soybean	0.06	0.03
sunflower	0.2	0.1
tree fruits	0.2	0.1
tree nuts	0.16	0.08
wheat	0.06	0.03

When the maximum application rate of lambda cyhalothrin is reached, no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Mixing Directions

Cobalt Advanced – Alone

To prepare the spray, add a portion of the required amount of water to the spray tank and with the spray tank agitator operating, add Cobalt Advanced. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

Cobalt Advanced – Tank Mix

Cobalt Advanced is compatible with insecticides, herbicides, miticides, and fungicides and non pressure fertilizer solutions commonly used except for alkaline materials such as Bordeaux mixture and lime. Conduct a small jar compatibility test prior to tank mixing. Prepare tank mixtures in the same manner as directed above for use of Cobalt Advanced alone. When tank mixing Cobalt Advanced with herbicides

add wettable powders first flowables second and emulsifiable concentrates last For best results when a fertilizer solution is involved use a fertilizer pesticide compatibility agent such as Unite or Complex Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture Do not allow spray mixtures to stand overnight

Tank Mix Compatibility Test Test compatibility of the intended tank mixture before adding Cobalt Advanced to the spray or mix tank Add proportional amounts of each tank mix ingredient to a pint or quart jar cap shake and let set 15 minutes Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used

Uses

Alfalfa

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

Apply as a broadcast foliar spray using aircraft or ground spray equipment Use a higher rate in the rate range for increased pest pressure Use a minimum spray volume of 2 gpa for aerial application (fixed wing or helicopter) or 10 gpa for ground equipment Use a spray volume of 5 gpa or more by air or up to 20 gpa by ground when foliage is dense and/or pest population is high and/or under high temperature and wind conditions Some reduction in insect control may occur under unusually cool conditions

Chemigation Cobalt Advanced may be applied through sprinkler irrigation systems to control listed foliar pests Use specified broadcast application rates See Chemigation Application section

Target Pests	Cobalt Advanced (fl oz/acre)
corn rootworm adults (spotted cucumber beetle) grasshoppers leafhoppers	6 13
alfalfa caterpillar blue alfalfa aphid cutworm spp green cloverworm pea aphid spotted alfalfa aphid (suppression) (not for use in California) velvetbean caterpillar webworm spp	11 26

alfalfa blotch leafminer	16 38
alfalfa seed chalcid adult	
alfalfa weevil larvae and adults	
armyworms	
bean leaf beetle	
blister beetle sp	
clover leaf weevil spp	
clover root borer adult	
clover root curculio spp adult	
clover stem borer adult	
corn earworm	
cowpea aphid	
cowpea curculio adult	
cowpea weevil adult	
Egyptian alfalfa weevil larvae and adults	
grape colapsis adult	
green June beetle adult	
green peach aphid	
Japanese beetle adult	
looper spp	
Mexican bean beetle	
pea weevil adult	
plant bugs	
spider mites	
spittlebug spp	
stink bug spp	
sweet clover weevil adult	
threecornered alfalfa hopper	
thrips spp	
whitefringed beetle spp adult	

Specific Use Precautions

- Do not tank mix Cobalt Advanced with other pesticides surfactants or fertilizer formulations unless prior use has shown the combination to be non injurious to alfalfa under current conditions of use Some phytotoxic symptoms may be observed on young tender rapidly growing alfalfa treated with Cobalt Advanced Alfalfa will outgrow these symptoms and no yield loss should be expected
- This product is highly toxic to bees exposed to direct treatment on alfalfa Do not apply if nearby bees are clustered outside of hives and bees are foraging in the treated area Protective information may be obtained from your Agricultural Extension Service
- To avoid contamination of irrigation tail waters do not flood irrigate within 24 hours following an application of Cobalt Advanced

Specific Use Restrictions

Preharvest Interval Do not cut or graze treated alfalfa within 7 days after application of 6 to 13 fl oz of Cobalt Advanced per acre within 14 days after application of 13 to 26 fl oz per acre or within 21 days after application of rates above 26 fl oz per acre

Maximum single application rate is 38 fl oz of Cobalt Advanced (0.75 lb ai chlorpyrifos and 0.04 lb ai lambda cyhalothrin) per acre

Do not apply more than a total of 119 fl oz of Cobalt Advanced (2.3 lb ai chlorpyrifos and 0.12 lb ai lambda cyhalothrin) per acre per season

Do not apply any product containing chlorpyrifos more than once per alfalfa cutting

Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products The maximum

rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
alfalfa	0 12	0 06

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin

Apple Tree Trunk

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn

Apply as a post bloom application to the lower 4 feet of the apple tree trunk for borer control in states east of the Rockies only Mix with water and apply directly to trunk from a distance of no more than 4 feet using low volume handgun or shielded spray equipment Do not allow spray to contact foliage or fruit

Target Pests	Cobalt Advanced (quart/100 gal)
American plum borer apple bark borer broad necked root borer dogwood borer flatheaded appletree borer roundheaded apple tree borer tilehorned prionus	1 3

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 28 days before harvest
- Maximum single application rate is 1 3 quarts of Cobalt Advanced (0 81 lb ai chlorpyrifos and 0 05 lb ai lambda cyhalothrin) per 100 gallons
- Do not make more than one application of Cobalt Advanced to the apple tree trunk per year as either a pre bloom or post bloom trunk application
- This product may not be used if a pre bloom application of any other product containing chlorpyrifos has been made during the year
- Do not allow meat or dairy animals to graze in treated orchards
- Treat only the lower 4 feet of the apple tree trunk
- Do not apply when wind speed is greater than 10 mph

Brussels Sprout

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

Apply with conventional power operated spray equipment in 20 to 150 gpa of water Use a higher rate in the rate range when there is increased pest pressure Consult your state agricultural experiment station extension service specialist or integrated pest control advisor for proper time to treat in your area

Target Pests	Cobalt Advanced (fl oz/acre)
cabbage aphid grasshoppers	11 19

cutworms fall armyworms imported cabbage worm	11 26
beet armyworm cabbage looper cabbage webworm southern cabbage worm stink bugs striped flea beetle (adult) yellowstriped armyworm	16 38

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 21 days before harvest
- Maximum single application rate is 38 fl oz of Cobalt Advanced (0.75 lb ai chlorpyrifos and 0.04 lb ai lambda cyhalothrin) per acre
- Do not make more than three applications of any product containing chlorpyrifos per crop
- If three applications are made at the maximum application rate of 38 fl oz then a fourth application of a product containing chlorpyrifos is not permitted. More applications with a product containing lambda cyhalothrin are permitted as long as no more than the maximum amount of 0.24 lb ai per acre per year is applied
- Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
Brussels sprout	0.24	0.12

When the maximum application rate of lambda cyhalothrin is reached, no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Conifer and Deciduous Trees (Plantations, Nurseries, and Seed Orchards)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use in nurseries is restricted to wholesale nursery operations. Wholesale nursery operations are commercial agricultural operations which do not sell or distribute directly to consumers or the general public through retail sales. Plants, trees, fruit, vegetables, or any parts of the plants, trees, fruits, or vegetables treated with this product cannot be sold or distributed directly to consumers or the general public through retail sales.

Apply Cobalt Advanced as required by scouting to control exposed foliage, flower, cone, seed, and bark feeding insects. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.

Unless otherwise indicated, apply only as a foliar spray using power operated ground equipment. Thorough coverage of foliage is essential. Use a minimum 10 gpa of finished spray with ground equipment. Use higher volume of finished spray, 20 gpa or more, when foliage is dense and/or pest density is high and/or under high temperature and wind conditions.

Target Pests		Cobalt Advanced (fl oz/acre)
aphids	pine needle scale (1)	45 51 1
European pine sawfly	pine tortoise scale (1)	
gypsy moth	spittlebugs	
pales weevil (adult)	spruce budworm	
pales weevil (2)		13 15 33 (130 2 153 3 fl oz/100 gal)
coneworm spp (3)	seed bug spp (3)	0 5 2 (5 12 – 20 fl oz/100 gal)

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- 1 For **scale control** apply when scale crawlers are active
- 2 Apply as a cut stump drench
- 3 **Coneworm/Seed Bug/Thrips spp in Seed Orchards**
 - For high volume sprayers dilute 5 12 fl oz of Cobalt Advanced per 100 gallons of water and apply 5 to 10 gallons of finished spray per tree
 - For low volume sprayers dilute 20 fl oz of Cobalt Advanced per 100 gallons of water and apply 100 gallons of finished spray volume per acre

Specific Use Precautions

Phytotoxicity Do not apply under conditions of extreme heat or drought stress. Environmental factors and varietal differences significantly influence potential phytotoxic expression. **Testing has shown that Cobalt Advanced may be used at specified rates on the following conifer species without serious phytotoxicity:** balsam fir, concolor fir, Douglas fir, eastern white pine, Fraser fir, grand fir, noble fir, Scotch pine, white spruce. Before treating large numbers of other conifer species, treat a small block of plants and observe 7 to 10 days for symptoms of phytotoxicity. **Note:** The user assumes responsibility for determining if it is safe to treat other conifer species with Cobalt Advanced under commercial growing conditions.

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 30 days before harvest
- **Chemigation** Do not apply this product through any type of irrigation system
- Maximum single application rate is 51 1 fl oz of Cobalt Advanced (1 lb ai chlorpyrifos and 0 06 lb ai lambda cyhalothrin) per acre
- Do not make more than three applications of Cobalt Advanced or other product containing chlorpyrifos per season
- Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 7 days of the first application
- Do not allow meat or dairy animals to graze in treated areas
- In plantations and nurseries do not apply more than 0 24 lb ai lambda cyhalothrin per acre per year
- In seed orchards do not apply more than 0 5 lb ai lambda cyhalothrin per acre per year

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table.

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
conifer	0 24	0 12

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Corn (Field Sweet, Seed)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Conservation Tillage Preplant At Plant or Preemergence Applications

Apply as a broadcast spray to surface trash and exposed soil using power operated ground spray equipment. Use a total spray volume of 20 gpa or more. Use a higher rate in the rate range to extend residual control.

Tank Mixing Cobalt Advanced may also be applied in tank mixtures with paraquat or glyphosate herbicide and/or liquid fertilizer solutions. See Mixing Directions section for tank mixing instructions. Read and carefully follow all applicable directions, restrictions and precautions on labeling for each product use in combination with Cobalt Advanced.

Target Pests	Cobalt Advanced (fl oz/acre)
armyworms cutworms	11 – 38

At Plant T Band Application

Apply as a liquid T band in fields with no more than 30% cover of crop residue remaining on the soil surface. Apply Cobalt Advanced at a rate of 3.8 fl oz per 1000 linear feet of row (51 fl oz per acre with 40 inch row spacing) in a minimum spray volume of 5 gpa. Position a flat fan nozzle over the open seed furrow immediately behind the planter shoe, in front of the press wheel, and adjust to provide a 5 to 6 inch band width centered over the row. Incorporate into the top 1 inch of soil using tines, chains or other suitable equipment.

The following table provides equivalent application rates for various row spacings when Cobalt Advanced is applied at the rate of 2.87 fl oz per 1000 ft of row for grubs, seed corn beetle, seed corn maggot and wireworms, or applied at a rate of 1.89 fl oz per 1000 ft of row for cutworms alone.

Target Pests	Amount of Cobalt Advanced Required	
	Row Spacing (inches)	fl oz/acre
cutworms	30	33
	36	28
	38	27
	40	26
grubs	30	50
seed corn beetle	36	42
seed corn maggot	38	40
wireworms	40	38

Sprayer Calibration Information for Band Application

Fluid Ounces of Spray Required Per 100 Feet of Row for Various Row Spacings and Spray Volumes				
Volume of Spray Per Acre (gal)	30	36	38	40
5	3.67	4.41	4.65	4.90
10	7.34	8.82	9.30	9.80

15	11 00	13 23	13 95	14 70
20	14 68	17 64	18 60	19 69

Postemergence Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment. Use higher carrier volume when foliage is dense and/or pest pressure is high and/or under high temperatures and wind conditions. Cobalt Advanced may be tank mixed with glyphosate products when application is to be made to glyphosate tolerant corn.

Chemigation Cobalt Advanced may be broadcast applied postemergence through sprinkler irrigation systems at specified application rates to control listed foliar pests. For best results, tank mix Cobalt Advanced with 2 pints of non emulsifiable oil. See Chemigation Application section.

Target Pests	Cobalt Advanced (fl oz/acre)
grasshoppers	6 13
aphids	11 26
armyworms	
bean leaf beetle	
cereal leaf beetle	
corn rootworm adults (2)	
cutworms (3)	
flea beetle adults (1)	
green cloverworm	
greenbug	
meadow spittlebug	
southern corn leaf beetle	
webworms (4)	
western bean cutworm	
aster leafhopper	16 38
chinch bugs (1)	
corn earworm	
European corn borer (5)	
hop vine borer	
sap beetle	
southwestern corn borer (6)	
stalk borer	
stink bug spp	
tarnished plant bug	
billbugs (1)	32 42
common stalk borer (9)	
corn rootworm larvae (7) (8)	
Japanese beetle adult	
lesser cornstalk borer	

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- Billbug, chinch bug, or flea beetle** For best control, ground apply in a minimum spray volume of 20 to 40 gpa at 40 psi. If corn is less than 6 inches tall, apply in a 9 to 12 inch wide band over the row. For corn more than 6 inches tall, apply using drop nozzles directed to the base of the plant. Do not reduce the application rate for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pest pressure, a second application may be needed.
- The specified dosage will control silk clipping by **corn rootworm adults**.

- 3 **Cutworms** It is preferable to apply Cobalt Advanced when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy, or crusted at time of treatment, worms may be protected from the spray and effectiveness will be reduced. Shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. A second application may be required if damage or density levels exceed economic thresholds established for your area.
- 4 **Webworm** For control, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary.
- 5 **European corn borer** For control, use 22 to 38 fl oz per acre when application is made with power operated ground or aerial equipment or 16 to 38 fl oz per acre when application is made through a sprinkler irrigation system. University research indicates that achieving greater than 50% control of first generation European borer with a single liquid insecticide treatment is highly dependent upon timing, insecticide placement, and weather conditions.
- 6 **Southwestern corn borer** A second application may be applied 21 days later if needed due to reinfestation.
- 7 **Corn rootworm larvae** For postemergence control, apply at cultivation. Direct the spray to both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. A cultivation application of Cobalt Advanced may be made in addition to an at planting application of Lorsban® 15G insecticide.
- 8 Cobalt Advanced may also be applied through sprinkler irrigation systems at the rate of 32 to 42 fl oz per acre to control **corn rootworm larvae**. Time application to coincide with the appearance of the second instar larvae. Apply with enough water to wet the root zone to the depth control needed. If soils are wet, allow enough soil drying to occur such that an application using a minimum amount of water will not produce surface runoff. See Chemigation Application section.
- 9 Do not use Cobalt Advanced in combination with a burndown herbicide for control of common stalk borer. For **common stalk borer** control, treat approximately 11 days after application of glyphosate or after burndown with paraquat herbicide is complete (3 to 5 days).

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 21 days before harvest of grain, ears, forage or fodder.
- Do not make more than three applications of any product containing chlorpyrifos per season, including the maximum allowed of two granular applications at the 1 lb ai chlorpyrifos rate.
- Do not apply more than a total of 129 fl oz of Cobalt Advanced (2.5 lb ai chlorpyrifos and 0.13 lb ai lambda cyhalothrin) per acre per season.
- Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application.
- If more than 1 lb ai granular chlorpyrifos per acre is applied at plant (for a maximum of 1.3 lb ai per acre per season), only one application of a liquid product containing chlorpyrifos at 42 fl oz (0.82 lb ai) per acre is allowed per season for a total of 2.12 lb ai chlorpyrifos per acre per season.
- Do not allow meat or dairy animals to graze in treated areas nor harvest treated corn silage as feed for meat or dairy animals within 14 days after last treatment.
- Do not apply in tank mixes with Steadfast or Lightning herbicides.
- Do not apply more than 79 fl oz after silk initiation.
- Do not apply more than 39.5 fl oz after corn has reached the milk stage (yellow kernels with milky fluid).

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table.

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
corn	0.12	0.06

When the maximum application rate of lambda cyhalothrin is reached, no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Cotton

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

Apply as a broadcast foliar spray using aircraft or ground spray equipment (see separate rate table for Arizona and California) Use a higher rate in the rate range when there is increased pest pressure Use sufficient spray volume to ensure thorough coverage of treated plants but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment Increase spray volume when foliage is dense and/or pest population is high and/or under high temperature and wind conditions Treat when field counts indicate damaging insect populations are developing or present

Chemigation Cobalt Advanced may be applied through sprinkler irrigation systems at specified broadcast application rates to control listed foliar pests See Chemigation Application section

Proper application methods are necessary to ensure thorough spray coverage and correct rate and minimize off target drift Follow Application Directions for ground and aerial application and Spray Drift Management recommendations in Product Information section of this label

All States Except Arizona and California

Target Pests	Cobalt Advanced (fl oz/acre)
cabbage looper cotton leafperforator cutworms grasshoppers	11 – 26
bollworm (2) cotton aphid cotton fleahopper (1) cotton leafworm cutworms spp European corn borer fall armyworm plant bugs (1) (<i>Lygus Mirids</i>) saltmarsh caterpillar thrips spp yellowstriped armyworm	16 38
beet armyworm boll weevil pink bollworm stink bug spp	22 38
bandedwing whitefly sweetpotato whitefly tobacco whitefly	32 – 42

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- 1 Suppression that will minimize damage from **plant bugs** and **cotton fleahoppers** Use a higher rate in the rate range for increased levels of control
- 2 **Bollworms** For best results scout fields twice per week and apply when worms are 1/4 inch or less in length

Arizona and California

Target Pests	Cobalt Advanced (fl oz/acre)
armyworms bollworm (2) cotton aphid cotton fleahopper cotton leaf perforator cutworms <i>Lygus</i> salt marsh caterpillar silverleaf whitefly (1) thrips	22 – 42
pink bollworm	32 42

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- 1 **Silverleaf whitefly** Apply in tank mix combination with the specified rate of another insecticide labeled for control or suppression
- 2 **Bollworms** For best results scout fields twice per week and apply when worms are 1/4 inch or less in length

Specific Use Restrictions

Preharvest Interval Do not apply within 21 days before harvest

- Maximum single application rate is 42 fl oz of Cobalt Advanced (0.82 lb ai chlorpyrifos and 0.05 lb ai lambda cyhalothrin) per acre
- Do not make more than three applications of Cobalt Advanced or other product containing chlorpyrifos per crop season
- Do not apply more than a total of 126 fl oz of Cobalt Advanced (2.5 lb ai chlorpyrifos and 0.127 lb ai lambda cyhalothrin) per acre per season
- Do not make a second application of Cobalt or other product containing chlorpyrifos within 10 days of the first application

Do not allow meat or dairy animals to graze in treated areas

Do not feed gin trash or treated forage to meat or dairy animals

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
cotton	0.2	0.1

When the maximum application rate of gamma cyhalothrin is reached, no lambda cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Sorghum Grain Sorghum (Milo)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment. Control may be reduced at low spray volumes under high temperature and wind conditions.

Chemigation Cobalt Advanced may be applied through sprinkler irrigation systems at specified broadcast application rates to control listed foliar pests. See Chemigation Application section.

Target Pests	Cobalt Advanced (fl oz/acre)
grasshoppers sorghum midge (1) yellow sugar cane aphid and other aphids	6 13
chinch bugs (3) cutworms fall armyworms flea beetle spp greenbug (2) lesser cornstalk borer (3) yellowstriped armyworm	11 38
beet armyworm corn earworm European and southwestern corn borer stink bug spp webworms	16 38

Numbers in parentheses () refer to Pest Specific Use Directions

Pest-Specific Use Directions

- 1 **Sorghum midge** Apply when 30% to 50% of the seed heads are in bloom
- 2 **Greenbug** Use a higher rate in the rate range when pest populations are high
- 3 **Chinch bugs and lesser cornstalk borer** Apply as a directed spray toward the base of the plant using power operated ground spray equipment with sufficient water to ensure coverage of an 8 to 12 inch band centered in the row. For plants less than 6 inches high, apply an 8 to 12 inch band centered over the row. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone.

Specific Use Precautions

To minimize the potential for chemical injury, do not apply Cobalt Advanced to drought stressed grain sorghum within three days following irrigation or rain except where the product is applied in irrigation water.

Be aware that sorghum lines used in seed production fields may be more susceptible to chemical injury. Susceptible inbred lines or hybrids are likely to be at greater risk of yield reducing chemical injury when treated at the higher application rates. Do not apply more than 26 fl oz of Cobalt Advanced per acre to seed sorghum if the additional risk of crop injury is unacceptable.

Specific Use Restrictions

- **Preharvest Interval** Do not harvest for grain, forage, fodder, hay, or silage within 30 days after application of 26 fl oz of Cobalt Advanced per acre or within 60 days after application of rates above 26 fl oz per acre.
- Maximum single application rate is 38 fl oz of Cobalt Advanced (0.75 lb a.i. chlorpyrifos and 0.04 lb a.i. lambda cyhalothrin) per acre per season.
- Do not make more than three applications of Cobalt Advanced or other product containing chlorpyrifos per use season.
- Do not apply more than a total of 77 fl oz of Cobalt Advanced (1.5 lb a.i. chlorpyrifos and 0.08 lb a.i. lambda cyhalothrin) per acre per season.
- Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application.
- Do not treat sweet varieties of sorghum.

- Do not apply more than 28 fl oz per acre per season once crop is in soft dough stage

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
sorghum	0.08	0.04

When the maximum application rate of lambda cyhalothrin is reached, no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Soybean

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Conservation Tillage At Plant or Preemergence Applications

Apply as a broadcast spray to surface trash and exposed soil using power operated ground spray equipment. Use a total spray volume of 20 gpa or more. Use a higher rate in the rate range to extend residual control.

Tank Mixing Cobalt Advanced may also be applied in tank mixtures with paraquat or glyphosate herbicide and/or liquid fertilizer solutions. See Mixing Directions section for tank mixing instructions. Read and carefully follow all applicable directions, restrictions and precautions on labeling for each product use in combination with Cobalt Advanced.

Target Pests	Cobalt Advanced (fl oz/acre)
cutworms grasshoppers lesser corn stalk borer	11 – 38

Postemergence Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment. Use higher carrier volume when foliage is dense and/or pest pressure is high and/or under high temperatures and wind conditions. Apply when field counts indicate damaging pest populations are developing or present. Cobalt Advanced may be tank mixed with glyphosate products when application is to be made to glyphosate tolerant soybeans. Use a higher rate in the rate range when there is increased pest pressure.

Chemigation Cobalt Advanced may be applied through sprinkler irrigation systems at specified broadcast application rates to control listed foliar pests. See Chemigation Application section.

Target Pests	Cobalt Advanced (fl oz/acre)
grasshoppers green cloverworm velvetbean caterpillar	6 – 13

blister beetle spp cabbage looper cutworms painted lady caterpillar saltmarsh caterpillar silverspotted skipper soybean aphid spider mites webworm spp woollybear caterpillar yellowstriped armyworm	11 26
bean leaf beetle corn earworm Japanese beetle adult Mexican bean beetle Mexican corn rootworm adult northern corn rootworm adult potato leafhopper southern corn rootworm beetle adult stink bug spp threecornered alfalfa hopper thrips spp western corn rootworm beetle adult	16 – 38
beet armyworm European corn borer lesser cornstalk borer	22 38

Specific Use Precaution

On determinate soybeans do not make more than one application after pod set

Specific Use Restrictions

Preharvest Interval Do not apply within 30 days before harvest

Maximum single application rate is 38 fl oz of Cobalt Advanced (0 75 lb ai chlorpyrifos and 0 04 lb ai lambda cyhalothrin) per acre

Do not make more than three applications per year of Cobalt Advanced or other product containing chlorpyrifos

Do not apply more than a total of 59 fl oz of Cobalt Advanced (1 15 lb ai chlorpyrifos and 0 06 lb ai lambda cyhalothrin) per acre per season

Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 14 days of the first application

Do not allow meat or dairy animals to graze in treated areas or otherwise feed treated soybean forage hay and straw to meat or dairy animals

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
soybean	0 06	0 03

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin

Sunflower

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

Postemergence Broadcast Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment Use a higher rate in the rate range when there is increased pest pressure

Target Pests	Cobalt Advanced (fl oz/acre)
grasshoppers	6 13
banded sunflower moth cutworms fall armyworm painted lady caterpillar seed weevil (4) stem weevil (2) sunflower beetle larvae and adults (1) sunflower moth (3) woolly bears	16 38
beet armyworm head clipper weevil adult Japanese beetle adult leafhopper spp meadow spittlebug spotted cabbage looper stink bug spp sunflower maggot adult tarnished plant bug (<i>Lygus</i>) (5)	22 – 38

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- Sunflower beetle** For control of larvae or adults treat when field counts indicate 10 larvae or 1 to 2 adults per seedling
- Stem weevil** Optimal treatment time is within 5 to 7 days after adult weevils begin to appear
- Sunflower moth** To control make first application during early 1% to 5% bloom stage
- Seed weevil** To control apply when field counts indicate 10 to 12 adults per plant for oil crop varieties and 1 to 3 adults per plant on confectionery crop varieties
- Tarnished plant bug (*Lygus*)** Use a higher rate in the rate range where populations are heavy Apply at the onset of pollen spread or approximately 10% bloom (R 5 growth stage) For best protection make a second application 10 days later Use sufficient water to ensure thorough coverage of treated plants

Specific Use Restrictions

Preharvest Interval Do not apply within 45 days before harvest

Maximum single application rate is 38 fl oz of Cobalt Advanced (0.75 lb ai chlorpyrifos and 0.04 lb ai lambda cyhalothrin) per acre

Do not make more than three applications of Cobalt Advanced or other products containing chlorpyrifos per season

Do not apply more than a total of 114 fl oz of Cobalt Advanced (2.3 lb ai chlorpyrifos and 0.12 lb ai lambda cyhalothrin) per acre per season or after bloom initiation

Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application

Do not allow meat or dairy animals to graze in treated areas

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
sunflower	0.2	0.1

When the maximum application rate of lambda cyhalothrin is reached, no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Tree Fruits¹ and Almond (Dormant/Delayed Dormant Sprays)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days for tree fruits and 24 hours for almond unless PPE required for early entry is worn.

¹Apple, cherry, nectarine, peach, pear, plum, prune

Apply as a dormant or delayed dormant spray. While Cobalt Advanced may be used without oil for best results, use oil to control additional pests. See precautions for use of oil below. Apply as a concentrate or dilute spray using conventional power operated spray equipment. For dilute sprays (greater than 200 gpa) use sufficient spray volume to completely wet tree foliage but not to point of runoff. For concentrate sprays (less than 200 gpa) uniformly apply an equivalent amount of Cobalt Advanced per acre.

Use a higher rate in the rate range when there is increased pest pressure.

Use Precautions for Tree Fruits and Almond

Cold or dry conditions may cause Cobalt Advanced plus oil sprays to infuse into trees, resulting in bud damage or bud drop. Do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated.

To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours of application of Cobalt Advanced.

Use Restrictions for Tree Fruits and Almond

Make only one application of chlorpyrifos during the dormant season.

Do not use more than a total of 6.25 pints of Cobalt Advanced (2 lb ai chlorpyrifos and 0.1 lb ai lambda cyhalothrin) per acre per season as a dormant/delayed dormant application.

Do not allow meat or dairy animals to graze in treated orchards.

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table.

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
tree fruits	0.2	0.1
almond	0.16	0.08

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Almond Cherry Nectarine Peach Pear Plum Prune

Target Pests	Cobalt Advanced (pint/acre)
American plum borer greater peach tree borer lesser peach tree borer peach twig borer	4 – 6 25

Specific Use Precautions for Almond Cherry Nectarine Peach Pear Plum Prune

Avoid contact with foliage in sweet cherries as premature leaf drop may result.

Specific Use Restrictions for Almond Cherry Nectarine Peach Pear Plum Prune

Do not make a soil or foliar application of Cobalt Advanced or product containing chlorpyrifos within 10 days of a dormant/delayed dormant application of chlorpyrifos to the orchard.

Do not exceed the maximum rate of 0.08 lb ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the Maximum Application Rate table.

Additional Restrictions Specific to California for Almond Cherry Nectarine Peach Pear Plum Prune

Do not use more than 1% dormant oil in almond orchards less than 4 years old.

Use a minimum of 100 gpa of total spray volume.

Use up to 2% supreme oil with no more than 4 gpa on almonds.

Use up to 2% supreme oil with no more than 6 gpa on peaches and nectarines.

Refer to the University of California pest management guide for pears, plums, and prunes.

Do not use any adjuvants or surfactants in addition to or as a substitute for a petroleum spray oil in a tank mix with Cobalt Advanced.

Do not apply on almonds in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo, and Yuba.

Apple

Target Pests	Cobalt Advanced (pint/acre)
rosy apple aphid san jose scale	4 – 6 25

Specific Use Restrictions for Apple

Only one application of any chlorpyrifos containing product can be made per year. The application can be either a prebloom dormant/delayed dormant spray to the canopy or the trunk, or a post bloom application to the lower 4 feet of the trunk [for post bloom application instructions and restrictions on apple, refer to Apple Tree Trunk section of the label].

Additional Restrictions Specific to California for Apple

Use a minimum of 100 gpa of total spray volume.

Refer to the University of California pest management guide for apples.

Do not use any adjuvants or surfactants in addition to or as a substitute for a petroleum spray oil in a tank mix with Cobalt Advanced.

Tree Fruits¹ and Almond (Trunk Spray or Preplant Dip)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days for tree fruits and 24 hours for almond unless PPE required for early entry is worn

¹Cherry nectarine peach plum

Apply Cobalt Advanced to tree trunks and lower branches using a coarse low pressure spray to control pests listed in the following table Use a higher rate in the rate range when there is increased pest pressure Unless otherwise specified a second application may be made after two weeks and a third application may be made after harvest Apply as required by scouting usually at intervals of five days or more Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds and IPM recommendations Avoid spray contact with foliage in sweet cherries as premature leaf drop may result Consult your state agricultural experiment station or extension service specialist for proper application timing for your area

Crops	Target Pests	Cobalt Advanced (quart/100 gal)
cherry	American plum borer greater peach tree borer lesser peach tree borer	2.4 – 4.8
almond peach nectarine plum	peach tree borers (1) (2)	4.8

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- Preplant Dip Application (Peaches and Nectarines Only)** For preplant control of peachtree borer use Cobalt Advanced at the equivalent application rate of 3 quarts per 100 gallons of water Dip trees several inches above the grafting bud scar and plant immediately or allow them to dry before returning to storage Do not allow peach trees to remain in contact with the dip solution
- Peach tree borer** For control in established trees apply before newly hatched borers enter the tree Use as a coarse low pressure trunk spray and thoroughly wet all bark areas from ground level to scaffold limbs Do not allow spray to contact fruit Consult written recommendations provided by your State agricultural experiment station or extension service specialist for proper time to treat in your area

Specific Use Restrictions

Preharvest Interval Do not apply within 14 days before harvest of almonds nectarines peaches and plums or within 21 days before harvest of cherries

Do not make more than one chlorpyrifos application per year in peaches and nectarines and no more than three chlorpyrifos applications per year in cherries

Do not exceed the maximum rate of 0.16 lb ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the Maximum Application Rate table

Do not allow meat or dairy animals to graze in treated orchards

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
------	---	--

tree fruits	0 2	0 1
tree nuts	0 16	0 08

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Tree Nuts¹ (Foliar Sprays)

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

¹Almond filbert pecan walnut

Apply Cobalt Advanced as a foliar spray at the dosages indicated to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop and apply as a concentrate or dilute spray using conventional power operated spray equipment. For dilute sprays applied to tree nut crops mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays apply an equivalent amount of Cobalt Advanced per acre. Treat when pests appear or in accordance with local conditions. Aerial application may result in less effective insect control because of reduced coverage. Consult your State agricultural experiment station, certified pest control advisor, or extension service specialist for specific use information in your area.

Crops	Target Pests	Cobalt Advanced (fl oz/acre)
almond	ant spp leaffooted bug leafroller spp navel orangeworm peach twig borer stink bug spp	22 – 57
filbert	eye spotted bud moth filbert aphid filbert leafroller filbert worm obliquebanded leafroller omnivorous leaftier stink bug spp winter moth	
pecan	blackmargined aphid fall webworm pecan nut casebearer yellow pecan aphid	16 – 57
	ant spp black pecan aphid hickory shuckworm (2) Phylloxera spp (3) pecan leaf scorch mite (suppression) (4) spittlebugs (1) stink bug spp	22 – 57
walnut	ant spp codling moth leaffooted bug leafrollers navel orangeworm stink bug spp walnut aphid walnut husk fly walnut scale	

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- Spittlebug** For control use a dosage of 57 fl oz per acre for concentrate sprays
- Hickory shuckworm** For best results make two applications 10 to 14 days apart
- Phylloxera spp** For best control make two applications at a 10 day interval using a minimum of 26 fl oz of Cobalt Advanced per acre starting at bud swell
- Pecan leaf scorch mite** For suppression use a preventative program

Specific Use Precautions

- Cobalt Advanced is highly toxic to bees exposed to direct treatment and should not be applied when bees are foraging in the treated area
- To avoid contamination of irrigation tail waters do not flood irrigate within 24 hours of application of Cobalt Advanced

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 14 days before harvest of almonds filberts and walnuts or 28 days of harvest of pecans
- Maximum single application rate is 57 fl oz of Cobalt Advanced (1 12 lb ai chlorpyrifos and 0 06 lb ai lambda cyhalothrin) per acre
- Do not make more than three total applications of Cobalt Advanced or other product containing chlorpyrifos to almonds pecans and filberts per season and no more than two applications on walnuts per season
- Do not apply more than a total of 158 fl oz of Cobalt Advanced (3 09 lb ai chlorpyrifos and 0 16 lb ai lambda cyhalothrin) per acre per season as a foliar spray
- Do not make a second application of Cobalt Advanced or other product containing chlorpyrifos within 10 days of the first application
- Do not allow meat or dairy animals to graze in treated orchards

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
tree nuts	0 16	0 08

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin

Wheat

Worker Restricted Entry Interval Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn

(For use only in Arizona California Colorado Idaho Kansas Minnesota Montana Nebraska New Mexico Nevada North Dakota Oklahoma Oregon South Dakota Texas Utah Washington and Wyoming)

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants but no less than 10 gpa for ground spray equipment or 2 gpa for aircraft equipment

Chemigation Cobalt Advanced may be applied through sprinkler irrigation systems at specified broadcast application rates to control listed foliar pests See Chemigation Application section

Target Pests	Cobalt Advanced (fl oz/acre)
aphid spp (1) English grain aphid greenbug oat bird cherry aphid Russian wheat aphid brown wheat mite grasshoppers	6 13

army cutworms (2)	11 ~ 25
armyworm spp	
cereal leaf beetle (3)	
cutworms (2)	
flea beetle spp	
grass sawfly	
orange wheat blossom midge	
wheat midge (4)	
stink bug spp	16 25

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions

- 1 Consult university extension bulletins for local treatment recommendations
- 2 Control may be reduced under high temperature conditions (greater than 80 F) under dry soil conditions or if larvae are more than 1/2 inch long
- 3 Target application when eggs are near hatching and larvae is emerging as monitored by plant inspection
- 4 **Wheat midge** For control treat when 75% of the wheat heads have emerged from the boot and when midge adults are found in the crop (1 midge per 4 to 5 heads) If possible apply in the late afternoon or early evening when temperatures exceed 50 F and wind speed is less than 7 mph

Specific Use Restrictions

- **Preharvest Interval** Do not apply within 14 days before harvest of forage and hay or within 30 days before harvest of grain and straw
- Maximum single application rate is 25 fl oz of Cobalt Advanced (0.5 lb ai chlorpyrifos and 0.025 lb ai lambda cyhalothrin) per acre
- Do not apply more than a total of 50 fl oz of Cobalt Advanced (1 lb ai chlorpyrifos and 0.05 lb ai lambda cyhalothrin) per acre per year
- Do not make more than two applications of Cobalt Advanced or product containing chlorpyrifos per season
- Do not allow meat or dairy animals to graze or otherwise feed on treated forage within 7 days after last treatment
- Do not feed straw from treated wheat within 30 days of application

Maximum Application Rate Do not exceed the maximum application rate of ai per acre per year allowed by using other lambda cyhalothrin or gamma cyhalothrin containing products. The maximum rate allowed for use if both lambda cyhalothrin and gamma cyhalothrin products are used during the same crop growing season can be calculated based upon the listed maximum rates in the following table

Crop	Lambda Cyhalothrin (lb ai/acre/season)	Gamma Cyhalothrin (lb ai/acre/season)
wheat	0.12	0.06

When the maximum application rate of lambda cyhalothrin is reached no gamma cyhalothrin product can be used. The gamma cyhalothrin quantity can be multiplied by 2 to calculate the total ai based upon lambda cyhalothrin.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions subject to the inherent risks set forth below TO THE EXTENT PERMITTED BY LAW Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product Crop injury lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures soil conditions etc) abnormal conditions (such as excessive rainfall drought tornadoes hurricanes) presence of other materials the manner of application or other factors all of which are beyond the control of Dow AgroSciences or the seller All such risks shall be assumed by buyer

Limitation of Remedies

To the extent permitted by law the exclusive remedy for losses or damages resulting from this product (including claims based on contract negligence strict liability or other legal theories) shall be limited to at Dow AgroSciences election one of the following

- (1) Refund of purchase price paid by buyer or user for product bought or
- (2) Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses

The terms of the Warranty Disclaimer Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner

™®Trademark of Dow AgroSciences LLC
EPA accepted __/__/__