

62719-621

07/12/2012

1/48



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF
CHEMICAL SAFETY
AND POLLUTION PREVENTION

JUL 12 2012

Dr Kerry Hastings
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268


Subject Entrust SC EPA Reg No 62719 621
 Date of Registrant Submission April 20 2012
 Decisions 464884

Dear Dr Hastings

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) is acceptable

Two (2) copies of the finished labeling must be submitted prior to releasing each product for shipment If you have any questions regarding this letter please contact Samantha Hulkower at (703) 603 0683

Sincerely


✓ Mark Suarez
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosures Copy of Label Stamped Accepted

T6P / Entrust SC / MSTR Amend / 04 12 12
file Entrust SC 621 MSTR 12Apr12d doc

Entrust[®] SC

EPA Reg No 62719 621

Registration Notes

Source label text based on EPA accepted text dated January 10 2012

Proposed changes by amendment

- 1 Added OMRI logo and statement Listed by the Organic Materials Review Institute (OMRI) for use in organic production
- 2 Product Information Changed General to Product
- 3 Use Precautions Deleted General from heading

[®]Trademark of Dow AgroSciences LLC

F
E
L
R 3

c

3/48

(Base label)

Entrust[®] SC**Naturalyte[○] Insect Control**

A Naturalyte[○] insect control product formulated for control of lepidopterous larvae (worms or caterpillars), leafminers, thrips, and red imported fire ants

Group	5	INSECTICIDE
-------	---	-------------


Active Ingredient

spinosad (a mixture of spinosyn A and spinosyn D) 22.5%

Other Ingredients 77.5%

Total 100.0%

Contains 2 lb of active ingredient per gallon



Listed by the Organic Materials Review Institute (OMRI) for use in organic production



For Organic Production

Keep Out of Reach of Children

Precautionary Statements

Personal Protective Equipment (PPE)

Applicators and other handlers must wear

- Long sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to bees exposed to treatment for 3 hours following treatment. Do not apply this pesticide to blooming, pollen shedding or nectar producing parts of plants if bees may forage on the plants during this time period. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on the label.

4/48

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 Refer to label booklet under Agricultural Use Requirements in the Directions for Use section for information about this standard

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water food or feed by storage and disposal

Pesticide Storage Store in original container only In case of leak or spill contain material with absorbent materials and dispose as waste

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

(Storage and Disposal for refillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water food or feed by storage and disposal

Pesticide Storage Store in original container only In case of leak or spill contain material with absorbent materials and dispose as waste

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Container Handling Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10% full with water and if possible spray all sides while adding water If practical agitate vigorously or recirculate water with the pump for two minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water food or feed by storage and disposal

Pesticide Storage Store in original container only In case of leak or spill contain material with absorbent materials and dispose as waste

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with

5/48

water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Refer to label booklet for Directions for Use

Notice Read the entire label before using Use only according to label directions **Before using this product read Warranty Disclaimer Inherent Risks of Use and Limitation of Remedies at end of label booklet If terms are unacceptable return at once unopened**

In case of emergency endangering health or the environment involving this product call 1 800 992 5994

Shake Well Before Use Avoid Freezing

EPA Reg No 62719-621

EPA Est _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268**

Net Contents ____

6/48

(Cover shipping container)

Entrust[®] SC

Naturalyte[®] Insect Control

A Naturalyte[®] insect control product formulated for control of lepidopterous larvae (worms or caterpillars), leafminers, thrips, and red imported fire ants

Group	5	INSECTICIDE
-------	---	-------------

Active Ingredient

spinosad (a mixture of spinosyn A and spinosyn D)	22.5%
---	-------

Other Ingredients	77.5%
-------------------	-------

Total	100.0%
-------	--------

Contains 2 lb of active ingredient per gallon



Listed by the Organic Materials Review Institute (OMRI) for use in organic production



For Organic Production

Keep Out of Reach of Children

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. Refer to label booklet under Agricultural Use Requirements in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use

Notice Read the entire label before using. Use only according to label directions. **Before using this product read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1 800 992 5994

Shake Well Before Use Avoid Freezing

EPA Reg No 62719 621

EPA Est _____

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis IN 46268**

Net Contents _____

(Page 1 through end)

Table of Contents	Page
Precautionary Statements	
Personal Protective Equipment (PPE)	
User Safety Recommendations	
Environmental Hazards	
Directions for Use	
Agricultural Use Requirements	
Non Agricultural Use Requirements	
Storage and Disposal	
Product Information	
Use Precautions	
Integrated Pest Management (IPM) Programs	
Insecticide Resistance Management (IRM)	
Mixing Directions	
Application Directions	
Uses	
Asparagus	
Banana and Plantain	
<i>Brassica</i> (Cole) Leafy Vegetables (Crop Group 5)	
Bulb Vegetables (Crop Group 3)	
Bushberries (Subgroup 13B)	
Caneberries (Subgroup 13A)	
Citrus (Crop Group 10)	
Corn (Field Sweet Pop and Seed) and Teosinte	
Cotton	
Cranberry	
Cucurbit Vegetables (Crop Group 9)	
Dates	
Fig	
Fruiting Vegetables (Crop Group 8) and Okra	
Grape	
Grass Crops Grass Grown for Seed Pastures and Rangeland	
Herbs (Subgroup 19A)	
Hops Dried Cones	
Leafy Vegetables (Except <i>Brassica</i>) (Crop Group 4) Leaves of Root and Tuber	
Vegetables (Crop Group 2) and Leaves of Legume Vegetables (Crop Group 7A)	
Turnip Greens and Watercress	
Legume Vegetables (Succulent and Dried Beans and Peas) (Crop Group 6)	
Nongrass Animal Feeds (Forage Fodder Straw and Hay) (Crop Group 18)	
Ornamentals (Herbaceous and Woody) Growing Outdoors in Nurseries (Including)	
Conifer Seed Orchards) or in Greenhouses	
Peanut	
Peppermint and Spearmint	
Pineapple	
Pome Fruits (Crop Group 11)	
Pomegranate	
Root and Tuber Vegetables (Crop Group 1) and Artichoke	
Small Cereal Grains and Grain Amaranth	
Soybean	
Spices (Except Black Pepper) (Subgroup 19B)	
Stone Fruits (Crop Group 12)	
Strawberry	
Tree Farms or Plantations	

Tree Nuts (Crop Group 14) and Pistachios
Tropical Tree Fruits
Fire Ants Mound Application in Turfgrass and Ornamentals
in Greenhouses and in Other Outdoor Production Areas
Terms and Conditions of Use
Warranty Disclaimer
Inherent Risks of Use
Limitation of Remedies

• •
•

9/48

Precautionary Statements

Personal Protective Equipment (PPE)

Applicators and other handlers must wear

- Long sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to bees exposed to treatment for 3 hours following treatment. Do not apply this pesticide to blooming, pollen shedding or nectar producing parts of plants if bees may forage on the plants during this time period. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on the label.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

Non Agricultural Use Requirements

10/48

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Do not enter or allow others to enter the treated area until sprays have dried

Storage and Disposal

Do not contaminate water food or feed by storage or disposal

Pesticide Storage Store in original container only In case of leak or spill contain material with absorbent materials and dispose as waste

Pesticide Disposal Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Nonrefillable containers 5 gallons or less

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Refillable containers 5 gallons or larger

Container Handling Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10% full with water and if possible spray all sides while adding water If practical agitate vigorously or recirculate water with the pump for two minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

Nonrefillable containers 5 gallons or larger

Container Handling Nonrefillable container Do not reuse or refill this container

Triple rinse or pressure rinse container (or equivalent) promptly after emptying **Triple rinse** as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times **Pressure rinse** as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities

11/48

Product Information

Entrust® SC is a Naturalyte® insect control product for control of many foliage feeding pests including lepidoptera larvae (worms or caterpillars) Colorado potato beetles leafminers and thrips infesting labeled crops This product's active ingredient spinosad is biologically derived from the fermentation of *Saccharopolyspora spinosa* a naturally occurring soil organism Mix Entrust SC with water and apply as a foliar spray with aerial or ground equipment equipped for conventional insecticide spraying

Use Precautions

Integrated Pest Management (IPM) Programs

Entrust SC is recommended for IPM programs in labeled crops Apply Entrust SC when field scouting indicates target pest densities have reached the economic threshold i.e. the point at which the insect population must be reduced to avoid economic losses beyond the cost of control Other than reducing the target pest species as a food source Entrust SC does not have a significant impact on certain parasitic insects or the natural predaceous arthropod complex in treated crops including big eyed bugs ladybird beetles flower bugs lacewings minute pirate bugs damsel bugs assassin bugs predatory mites or spiders The feeding activities of these beneficials will aid in natural control of other insects and reduce the likelihood of secondary pest outbreaks If Entrust SC is tank mixed with any insecticide that reduces its selectivity in preserving beneficial predatory insects the full benefit of Entrust SC in an IPM program may be reduced

Insecticide Resistance Management (IRM)

Entrust SC contains spinosad a Group 5 insecticide Insect/mite biotypes with acquired resistance to Group 5 insecticides may eventually dominate the insect/mite population if Group 5 insecticides are used repeatedly in the same field or area or in successive years as the primary method of control for targeted species This may result in partial or total loss of control of those species by Entrust SC or other Group 5 insecticides Currently only spinetoram and spinosad active ingredients are classified as Group 5 insecticides These two insecticide active ingredients share a common mode of action and must not be rotated with each other for control of pests listed on this label Spinetoram and spinosad may be rotated with all other labeled insecticide active ingredients

To delay development of insecticide resistance the following practices are recommended

- Carefully follow the specific label guidelines within the use directions sections of this label especially in regard to IRM recommendations
- Avoid use of the same active ingredient or mode of action (same insecticide group) on consecutive generations of insects However multiple applications to reduce a single generation are acceptable Treat the next generation with a different active ingredient that has a different mode of action or use no treatment for the next generation
- Avoid using less than labeled rates of any insecticide when applied alone or in tank mixtures
- Applications should be targeted against early insect developmental stages whenever possible
- Base insecticide use on comprehensive IPM programs including crop rotations
- Monitor treated insect populations in the field for loss of effectiveness
- Contact your local extension specialist certified crop advisor and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems
- For further information or to report suspected resistance contact your local Dow AgroSciences representative or by calling 800 258 3033

Mixing Directions

Always shake well before use Avoid freezing

Application Rate Reference Table

Application Rate of Entrust SC (fl oz/acre)	Active Ingredient Equivalent (lb ai/acre)	Acres per Gallon of Entrust SC
---	---	--------------------------------

12/48

Application Rate of Entrust SC (fl oz/acre)	Active Ingredient Equivalent (lb ai/acre)	Acres per Gallon of Entrust SC
1.5	0.023	85
3	0.047	43
4	0.062	32
6	0.094	21
8	0.125	16
10	0.156	13

Entrust SC Alone Fill the spray tank with water to about 1/2 of the required spray volume. Start agitation and add the required amount of Entrust SC. Continue agitation while mixing and filling the spray tank to the required spray volume. Maintain sufficient agitation during application to ensure uniformity of the spray mix. Do not allow water or spray mixture to back siphon into the water source.

Entrust SC Tank Mix When tank mixing Entrust SC with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. If foliar fertilizers are used, the jar test should be repeated with each batch of fertilizer utilizing the mixing water source. Do not use acidifying buffering agents in tank mixes with Entrust SC. Vigorous, continuous agitation during mixing, filling, and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes Fill the spray tank with water to 1/4 to 1/3 of the required spray volume. Start agitation. Add different formulation types in the order indicated below, allowing time for complete dispersion and mixing after addition of each product. Allow extra dispersion and mixing time for dry flowable products.

Add different formulation types in the following order:

- 1 Water dispersible granules
- 2 Wettable powders
- 3 Entrust SC and other aqueous suspensions

Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add:

- 4 Emulsifiable concentrates and water based solutions
- 5 Spray adjuvants, surfactants and oils
- 6 Foliar fertilizers

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling, and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose.

Premixing Dry and flowable formulations may be premixed with water (slurried) and added to the spray tank through a 20 to 35 mesh screen. This procedure assures good initial dispersion of these formulation types.

Spray Tank pH A spray tank pH between 6.0 and 9.0 is suggested to achieve maximum performance of Entrust SC. If the water source is outside of this pH range, or if tank mixing other pesticides, adjuvants, or foliar nutrients will cause the pH to fall outside of this range, consider adjusting the spray tank pH to be between 6.0 and 9.0 before adding Entrust SC. To do this, add all other tank mix components first, then check the spray tank pH, adjust if desired, and then add Entrust SC. If you require additional information on how to adjust spray tank pH, contact your Dow AgroSciences representative.

Use of Adjuvants Adjuvants may be used to improve the control of leafminers and thrips in situations

13/48

where achieving uniform plant coverage is difficult (such as closed crop canopy or dense foliage) or penetration into waxy leaf surfaces is required for pest control

- Use only adjuvant products labeled for agricultural use and follow the manufacturer's label directions. A nominal concentration of 1 to 2 quarts per 100 gallons (0.25 to 0.5% v/v) is generally sufficient.
- For leafminers and thrips, emulsified crop oils or methylated crop oil plus organosilicone combination products are recommended.
- When using adjuvants, always conduct a jar test to determine the compatibility of the various components in the spray mixture. Crop safety should be evaluated in a small area of the crop whenever there is a significant change in spray mixture ingredients or source of water for the spray mixture.
- Do not use diesel fuel or pure mineral oil.
- Adjuvants may require organic certification; consult your organic certifier. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Proper application techniques help ensure thorough spray coverage and correct dosage for optimum insect control. The following directions are provided for ground and aerial application of Entrust SC. Attention should be given to sprayer speed and calibration, wind speed, and foliar canopy to ensure adequate spray coverage.

Row Crop Application

Use calibrated power operated ground spray equipment capable of providing uniform coverage of the target crop. Orient the boom and nozzles to obtain uniform crop coverage. A minimum of 5 to 10 gallons per acre should be utilized, increasing volume with crop size and/or pest pressure. Use hollow cone, twin jet, flat fan nozzles or other atomizer suitable for insecticide spraying to provide a fine to coarse spray quality (per ASABE S 572; see nozzle catalogs). Under certain conditions, drop nozzles may be required to obtain complete coverage of plant surfaces. Follow manufacturer's specifications for ideal nozzle spacing and spray pressure. Minimize boom height to optimize uniformity of coverage and maximize deposition (optimize on target deposition) to reduce drift.

Orchard Spraying Application

Dilute Spray Application This application method is based upon the premise that all plant parts are thoroughly wetted to the point of runoff with spray solution. To determine the number of gallons of dilute spray required per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance.

Concentrate Spray Application Apply Entrust SC in a manner that achieves uniform coverage of the entire crop canopy but not past the point of runoff. For optimum control of target pests, complete and uniform spray coverage is essential. The spray volume required to achieve complete and uniform coverage will depend upon tree size and shape, leaf size, and density, and the application equipment used. To determine the required spray volume per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance. Use of tree row volume is appropriate.

Aerial Application

Apply in spray volume of 5 gallons or more per acre (10 gallons or more per acre for trees, vines, or orchard crops). Nozzle configuration should provide a medium to fine droplet size per ASABE S 572 standard (see USDA ARS or NAAA handbook). Guidance for ASABE S 572 nozzle configuration can be found at the following web site: www.cproductsinc.com. Boom length must be less than 75% of wing or 85% of rotor span and swath adjustment (offset) to compensate for crosswinds. Observe minimum safe application height (maximum 12 feet for ag canopies). Use GPS equipment, swath markers, or flagging to ensure proper application to the target area. Configure the boom/nozzle used (e.g., at NAAA Fly In) for both crosswind and near parallel winds. If application is made parallel to the wind direction, adjust swath

14/48

width downward. Use swath adjustment (offset) to compensate for crosswinds. Do not apply under completely calm wind conditions. It is best to apply when wind speed is between 2 to 10 mph. Under conditions of low humidity and high temperatures, adjust spray volume and droplet size upward to compensate for evaporation of spray droplets. Insect control by aerial application may be less than control by ground application because of reduced coverage.

Chemigation Application

Entrust SC may be applied through properly equipped chemigation systems for insect control in corn, cranberries, ornamentals, and potatoes. Follow use directions for these crops in the Uses section of this label. Do not apply Entrust SC by chemigation to other labeled crops except as specified in Dow AgroSciences supplemental labeling or product bulletins.

General Directions for Sprinkler Chemigation Entrust SC may be applied through overhead sprinkler irrigation systems that will apply water uniformly, including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system. Sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units are not recommended.

For continuously moving systems, the mixture containing Entrust SC must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. If continuously moving irrigation equipment is used, apply in no more than 0.25 inch of water. For irrigation systems that do not move during operation, apply in no more than 0.25 inch of irrigation immediately before the end of the irrigation cycle.

General Directions for Drip Chemigation Entrust SC may be applied through surface or buried drip systems or micro sprinklers. Drip irrigation application procedures are designed to minimize soil adsorption and maximize the bio-availability of Entrust SC to target pests. For best results, make the application in conjunction with continuous drip irrigation or a normal drip irrigation cycle as described in the following steps:

- **Pre irrigation** Moderate pre irrigation is required. Soil in the vicinity of emitters should be at or above field capacity prior to injection of Entrust SC.
- **Application Rate** Apply an amount equivalent to the labeled broadcast application rate for the labeled crop.
- **Injection** Prior to injection, fully charge the drip irrigation system and make sure it is in operation. Injection of Entrust SC should occur without interruption following pre irrigation. Mix Entrust SC in a dilution volume sufficient for a 1 to 4 hour injection period based upon the system calibration. Continuously agitate the mixture in the injection system supply tank throughout the injection cycle. Inject the diluted mixture of Entrust SC into the center of the irrigation water stream using a suitable dip tube to encourage thorough mixing and even distribution within the drip irrigation system. This is especially important if flow is slow or laminar.

Chemigation Preparation The following use directions are to be followed when this product is applied through sprinkler irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of Entrust SC needed to cover the desired acreage. Mix according to instructions in the Mixing Directions section above. Continuously agitate the mixture during mixing and application.

Chemigation Equipment Calibration In order to calibrate the irrigation system and inject or to apply the mixture containing Entrust SC, determine the following: 1) Calculate the number of acres irrigated by the system. 2) Calculate the amount of product required and premix. 3) Determine the irrigation rate and determine the number of minutes for the system to cover the intended treatment area. 4) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes (minus time to flush out) to cover the treatment

area This value equals the gallons per minute output that the injector or eductor must deliver Convert the gallons per minute to milliliters or ounces per minute if needed Calibrate the injector system with the system in operation at the desired irrigation rate It is suggested that the injection pump/system be calibrated at least twice before operation and that the system should be monitored during operation

Chemigation Operation Start the water pump and irrigation system and let the system achieve the desired pressure and speed before starting the injector Check for leaks and uniformity and make repairs before any chemigation takes place Start the injection system and calibrate according to manufacturer's specifications This procedure is necessary to deliver the desired rate per acre in a uniform manner When the application is finished allow the entire irrigation and injection system to be thoroughly flushed clean before stopping the system

Chemigation Precautions

- Lack of effectiveness or illegal pesticide residues in the crop can result from non uniform distribution of treated water
- If you have questions about calibration contact state extension service specialists equipment manufacturers or other experts
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide label prescribed safety devices for public water systems are in place with current certification Specific local regulations may apply and must be followed
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection
- Do not apply when wind speed favors drift beyond the area intended for treatment End guns must be turned off during the application if they irrigate nontarget areas
- Do not allow irrigation water to collect or run off and pose a hazard to livestock wells or adjoining crops
- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units
- Limit application of Entrust SC by drip irrigation systems to coarse textured soils with low organic matter content Product effectiveness is reduced in soils with significant clay or organic matter
- Do not tank mix Entrust SC with other pesticides or agricultural products when applying through drip irrigation systems
- If Entrust SC is applied by drip irrigation do not make broadcast foliar applications of Entrust SC during the crop cycle

Chemigation Specific Equipment Requirements

- The system must contain an air gap or approved backflow prevention device or approved functional check valve vacuum relief valve (including inspection port) and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations
The pesticide injection line must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection chemical supply
- A pesticide injection pump must also contain a functional interlock e.g. mechanical or electrical to shut off chemical supply when the irrigation system is either automatically or manually shut down
- The system must contain functional interlocking controls to automatically shut off the pesticide injection when the water pressure drops too low or water flow stops
- Use of public water supply requires approval of a backflow prevention device or air gap (preferred) by both state and local authorities
- Systems must use a metering device such as a positive displacement injection pump (or flow meter on eductor) effectively designed and constructed of materials that are compatible with pesticides and

capable of being fitted with a system interlock. An electric powered pump must meet Section 675 for Electrically Driven or Controlled Irrigation Machines, NEC 70.

- To insure uniform mixing of the insecticide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.
- The tank holding the insecticide mixture should be free of rust, fertilizer, sediment, and foreign material and equipped with an in-line strainer situated between the tank and the injection point.

Uses

Asparagus

(Post Harvest Protection of Ferns Only)

Pest and Application Rates

Pest	Entrust SC (fl oz/acre)
asparagus beetle	4 – 6

Application Timing For determining when to treat, scout with enough regularity to monitor the population size of the labeled pest. Make applications **only to asparagus ferns**. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control asparagus beetle in asparagus fern. Use a higher rate in the rate range for heavy infestations or advanced growth stages of the beetle. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval This use is only for asparagus ferns; do not apply within 60 days of spear harvest.

Minimum Treatment Interval Do not make applications less than 4 days apart.

Do not apply more than a total of 18 fl oz of Entrust SC (0.28 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not make more than three applications per crop.

Do not feed treated ferns to meat or dairy animals.

Banana and Plantain

(For use in California, Florida, Hawaii, and Texas only)

Pests and Application Rates

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (oz/100 gal)
banana rust thrips ¹	8	3.3
caterpillars		
Hawaiian flower thrips ¹		

17/48

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions. Dilute sprays assume an average volume of 300 gallons per acre.

Application Timing Apply no later than two weeks after bunch emergence and before flower petals senesce and again one to two days before bunch cover.

Application Rate Apply as a directed fine spray toward bunches and spray to runoff.

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate range for dilute sprays in the table is based upon a spray volume of 300 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 8 weeks of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not make more than four applications per crop or apply more than six times per calendar year.

Brassica (Cole) Leafy Vegetables (Crop Group 5)¹

¹Brassica (cole) leafy vegetables (crop group 5) broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens.

In the state of Georgia do not apply Entrust SC to broccoli raab, Chinese cabbage (bok choy), collards, kale, mizuna, mustard greens, mustard spinach, rape greens.

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
diamondback moth	1.5 – 4
cabbage looper imported cabbageworm	3 – 6
armyworms (including beet armyworm) leafminers ¹ thrips ²	4 – 10
flea beetle (suppression)	4 – 8

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing Treat when pests appear, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications, but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor, or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate

in the rate range for heavy infestations or advanced growth stages of target pests

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application For **diamondback moth** if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area Make treatment decisions for the entire farm and consider area wide programs if other growers are in close proximity Do not make more than six applications of Entrust SC per calendar year for diamondback moth over an entire farm (an area of abutting or nearby fields)

Restrictions

Preharvest Interval Do not apply within 1 day of harvest

Minimum Treatment Interval Do not make applications less than 4 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than six applications per calendar year

Do not apply to seedling cole crops grown for transplant within a greenhouse shade house or field plot

Bulb Vegetables (Crop Group 3)¹

¹Bulb vegetables (crop group 3) dry bulb onion garlic great headed (elephant) garlic green onion leek shallot welch onion

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms dipteran leafminers European corn borer fleabeetle loopers	3 – 6
thrips (suppression) ¹	4 – 8

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture See Use of Adjuvants under Mixing Directions If thorough coverage is desired then high pressure (>70 psi) directed sprays with dual directed nozzles can assist leaf penetration of onion

Application Timing For determining when to treat scout with enough regularity to monitor the population size of each of the labeled pests Treat when pests appear targeting eggs at hatch or small larvae Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for larger larvae or heavier infestations

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

19/48

Preharvest Interval Do not apply within 1 day of harvest

Minimum Treatment Interval Do not make applications less than 4 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than five applications per calendar year

Bushberries (Subgroup 13B)¹

(Insect Suppression)

¹Bushberries (subgroup 13B) blueberry currant elderberry gooseberry huckleberry juneberry lingonberry salal

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms	4-6
blueberry gall midge	
cherry fruitworm	
cranberry fruitworm	
currant fruitfly	
European grapevine moth	
fireworms	
leafrollers	
light brown apple moth	
loopers	
thrips ¹	

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing Treat when pests appear targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate The amount of Entrust SC applied per acre will depend upon plant size and volume of foliage present and pest pressure. Use a lower rate in the rate range for light infestations and/or small plants and a higher rate in the rate range for heavy infestations and/or larger plants.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 3 days of harvest

Minimum Treatment Interval Do not make applications less than 6 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than six applications per calendar year or more than three applications per crop

Caneberries (Subgroup 13A)¹

¹Caneberries (subgroup 13A) blackberry loganberry red and black raspberry cultivars and/or hybrids of these.

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms European grapevine moth green fruitworm leafrollers light brown apple moth raspberry fruitworm sawfly	4 - 6

Application Timing Treat when pests appear targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate The amount of Entrust SC applied per acre will depend upon plant size and volume of foliage present and pest pressure. Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 1 day of harvest.

Minimum Treatment Interval Do not make applications less than 5 days apart.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not make more than six applications per calendar year.

Citrus (Crop Group 10)¹

¹Citrus (crop group 10) grapefruit, lemons, limes, oranges, tangerines.

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
citrus leafminer citrus orangedog citrus thrips ¹ European grapevine moth light brown apple moth	4 - 10

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing Treat when pests appear or in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional area use recommendations for your area.

Application Rate The amount of Entrust SC applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or small trees and a higher rate in the rate range for heavy infestations and/or large trees.

Resistance Management Citrus thrips are present most of the time on the crop during the growing season and have demonstrated a high potential to develop resistance to insect control products. Do not

make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application For **citrus thrips** rotate to another class of effective products for the next two applications after using two applications of Entrust SC within a season Consult your Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Preharvest Interval Do not apply within 1 day of harvest

Minimum Treatment Interval Do not make applications less than 6 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than three applications per calendar year
Do not apply to citrus nurseries or citrus in greenhouses

Corn (Field Sweet Seed Popcorn) and Teosinte

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms (including fall armyworm) European corn borer	1.5 - 6
beet armyworm corn earworm southwestern corn borer western bean cutworm	3 - 6

Application Timing Scout for **European corn borer** and **armyworms** with enough regularity to monitor egg laying and egg hatch Time applications of Entrust SC to coincide with peak egg hatch of each generation Frequent treatments may be necessary when the crop is growing rapidly during silking or under heavy pest pressure For **corn earworm** control a 1 to 2 day re treatment schedule may be necessary at silking For control of all other pests a 5 to 7 day re treatment schedule may be necessary if the crop is growing rapidly or if there is heavy pest pressure

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests

Spray Delivery For control of **first generation European corn borer** and **armyworms** apply broadcast or as a directed spray into the leaf whorls For control of **corn earworm** apply broadcast or direct spray to ear zone Use sufficient spray volume and nozzle pressure to ensure thorough wetting of the silks

Chemigation Entrust SC may be applied to corn by chemigation at labeled rates Refer to the Chemigation Application section

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Sweet Corn Popcorn Seed Corn

Preharvest Interval Do not apply within 28 days of fodder harvest 1 day of grains harvest or 7 days of forage harvest

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year

Maximum Number of Applications Do not make more than six applications per calendar year
Field Corn and Teosinte

- **Preharvest Interval** Do not apply within 28 days of grain or fodder harvest or 7 days of forage harvest
Do not apply more than a total of 12 fl oz of Entrust SC (0.188 lb ai spinosad) per acre per year
- **Maximum Number of Applications** Do not make more than three applications per calendar year

Cotton

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
European corn borer cotton bollworm (pre bloom) cotton leafperforator tobacco budworm	3 6
armyworms (including beet armyworm fall armyworm) cotton bollworm (post bloom) leafminers loopers (including soybean looper cabbage looper) saltmarsh caterpillar thrips	4 6

Application Timing

Tobacco Budworm and/or Cotton Bollworm For the most effective control scout fields twice per week and apply Entrust SC when the majority of the population is within the time of blackhead egg stage to 1/8 inch larval length. The following table illustrates the size of development of worms in relation to age and stage of development (instar) as a guide to timing treatments for optimum control

Age (Days)	Average Size (Inches)	Instar¹
Hatch	1/16	1st
3	1/4	2nd
5	1/2	3rd
8	7/8	4th
10	1	5th

¹**Note** A scouting schedule of only once per week is risky since hatching worms will have grown to 3rd instar before the next scouting observation has determined the need to spray

Beet Armyworm Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply Entrust SC when field scouting reveals three or more occurrences of egg hatch or larval feeding per 100 feet of row

Loopers Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply Entrust SC when field scouting reveals four larvae per 1 foot of row or 25% defoliation

Application Rate Use a higher rate in the rate range and higher spray volume when one or more of the following is true: tobacco budworms or bollworms are more than 1/4 inch in length; target pest population

is 2X above state threshold level or foliage canopy is tall/dense and worms are present in the lower part of the canopy Heavy infestations may require repeat applications but follow resistance management guidelines

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area For **tobacco budworm** and/or **cotton bollworm** where early season conservation of beneficial insects is practical use Entrust SC to control the 1st and 3rd generation of tobacco budworm and/or cotton bollworm Where conservation of beneficial insects is not as critical (for example fields have received non selective early season treatments for boll weevil or lygus bugs) use Entrust SC to control either the 2nd or 3rd generation of tobacco budworm and/or cotton bollworm

Restrictions

- **Preharvest Interval** Do not apply within 28 days of harvest
- **Minimum Treatment Interval** Do not make applications less than 5 days apart for high rates of application
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per growing season

Cranberry

(Insect Suppression)

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms	4 - 10
currant fruitfly	
European grapevine moth	
fireworms	
leafrollers	
light brown apple moth	
loopers	
sparganothis fruitworm	
thrips	

Application Timing For determining when to treat scout with enough regularity to monitor the population size of each of the labeled pests Treat when pests appear targeting eggs at hatch or small larvae Consult your Dow AgroSciences representative extension specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area

Application Rate Application rate within the rate range will depend upon plant size and volume of foliage present and pest pressure Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume

Chemigation Entrust SC may be applied to cranberries by chemigation at labeled rates Refer to the Chemigation Application section

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

- Preharvest Interval** Do not apply within 21 days of harvest
- Minimum Treatment Interval** Do not make applications less than 7 days apart
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop
- Maximum Number of Applications** Do not make more than six applications per calendar year

Cucurbit Vegetables (Crop Group 9)¹

¹Cucurbit vegetables (crop group 9) cucumber edible gourds muskmelons (cantaloupe honeydew etc) pumpkin summer squash watermelon winter squash

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms cabbage looper melon worm pickleworm rindworms	4-8
leafminers ¹ thrips ¹	6-8

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture See Use of Adjuvants section under Mixing Directions

Application Timing Treat when pests appear targeting eggs at hatch or small larvae Heavy infestations may require repeat applications but follow resistance management guidelines Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional area local use recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

- Preharvest Interval** Do not apply within 3 days of harvest for all crops except cucumbers Do not apply within 1 day of harvest for cucumbers
- Minimum Treatment Interval** Do not make applications less than 5 days apart
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per season
- Maximum Number of Applications** Do not make more than six applications per crop

Dates

Pests and Application Rates

Pest	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
carob moth	8	2-7

25/48

Application Timing Apply Entrust SC as a foliar spray when pests appear or in accordance with local conditions. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a concentrate or dilute spray using conventional power operated spray equipment ensuring good coverage (see Orchard Spraying Application section under Application Directions).

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate for dilute sprays in the table is based upon a spray volume of 300 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than three consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after three consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 7 days of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year.

Maximum Number of Applications Do not make more than three applications of Entrust SC for carob moth control.

Fig

Pests and Application Rates

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
navel orangeworm	4 - 10	1 - 2.5

Application Timing Apply Entrust SC as a foliar spray when pests appear or in accordance with local conditions. Apply as a concentrate or dilute spray using conventional power operated spray equipment (see Orchard Spraying Application section under Application Directions). Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate The amount of Entrust SC applied per acre will depend upon tree size and volume of foliage present and pest pressure. Use a higher rate in the rate range for large trees or heavy infestations.

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate range for dilute sprays in the table is based upon a spray volume of 400 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 7 days of harvest

Minimum Treatment Interval Do not make applications less than 7 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

- **Maximum Number of Applications** Do not make more than four applications per calendar year

Fruiting Vegetables (Crop Group 8)¹ and Okra

¹Fruiting vegetables (crop group 8) eggplant groundcherry pepino pepper (except black) tomatillo tomato

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
lepidopterous larvae (maintenance only)	1.5 - 3
Colorado potato beetle European corn borer European grapevine moth hornworms light brown apple moth loopers tomato fruitworm	3 - 6
armyworms (including beet armyworm) flea beetle (suppression) flower thrips ^{1 2} thrips palmi ^{1 2} tomato pinworm	4 - 8
leafminers ¹ (<i>Linomyza</i> spp)	6 - 10

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture See Use of Adjuvants section under Mixing Directions

²For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications

Application Timing Scout weekly throughout the season to monitor and track populations of leafminers and thrips to determine when economic thresholds are exceeded Scout weekly throughout the season to monitor and track pest and beneficial populations For tracking lepidopterous larvae scout with enough regularity to monitor the population size of each of the labeled pests Time applications of Entrust SC to coincide with peak egg hatch in species without overlapping generations Consult current pest management recommendations for specific guidelines

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area Do not apply Group 5 insecticides to consecutive generations of Colorado potato beetle and do not make more than two applications per single generation of Colorado potato beetle

27/48

Restrictions

- Preharvest Interval** Do not apply within 1 day of harvest
- Minimum Treatment Interval** Do not make applications less than 4 days apart
Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop
- Maximum Number of Applications** Do not make more than six applications per calendar year
Do not apply to seedling fruiting vegetables and okra grown for transplant within a greenhouse shade house or field plot

Grape

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
climbing cutworms	4 - 8
European grapevine moth	
grape berry moth	
grape leafroller	
grape leaf skeletonizer	
light brown apple moth	
omnivorous leafroller	
orange tortrix	
redbanded leafroller	
thrips	

Application Timing Treat when pests appear targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Carefully adjust equipment and spray volume to assure thorough uniform coverage of infested parts of the crop. Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

- Preharvest Interval** Do not apply within 7 days of harvest
- Minimum Treatment Interval** Do not make applications less than 5 days apart
Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per season west of the Rocky Mountains and no more than 0.36 lb ai spinosad per acre per season east of the Rocky Mountains
- Maximum Number of Applications** Do not make more than five applications per calendar year

Grass Crops, Grass Grown for Seed, Pastures and Rangeland

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)

Pests	Entrust SC (fl oz/acre)
beet armyworm fall armyworm southern armyworm true armyworm	2 – 4

Application Timing Scout at least weekly and consider the impact of both pests and beneficials. Treat when economic thresholds are exceeded targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations applicable to your area.

Application Rate Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

Resistance Management Do not apply more than three times in any 21 day period. Whenever Entrust SC is applied up to three times in succession, follow it by no use of Entrust SC for a 21 day period or rotation to another insecticide class.

Restrictions

Preharvest Interval Do not apply within 3 days of harvest for hay or fodder. There is no preharvest interval for forage.

Do not apply more than a total of 12 fl oz of Entrust SC (0.186 lb ai spinosad) per acre per season.

Maximum Number of Applications Do not make more than six applications per season.

Herbs (Subgroup 19A)¹

(Insect Suppression)

¹Herbs (subgroup 19A) angelica balm basil borage burnet camomile catnip chervil (dried) chive chive (Chinese) cilantro cilantro (leaf) clary coriander (leaf) costmary curry (leaf) dillweed horehound hyssop lavender lemongrass lovage (leaf) marigold marjoram nasturtium parsley (dried) pennyroyal rosemary rue sage savory (summer and winter) sweet bay tansy tarragon thyme wintergreen woodruff wormwood

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms loopers thrips (suppression)	4 – 6

Application Timing For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for larger larvae or high infestations and/or larger plant volume. Heavy infestations may require repeat applications but follow resistance management guidelines.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

29/48

Restrictions

- Preharvest Interval** Do not apply within 1 day of harvest
- Minimum Treatment Interval** Do not make applications less than 5 days apart
Do not apply more than a total of 30 fl oz of Entrust SC (0.47 lb ai spinosad) per acre per crop
- Maximum Number of Applications** Do not make more than five applications per calendar year or more than three applications per crop

Hops Dried Cones

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms loopers thrips (suppression)	4 – 6

Application Timing For determining when to treat scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative extension specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for larger larvae or high infestations and/or larger plant volume. Heavy infestations may require repeat applications but follow resistance management guidelines.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

- Preharvest Interval** Do not apply within 1 day of harvest
- Minimum Treatment Interval** Do not make applications less than 5 days apart
Do not apply more than a total of 30 fl oz of Entrust SC (0.47 lb ai spinosad) per acre per crop
- Maximum Number of Applications** Do not make more than five applications per calendar year

Leafy Vegetables (Except *Brassica*) (Crop Group 4)¹, Leaves of Root and Tuber Vegetables (Crop Group 2)² and Leaves of Legume Vegetables (Crop Group 7A)³, Turnip Greens, and Watercress

¹Leafy vegetables (except *Brassica*) (crop group 4) amaranth arugula cardoon celery celtuce chervil Chinese celery Chinese spinach corn salad dandelion dock edible leaved chrysanthemum endive (swiss chard) fennel garden cress garden purslane garland chrysanthemum head lettuce leaf lettuce leafy amaranth New Zealand spinach orach parsley radicchio (red chicory) rhubarb spinach Swiss chard taramola upland cress vine spinach winter cress winter purslane yellow rocket

²Leaves of root and tuber vegetables (crop group 2) bitter cassava black salsify carrot celeriac (celery root) chicory washeen (taro) edible burdock garden beet oriental radish (daikon) parsnip radish rutabaga sugar beet sweet cassava sweet potato tanier true yam turnip turnip rooted chervil

³Leaves of legume vegetables (crop group 7A) any cultivar of bean and field pea (except soybean)

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
-------	-------------------------

30/48

Pests	Entrust SC (fl oz/acre)
diamondback moth	1 5 3
cabbage looper imported cabbage worm	3 6
armyworms (including beet armyworm)	4 8
leafminers ¹ thrips ¹	6 10

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture
See Use of Adjuvants section under Mixing Directions

Application Timing Scout at least weekly and consider the impact of both pests and beneficials. Treat when economic thresholds are exceeded targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Intervals

Leafy vegetables (including watercress) Do not apply within 1 day of harvest

Leaves of root, tuber and legume vegetables Do not apply within 3 days of harvest

Minimum Treatment Interval Do not make applications less than 4 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than six applications per year


Do not apply to seedling leafy crops grown for transplant within a greenhouse or shade house

Legume Vegetables (Succulent and Dried Beans and Peas) (Crop Group 6)¹

¹Legume vegetables (succulent and dried beans and peas) (crop group 6) adzuki bean, blackeye pea, chickpea, cowpea, crowder pea, edible pod pea, English pea, fava bean, field bean, field pea, garbanzo bean, garden pea, green pea, kidney bean, lentil, lima bean, lupins, mungbean, navy bean, pigeon pea, pinto bean, runner bean, snap bean, snow pea, sugar snap pea, tepary bean, wax bean, yardlong bean

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
European corn borer (eggs and larvae)	3 6
armyworms corn earworm loopers	4 6
leafminers ¹ thrips ¹	4 5 6



¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture
See Use of Adjuvants section under Mixing Directions

Application Timing For determining when to treat scout with enough regularity to monitor the population size of each of the labeled pests Heavy infestations may require repeat applications but follow resistance management guidelines Treat when pests appear targeting eggs at hatch or small larvae For **European corn borer** initiate when moth flights first appear and use the lower rate of the rate range to control eggs and larvae every three days before they enter the plant Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Minimum Treatment Interval Do not make applications less than 5 days apart

Maximum Number of Applications Do not make more than six applications per crop

Succulent Beans and Peas

Preharvest Interval Do not apply within 3 days of harvest

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per season

Dried Beans and Peas

Preharvest Interval Do not apply within 28 days of harvest

Do not apply more than a total of 12 fl oz of Entrust SC (0.188 lb ai spinosad) per acre per season

Do not feed forage or hay to meat or dairy animals

Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) (Crop Group 18)¹

¹Nongrass animal feeds (forage fodder straw and hay) (crop group 18) alfalfa clover crown vetch kudzu lespedeza lupin milk vetch sainfoin trefoil vetch velvet bean

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
alfalfa weevil (suppression) beet armyworm fall armyworm southern armyworm true armyworm	2 – 4

Application Timing Scout at least weekly and consider the impact of both pests and beneficials Treat when economic thresholds are exceeded targeting eggs at hatch or small larvae Heavy infestations may require repeat applications but follow resistance management guidelines Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations applicable to your area

Application Rate Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests

oo

Resistance Management Do not apply more than three times in any 21 day period Whenever Entrust SC is applied up to three times in succession this should be followed by no use of Entrust SC for a 21 day period or rotation to another insecticide class

Restrictions

Preharvest Interval Do not apply within 3 days of harvest for hay or fodder There is no preharvest interval for forage

Do not apply more than a total of 12 fl oz of Entrust SC (0 186 lb ai spinosad) per acre per season

Maximum Number of Applications Do not make more than six applications per season

Do not allow cattle to graze from treated area until spray has dried

Ornamentals (Herbaceous and Woody) Growing Outdoors, in Nurseries (Including Conifer Seed Orchards), or in Greenhouses

Pests and Application Rates

Pests	Entrust SC fl oz/gallon	Entrust SC fl oz/100 gallons	Entrust SC fl oz/acre
chrysomelid leaf feeding beetles such as elm leaf (1) viburnum leaf (larvae) willow leaf (1) European grapevine moth lepidopterous larvae such as azalea caterpillar bagworm beet armyworm cabbage looper California oakworm cankerworm diamondback moth E tent caterpillar fall webworm Florida fern caterpillar geranium budworm gypsy moth light brown apple moth oblique banded leafroller oleander caterpillar orange striped oakworm spruce budworm tussock moths (hickory whitemarked) W tent caterpillar winter moth yellownecked caterpillar (2) sawfly larvae such as European pine pear redheaded pine shore fly thrips (exposed) in greenhouse settings such as (3) Cuban laurel western flower	0.03 (0.9 mL)	3 (88.7 mL)	12 (354.9 mL)

Pests	Entrust SC fl oz/gallon	Entrust SC fl oz/100 gallons	Entrust SC fl oz/acre
dipterous gall midges pinyon spindlegall thrips (exposed) in outdoor settings such as Cuban laurel western flower (3)	0.05 (1.5 mL)	5.5 (162.7 mL)	22 (650.6 mL)
dipterous leafminers such as serpentine (4) emerald ash borer (5) lewis mites Nantucket pine tip moth spider mites such as spruce two spotted (6) (See 6 below for mite suppression/control expectations)	0.1 (3 mL)	10 (296 mL)	29 (858 mL)

Numbers in parentheses () refer to Pest Specific Use Directions

Pest Specific Use Directions (for pest control in the greenhouse or nursery also refer to Greenhouse Pest Resistance Avoidance Recommendations)

- Elm leaf beetle and willow leaf beetle** (adults and larvae) For effective control apply in the spring or early summer when feeding is observed
- For effective control of the following lepidopterous larvae
 - Bagworms** Apply when bags are small and larvae are actively feeding
 - Beet armyworms** Apply when larvae are small
 - Tent caterpillars and fall webworms** Apply early when webs are first observed and direct the spray into the web and surrounding foliage within at least 3 feet of the nest
 - Gypsy moth larvae** Apply when larvae are small and all eggs have hatched
 - Spruce budworms** Apply when larvae are exposed and actively feeding
- Exposed thrips (Cuban laurel and western flower)** For effective control apply early at first signs of infestation and repeat until infestation is controlled For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications
- Serpentine leafminers** For effective control apply early when stippling or mining of leaves is first observed and repeat until infestation is controlled Three sequential applications at 7 day intervals can maximize control Addition of a nonionic spray adjuvant such as DYNE AMIC spray adjuvant at 0.1% v/v in greenhouse settings has been shown to enhance control of leafminers (follow surfactant manufacturer's label directions)
- Apply to foliage and bark of tree when adult **emerald ash borer** are first observed emerging from the bark or when adult emerald ash borer are first noticed feeding on the leaves of the tree Reapply every 7 to 10 days until no additional adult emerald ash borer activity is observed Application to trees already heavily infested may not prevent the eventual loss of the tree due to existing pest damage and tree stress
- Spruce spider mites and two spotted spider mites** Apply when spider mites are first observed prior to webbing and before mite populations have become severe Reapply after 7 to 10 days (3 to 5 days in greenhouses and structures that can be altered to be closed or open) to contact newly hatched nymphs and repeat until infestation is managed **Uniform coverage of both upper and lower leaf surfaces is critical**

Note Control of spider mites with Entrust SC in certain research trials has been variable The variability between these evaluations is not well understood but may be due to late application timing when mite populations and webbing were severe poor spray coverage of both the upper and lower leaf surfaces or interaction of the leaf surface with residues of Entrust SC Addition of a nonionic spray adjuvant such as Activate Plus DYNE AMIC Joint Venture Phase and Thoroughbred at 0.1%

34/48

v/v in greenhouse settings and at label rates in outdoor settings has been shown to improve spray coverage and enhance control of spider mites (follow surfactant manufacturer s label directions)

Application Timing Dilute Entrust SC in water and apply using suitable hand or power operated application equipment (such as portable pump up backpack hydraulic boom) in a manner to provide complete and uniform plant coverage Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area Use of Entrust SC in lath and shadehouses is permitted

Application Rate Entrust SC may be used up to a maximum labeled rate of 0.1 fl oz per gallon (10 fl oz per 100 gallons 29 fl oz per acre) per application on trees and ornamentals as a general treatment regardless of the target insect pest Use pest specific rates when a single insect pest or group of insect pests within a rate category is the only intended target

Spray Volume Attempt to penetrate dense foliage but avoid over spraying to the point of excessive runoff Uniform coverage of both upper and lower leaf surfaces is critical for effective insect control

Chemigation Entrust SC may be applied to ornamentals by chemigation at labeled rates Refer to the Chemigation Application section

Phytotoxicity Entrust SC has been tested alone on a wide variety of herbaceous and woody ornamental plants without phytotoxic symptoms However because it is not possible to test all possible tank mix combinations (including adjuvants) and ornamental plant species varieties and cultivars and because environmental factors and varietal and plant stage of growth may affect phytotoxic expression it is recommended that a small group of test plants be treated at the anticipated use rate of Entrust SC either alone or in tank mix combinations and observed for at least 5 to 7 days to determine phytotoxicity before treating large numbers of those plants **Note** The professional user assumes responsibility for determining if Entrust SC is safe to treated plants when applied either alone or in tank mixtures under commercial growing conditions Research has demonstrated that some spotting of saintpaulia (African violet) flowers may occur

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Minimum Treatment Interval Except for greenhouses and structures that can be altered to be closed or open do not make applications less than 7 days apart
Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year

Peanut

(Not for Use in California)

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
-------	----------------------------

Pests	Entrust SC (fl oz/acre)
armyworms including beet armyworm fall armyworm true armyworm yellowstriped armyworm cabbage looper corn earworm European corn borer green cloverleaf worm red necked peanut worm saltmarsh caterpillar soybean looper tobacco budworm velvetbean caterpillar	3 - 6

Application Timing Regularly monitor the population size of each of the labeled pests. Treat when pests appear targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Use a higher rate in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

- Preharvest Interval** Do not apply within 3 days of nut harvest or within 14 days of forage.
- Minimum Treatment Interval** Do not make applications less than 7 days apart.
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop.
- Maximum Number of Applications** Do not make more than three applications per calendar year.
- Grazing Restrictions** Do not allow grazing of crop residue or harvest of crop residue for hay until 14 days after the last application.

Peppermint and Spearmint

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms cutworms dipteran leafminers ¹ loopers thrips (suppression) ¹	4 - 10

¹ Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions. Control in peppermint and spearmint has been variable. High pressure directed sprays can assist leaf penetration of peppermint and spearmint.

Application Timing For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear targeting eggs at hatch or small

36/48

larvae Consult your Dow AgroSciences representative extension specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Heavy infestations may require repeat applications but follow resistance management guidelines Use a higher rate in the rate range for heavier infestations or advanced growth stages of target pests

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Preharvest Interval Do not apply within 7 days of harvest

Minimum Treatment Interval Do not make applications less than 4 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than four applications per calendar year or more than three applications per crop

Pineapple

(Insect Suppression)

(For use in Hawaii only)

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
lepidopteran larvae such as armyworms banana moth fruit borer caterpillar (<i>Thecia basillides</i>) <i>Gummosos Batrachedra commosae</i> pineapple caterpillar pink cornworm sugarcane bud moth	4-6

Application Timing For determining when to treat scout with enough regularity to monitor the population size of each of the labeled pests Treat when pests appear targeting eggs at hatch or small larvae Consult your Dow AgroSciences representative extension specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Heavy infestations may require repeat applications but follow resistance management guidelines

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Preharvest Interval Do not apply within 7 days of harvest

Minimum Treatment Interval Do not make applications less than 7 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year

Maximum Number of Applications Do not make more than six applications per calendar year

Pome Fruits (Crop Group 11)¹

¹Pome fruits (crop group 11) apples crabapple loquat mayhaw pears quince

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
leafminers ¹ spotted tentiform western tentiform	4 - 10
apple maggot (suppression) codling moth European grapevine moth leafrollers oblique banded pandemis light brown apple moth oriental fruit moth thrips ¹ tufted apple budmoth	6 - 10

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture
See Use of Adjuvants section under Mixing Directions

Application Timing Optimal timing for **leafminers** and **leafrollers** may vary between species and geographic location. For **leafminers** monitor the moth flights and infestation densities of both the sap feeding and tissue feeding stage. For optimum control treat at first appearance of leaf mining activity. For **leafrollers** monitor the moth flights and the infestation densities of the larval stages. Repeat application as necessary to maintain control. Closely follow regional spray recommendations for **codling moth** and **oriental fruit moth** treatments based upon biofix dates and pheromone trap catches. **Codling moth** and **oriental fruit moth** larvae must be controlled before they penetrate the fruit. **Codling moth** and **oriental fruit moth** applications will provide control for no more than 10 days. Repeat application as necessary to maintain control. Consult with your Dow AgroSciences representative, state agricultural experiment station, certified pest control advisor or extension specialist for specific application timings in your area.

Application Rate The amount of Entrust SC applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or small trees and a higher rate in the rate range for heavy infestations and/or larger trees.

Resistance Management Do not make more than three consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after three consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Do not treat consecutive generations of codling moth, oriental fruit moth and leafrollers.

Restrictions

Preharvest Interval Do not apply within 7 days of harvest

Minimum Treatment Interval Do not make applications less than 10 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Do not apply more than three sprays targeted at leafrollers per season

Maximum Number of Applications Do not make more than four applications per calendar year

Pomegranate

Pests and Application Rates

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
carob moth	4 – 8	13 – 27
filbert moth		
leafrollers such as		
oblique banded		
omnivorous		
fruit tree		
pandemis		
redbanded		
variegated		
light brown apple moth		
navel orangeworm		
oriental fruit moth		
peach twig borer		
thrips ¹		
western cherry fruit fly		

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing

Application Timing Optimal timing for leafrollers may vary between species and geographic location. Monitor the moth flights and the infestation densities of the larval stages. Thorough coverage is necessary for optimal control. Consult with your Dow AgroSciences representative, state agricultural experiment station, certified pest control advisor or extension specialist for specific application timings in your area.

Application Rate Use a higher rate in the rate range for large trees, heavy infestations, or advanced growth stages of target pest, especially if spray volume or coverage is marginal.

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate range in the table is based upon a spray volume of 300 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than three consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after three consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid treating consecutive generations of oriental fruit moth and leafrollers.

Restrictions

Preharvest Interval Do not apply within 7 days of harvest

Minimum Treatment Interval A 10 to 14 day re-treatment schedule may be necessary to maintain control if the crop is growing rapidly or if there is heavy pest pressure.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year.

Maximum Number of Applications Do not apply more than three sprays targeted at leafrollers per season.

Root and Tuber Vegetables (Crop Group 1)¹ and Artichoke

¹Root and tuber vegetables (crop group 1) arracacha arrowroot bitter cassava black salsify carrot celeriac chayote root chicory Chinese artichoke chufa dasheen edible burdock edible canna garden beet ginger ginseng horseradish Jerusalem artichoke leren oriental radish parsnip potato radish rutabaga salsify skirret Spanish salsify sugar beet sweet cassava sweet potato tanier true yam turmeric turnip turnip rooted chervil turnip rooted parsley yam bean

Pests and Application Rates

Crops	Pests	Entrust SC (fl oz/acre)
black salsify carrot chicory ginseng horseradish parsnip salsify skirret Spanish salsify turnip rooted chervil turnip rooted parsley	armyworms dipteran leafminers European corn borer fleabeetle loopers thrips ¹	3 6
celeriac edible burdock oriental radish radish rutabaga turnip		
arracacha arrowroot artichoke bitter cassava chayote root Chinese artichoke chufa dasheen edible canna garden beet ginger Jerusalem artichoke leren potato sugar beet sweet cassava sweet potato tanier true yam turmeric yam bean	Colorado potato beetle European corn borer armyworms artichoke plume moth dipteran leafminers (<i>Linomyza</i>) loopers thrips ¹	3 10 4 5 10

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. When plants are growing rapidly, repeat applications may be necessary to protect new foliage. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or your state agricultural experiment station for any additional local use recommendations for your area.

40/48

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for larger larvae or heavier infestations. Heavy infestations may require repeat applications but follow resistance management guidelines.

Chemigation Entrust SC may be applied to potatoes by chemigation at labeled rates. Refer to the Chemigation Application section.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Do not apply Entrust SC to consecutive generations of **Colorado potato beetle** and do not make more than two applications per single generation of Colorado potato beetle.

Restrictions

Garden beet and sugar beet

Preharvest Interval Do not apply within 3 days of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 21 fl oz of Entrust SC (0.33 lb ai spinosad) per crop.

Maximum Number of Applications Do not make more than four applications per crop.

Black salsify, carrot, chicory, ginseng, horseradish, parsnip, salsify, skirret, Spanish salsify, turnip, rooted chervil, turnip, rooted parsley

Preharvest Interval Do not apply within 3 days of harvest.

Minimum Treatment Interval Do not make applications less than 5 days apart.

Do not apply more than a total of 21 fl oz of Entrust SC (0.3 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not make more than four applications per calendar year.

Arracacha, arrowroot, bitter cassava, chayote root, Chinese artichoke, chufa, dasheen, edible canna, ginger, Jerusalem artichoke, leren, potato, sweet cassava, sweet potato, tanager, true yam, turmeric, yam bean

Preharvest Interval Do not apply within 7 days of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 21 fl oz of Entrust SC (0.33 lb ai spinosad) per crop.

Maximum Number of Applications Do not make more than four applications per crop.

Artichoke

Preharvest Interval Do not apply within 2 days of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 7.5 oz of Entrust SC (0.33 lb ai spinosad) per crop.

Maximum Number of Applications Do not make more than four applications per crop.

Celeriac, edible burdock, Oriental radish, radish, rutabaga, turnip and other root vegetables not specifically listed

Preharvest Interval Do not apply within 3 days of harvest.

Minimum Treatment Interval Do not make applications less than 5 days apart.

Do not apply more than a total of 18 fl oz of Entrust SC (0.28 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not make more than three applications per calendar year.

Small Cereal Grains and Grain Amaranth

Small cereal grains: barley, buckwheat, milo, oats, pearl millet, proso millet, rye, sorghum, triticale, wheat.

Pests and Application Rates

41/48

Pests	Entrust SC (fl oz/acre)
cereal leaf beetle	2 6
armyworms	3 6
corn earworm (headworm)	
grasshoppers (suppression)	
southwestern corn borer	

Application Timing Scout for **armyworms** and **grasshoppers** with enough regularity to monitor egg laying and egg hatch and treat when thresholds are reached. Time applications of Entrust SC to coincide with peak egg hatch and/or small larval stage of growth of each generation.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations, advanced growth stages of target pests, or difficult spray coverage situations.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 21 days of grain or straw harvest or within 3 days of forage, fodder, or hay harvest.

Minimum Treatment Interval Do not make applications less than 4 days apart.

- Do not apply more than a total of 19 fl oz of Entrust SC (0.28 lb ai spinosad) per acre per year.
- **Maximum Number of Applications** Do not make more than three applications per calendar year. Do not allow cattle to graze treated area until spray has dried.

Soybean

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
green clover worm	2 25 3
soybean looper	
true armyworm	
velvet bean caterpillar	
armyworms such as	3 4
beet armyworm	
fall armyworm	
yellowstriped armyworm	
corn earworm (podworm)	
marsh caterpillar	

Application Timing Treat when field counts or crop injury indicates damaging pest populations are present or developing. Time applications to treat small larvae and use sufficient spray volume to ensure good coverage.

Application Rate Use a higher rate in rate range for heavy infestations and/or difficult spray coverage situations.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of

42/48

Group 5 insecticides rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

- **Preharvest Interval** Do not apply within 28 days of harvest
- **Minimum Treatment Interval** Do not make applications less than 4 days apart
- Do not apply more than a total of 10 fl oz of Entrust SC (0.186 lb ai spinosad) per acre per year
- **Maximum Number of Applications** Do not make more than four applications per calendar year
- Do not feed treated forage or hay to meat or dairy animals

Spices (Except Black Pepper) (Subgroup 19B)¹

¹Spices (except black pepper) (subgroup 19B) allspice, anise (seed), annatto (seed), black caraway, caper (buds), caraway, cardamom, cassia (buds), celery (seed), cinnamon, clove (buds), common fennel, coriander (seed), culantro (seed), cumin, dill (seed), Florence fennel (seed), fenugreek, grains of paradise, juniper (berry), lovage (seed), mace, mustard (seed), nutmeg, poppy (seed), saffron, star anise, vanilla, white pepper

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
lepidopteran larvae	4 - 6
flea beetles dipteran leafminers ¹ thrips ¹	6 - 10

¹Suppression of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing Directions

Application Timing For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for larger larvae or heavy infestations. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

- **Preharvest Interval** Do not apply within 14 days of harvest
- **Minimum Treatment Interval** Do not make applications less than 10 days apart
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai of spinosad) per acre per crop
- **Maximum Number of Applications** Do not make more than five applications per calendar year

Stone Fruits (Crop Group 12)¹

¹Stone fruits (crop group 12) apricot, cherries, nectarine, peach, plum, prune

Pests and Application Rates

43/48

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
cherry fruit fly such as black cherry eastern cherry western cherry fruit fly (suppression) European grapevine moth green fruitworm lepidopteran leafminers ¹ such as spotted tentiform western tentiform leafrollers such as fruit tree oblique banded pandemis redbanded variegated light brown apple moth oriental fruit moth peach twig borer thrips ¹ western cherry fruit fly	4 - 8	1.3 - 2.7

¹Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing Peach twig borer applications can be made dormant, delayed dormant or as summer sprays. Optimal timing for lepidopteran leafminers and leafrollers may vary between species and geographic location. For lepidopteran leafminers, monitor the moth flights and the infestation densities of both the sap feeding and tissue feeding stages, but for optimal control, treat before significant tissue feeding miners are observed. For leafrollers, monitor the moth flights and the infestation densities of the larval stages and re-treat as necessary to maintain control; thorough coverage is necessary for optimal control. For cherry fruit fly, western cherry fruit fly and other related species, maintain protective sprays at 7 day intervals while adults are present and fruit is susceptible to attack. For oriental fruit moth, no more than 10 days of residual control can be expected. If longer residual is required, make a second application of Entrust SC or other insecticide labeled for oriental fruit moth. For thrips, a 3 to 4 day re-treatment schedule may be necessary at flowering. After flowering, a 5 to 7 day re-treatment schedule may be followed. For cherry fruit fly, maintain protective sprays at 7 day intervals while adults are present and fruit is susceptible to attack. For all pests, consult with your Dow AgroSciences representative, state agricultural experiment station, certified pest control advisor or extension specialist for specific application timings in your area.

Application Rate Use a higher rate in the rate range for large trees, heavy infestations, or advanced growth stages of target pest, especially if spray volume or coverage is marginal.

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate range in the table is based upon a spray volume of 300 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than three consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after three consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist.

44/48

certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area Do not treat consecutive generations of oriental fruit moth and leafrollers

Restrictions

Preharvest Interval Do not apply within 14 days of harvest for apricots within 7 days of harvest for cherries plums prunes and other stone fruit crops or within 1 day of harvest for nectarines and peaches

Minimum Treatment Interval Do not make applications less than 7 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year

Maximum Number of Applications Do not apply more than three sprays targeted at leafrollers per season

Strawberry

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
armyworms including beet armyworms European grapevine moth leafrollers light brown apple moth thrips ¹	4-6

¹For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications

Application Timing Treat when pests appear targeting eggs at hatch or small larvae For **thrips** a 3 to 4 day re treatment schedule may be necessary if there is heavy pest pressure or if the pest population is increasing rapidly For control of all other pests a 5 to 7 day re treatment schedule may be necessary if the crop is growing rapidly or if there is heavy pest pressure Consult your Dow AgroSciences representative extension service specialist certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area

Application Rate Apply as a foliar spray at the rate specified to control target pests Use a higher rate in the rate range for larger larvae or moderate to severe pest infestations Heavy infestations may require repeat applications but follow resistance management guidelines

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application For thrips if additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least two applications Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

Preharvest Interval Do not apply within 1 day of harvest

Minimum Treatment Interval Do not make applications less than 5 days apart

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop

Maximum Number of Applications Do not make more than five applications per year

Tree Farms or Plantations

Conifers including Christmas trees and deciduous trees

Pests and Application Rates

45/48

Pests	Entrust SC (fl oz/acre)
European grapevine moth lepidopterous larvae such as bagworm fall webworm gypsy moth hemlock looper jackpine budworm pine tip moth redhumped caterpillar spruce budworm tent caterpillar tussock moths light brown apple moth sawfly larvae such as European pine pear redheaded pine	2 8

Application Timing Time applications to reach larvae when small or just hatching Repeat application as necessary to maintain control but follow resistance management guidelines Consult with your Dow AgroSciences representative state agricultural experiment station certified pest control advisor or extension specialist for information on application timing for specific pests in your area

Application Rates The rate of Entrust SC applied per acre will depend upon tree size and severity of infestation Use a higher rate in the rate range for large trees or heavy infestations Apply in sufficient volume to ensure thorough coverage

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad) If additional treatments are required after two consecutive applications of Group 5 insecticides rotate to another class of effective insecticides for at least one application Consult your local Dow AgroSciences representative extension specialist certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions

- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per year

Tree Nuts (Crop Group 14)¹ and Pistachios

¹Tree nuts (crop group 14) almonds cashew chestnut filbert (hazelnut) macadamia nut pecan walnut

Pests and Application Rates

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
codling moth European grapevine moth fall webworm filbert worm hickory shuckworm light brown apple moth navel orange worm olive banded leafroller peach twig borer pecan nut casebearer	4 10	1 2.5

Pests	Entrust SC	
	(fl oz/acre)	Dilute Spray (fl oz/100 gal)
redhumped caterpillar walnut caterpillar walnut husk fly		

Application Timing Apply Entrust SC as either a dormant or a foliar spray when pests appear or in accordance with local conditions. Apply as a concentrate or dilute spray using conventional power operated spray equipment (see Orchard Spraying Application section under Application Directions). Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Use of Crop Oils Crop oils labeled for agricultural use may be added to the dormant spray solution for suppression of overwintering mites and scale insects. Consult specific oil labels and University of California recommendations for precautions and restrictions regarding the use of oils in nut and fruit trees.

Application Rate The amount of Entrust SC applied per acre will depend upon tree size and volume of foliage present and pest pressure. Use a higher rate in the rate range for large trees or heavy infestations.

Spray Volume Dilute sprays are sprayed to the point of runoff. The application rate range in the table is based upon a spray volume of 400 gallons per acre. Gallonage of dilute sprays will vary depending upon tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management Do not make more than three consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after three consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Interval Do not apply within 1 day of harvest of all tree nuts and pistachios.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop.

Maximum Number of Applications Do not apply more than three sprays targeted at leafrollers per season.

Tropical Tree Fruits¹

(Insect Suppression)

¹Tropical tree fruits: acerola, atemoya, avocado, biriba, black sapote, canistel, cherimoya, custard apple, feijoa, guava, ilama, jaboticaba, longan, lychee, mamey sapote, mango, papaya, passionfruit, pulasan, rambutan, sapodilla, soursop, Spanish lime, star apple, starfruit, sugar apple, ti leaves, wax jambu (wax apple), white sapote.

Pests and Application Rates

Pests	Entrust SC (fl oz/acre)
-------	----------------------------

Pests	Entrust SC (fl oz/acre)
European grapevine moth katydids lepidopterous larvae avocado leafroller citrus peelminer cutworms fruit tree leafroller naval orange worm orange tortrix western tussock moth light brown apple moth thrips ¹	4 - 10

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing Treat when pests appear or in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate The amount of Entrust SC applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or small trees and a higher rate in the rate range for heavy infestations and/or large trees.

Resistance Management Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions

Preharvest Intervals Do not apply within 1 day of harvest.

Minimum Treatment Interval Do not make applications less than 7 days apart.

Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop.

Maximum Number of Applications In order to prevent or delay resistance development in thrips, do not apply Entrust SC more than two times per year.

For resistance management purposes, do not apply to tropical tree fruits grown in nurseries or in greenhouses.

Fire Ants – Mound Application in Turfgrass and Ornamentals, in Greenhouses, and in Other Outdoor Production Areas

Dilution Rate	
Entrust SC fl oz per 1 gallon (mL/gal)	Entrust SC fl oz per 10 gallons (mL/10 gal)
0.05 (1.5)	0.5 (14.8)

Fire ants such as imported Apply diluted Entrust SC to individual fire ant mounds as a drench application. Use 1 to 2 gallons per mound depending upon the mound size. For mounds less than 8 inches in diameter, use 1 gallon of dilution per mound. Use a higher volume, up to 2 gallons, on mounds 8 inches or larger in diameter. Apply approximately 10% of the dilution volume around the perimeter of the mound out to about 12 inches and pour the remaining volume directly on the mound. Do not disturb

48/48

mounds prior to application. If possible, apply following a recent rainfall. For best results, apply in cool weather, 65 to 85 F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays should not be used as they may disturb the ants and cause migration, reducing control.

Terms and Conditions of Use

To the extent permitted by law, if terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies.

Warranty Disclaimer

To the extent permitted by law, Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. **TO THE EXTENT PERMITTED BY LAW, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®™ Trademark of Dow AgroSciences LLC
EPA accepted ___/___/___