



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

62719-681

Date of Issuance:

2/6/2015

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

GF-3052 WG

Name and Address of Registrant (include ZIP Code):

Jamey Thomas
Dow AgroSciences LLC
9330 Zionsville, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate &Vertebrate Branch 3, Registration Division (7505P)

Date:

2/6/2015

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 62719-681.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 1/31/14

If you have any questions, you may contact Dr. Jennifer Urbanski at 703-347-0156 or via email at urbanski.jennifer@epa.gov.

Venus Eagle, Product Manager 01
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)

Enclosure

(Base label):

RESTRICTED USE PESTICIDE
 For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

GF-3052 WG

INSECTICIDE


For control or suppression of aphids, leaf feeding beetles, lepidopterous larvae (worms, caterpillars), dipterous leafminers, foliage feeding worms, lacebugs, plant bugs, thrips, whiteflies and certain scales and psyllids in *Brassica* (cole) crops, bulb vegetables, cereal grains (except rice, millet and sorghum), citrus, cotton, cucurbits, fruiting vegetables (tomato, peppers, and eggplant), grape, leafy vegetables, leaves of root and tuber vegetables, legume vegetables (succulent and dried beans and peas), okra, pistachios, root and tuber vegetables, soybean, stone fruits, strawberry, tree nuts, triticale and wheat.

Group	4C & 5	INSECTICIDE
-------	--------	-------------

Active Ingredient:

spinetoram (a mixture of spinetoram-J and spinetoram-L)	10%
sulfoxaflor	30%
Other Ingredients	60%
Total	100.0%

Contains 40% active ingredient on a weight basis



Keep Out of Reach of Children

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

First Aid
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

Restricted Use Pesticide due to toxicity categories. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Causes irreversible eye damage. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Protective eyewear (goggles or face shield)
- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is highly toxic to bees exposed through contact during spraying and while spray droplets are still wet. This product may be toxic to bees exposed to treated foliage for up to 3 hours following application. Toxicity is reduced when spray droplets are dry.

Risk to managed bees and native pollinators from contact with pesticide spray or residues can be minimized when applications are made before 7:00 am or after 7:00 pm local time or when the temperature is below 55° F at the site of application. Do not apply this pesticide to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period.

Refer to the Directions for Use for crop specific restrictions and additional advisory statements to protect pollinators.

This product is toxic to aquatic invertebrates. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as specified on the label and according to Directions for Use. See the attached label booklet.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Nonrefillable rigid containers 5 gallons or less:**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the

rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feed, drugs or clothing.

EPA Reg. No. 62719-XXX

EPA Est. _____

TM® Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

NET WEIGHT __

(booklet cover, shipping container):

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

GF-3052 WG

INSECTICIDE

For control or suppression of aphids, leaf feeding beetles, lepidopterous larvae (worms, caterpillars), dipterous leafminers, foliage feeding worms, lacebugs, plant bugs, thrips, whiteflies and certain scales and psyllids in *Brassica* (cole) crops, bulb vegetables, cereal grains (except rice, millet and sorghum), citrus, cotton, cucurbits, fruiting vegetables (tomato, peppers, and eggplant), grape, leafy vegetables, leaves of root and tuber vegetables, legume vegetables (succulent and dried beans and peas), okra, pistachios, root and tuber vegetables, soybean, stone fruits, strawberry, tree nuts, triticale and wheat.

Active Ingredient:

spinetoram (a mixture of spinetoram-J and spinetoram-L)	10%
sulfoxaflor	30%
Other Ingredients	60%
Total	100.0%

Contains 40% active ingredient on a weight basis

Keep Out of Reach of Children

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feed, drugs or clothing.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

NET WEIGHT ___

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Restricted Use Pesticide due to toxicity categories. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Causes irreversible eye damage. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Protective eyewear (goggles or face shield)
- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and out on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is highly toxic to bees exposed through contact during spraying and while spray droplets are still wet. This product may be toxic to bees exposed to treated foliage for up to 3 hours following application. Toxicity is reduced when spray droplets are dry.

Risk to managed bees and native pollinators from contact with pesticide spray or residues can be minimized when applications are made before 7:00 am or after 7:00 pm local time or when the

temperature is below 55° F at the site of application. Do not apply this pesticide to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period.

Refer to the Directions for Use for crop specific restrictions and additional advisory statements to protect pollinators.

This product is toxic to aquatic invertebrates. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as specified on the label and according to the Directions for Use in this booklet.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Protective eyewear (goggles or faceshield)
- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty

the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

GF-3052 WG insecticide is used for control or suppression of many listed foliage feeding pests including aphids, lepidopterous larvae (worms or caterpillars), whiteflies, thrips, Colorado potato beetles, dipterous leafminers, and certain psyllids infesting labeled crops. Mix the suspension concentrate of GF-3052 WG with water and apply as a foliar spray with aerial or ground equipment suitable for conventional insecticide spraying.

Use Precautions**Integrated Pest Management (IPM) Programs**

GF-3052 WG is recommended for IPM programs in labeled crops. Apply GF-3052 WG when field scouting indicates target pest densities have reached the economic threshold, i.e., the point at which the insect population must be reduced to avoid economic losses beyond the cost of control. Other than reducing the target pest species as a food source, GF-3052 WG does not have a significant impact on certain parasitic insects or the natural predaceous arthropod complex in treated crops, including big-eyed bugs, ladybird beetles, flower bugs, lacewings, minute pirate bugs, damsel bugs, assassin bugs, predatory mites or spiders. The feeding activities of these beneficials will aid in natural control of other insects and reduce the likelihood of secondary pest outbreaks. If GF-3052 WG is tank mixed with any

insecticide that reduces its selectivity in preserving beneficial predatory insects, the full benefit of GF-3052 WG in an IPM program may be reduced.

Insecticide Resistance Management (IRM)

GF-3052 WG contains sulfoxaflor, a Group 4C insecticide, and spinetoram, a Group 5 insecticide. Insect biotypes with acquired resistance to Group 4C and 5 insecticides may eventually dominate the insect population if Group 4C and 5 insecticides are used repeatedly in the same field or area, or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by GF-3052 WG. Currently, only spinetoram and spinosad active ingredients are classified as Group 5 insecticides. These two insecticide active ingredients share a common mode of action and must not be rotated with each other for control of pests listed on this label. Spinetoram and spinosad may be rotated with all other labeled insecticide active ingredients.

To delay development of insecticide resistance, the following practices are recommended:

- Carefully follow the specific label guidelines within the Use Direction sections of this label, especially in regard to IRM recommendations.
- Avoid use of the same active ingredient or mode of action (same insecticide group) on consecutive generations of insects. However, multiple applications to reduce a single generation are acceptable. Treat the next generation with a different active ingredient that has a different mode of action, or use no treatment for the next generation.
- Avoid using less than labeled rates of any insecticide when applied alone or in tank mixtures.
- Applications should be targeted against early insect developmental stages whenever possible.
- Base insecticide use on comprehensive IPM programs including crop rotations.
- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problem.
- For further information or to report suspected resistance, contact your local Dow AgroSciences by calling 800-258-3033.

Use Restrictions

- Do not treat pets.
- Do not graze livestock in treated areas.
- Do not apply directly to fish pools and other bodies of water that may contain fish.
- Do not apply to seedlings of edible crops for transplanting or to any other stage of edible crops growing in greenhouses
- Do not apply GF-3052 WG in greenhouses or other enclosed structures used for growing crops.

Mixing Directions

Application Rate Chart for Crop Uses

Application Rate of GF-3052 WG (oz/acre)	Total Active Ingredient Equivalent (lb ai/acre)	Sulfoxaflor lb ai/acre	Spinetoram lb ai/acre
5.5	0.14	0.103	0.034
5	0.13	0.094	0.031
4.5	0.11	0.084	0.028
4	0.10	0.075	0.025
3.5	0.09	0.066	0.022
3	0.08	0.056	0.019
2.5	0.06	0.047	0.016
2	0.05	0.038	0.013

1.5	0.038	0.028	0.009
1	0.025	0.019	0.006

GF-3052 WG - Alone: Fill the spray tank with water to about 1/2 of the required spray volume. Start agitation and add the required amount of GF-3052 WG. Continue agitation while mixing and filling the spray tank to the required spray volume. Maintain sufficient agitation during application to ensure uniformity of the spray mix. Do not allow water or spray mixture to back-siphon into the water source.

GF-3052 WG - Tank Mix: When tank mixing GF-3052 WG with other materials, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. If foliar fertilizers are used, the jar test should be repeated with each batch of fertilizer utilizing the mixing water source. Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes: Fill the spray tank with water to 1/4 to 1/3 of the required spray volume. Start agitation. Add different formulation types in the order indicated below, allowing time for complete dispersion and mixing after addition of each product. Allow extra dispersion and mixing time for dry flowable products.

Add different formulation types in the following order:

1. GF-3052 WG and other water dispersible granules
2. Wettable powders

Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add:

1. Emulsifiable concentrates and water-based solutions
2. Spray adjuvants, surfactants, and oils
3. Foliar fertilizers

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be re-suspended before spraying is resumed. A sparger agitator is particularly useful for this purpose.

Premixing: Dry and flowable formulations may be premixed with water (slurried) and added to the spray tank through a 20 to 35 mesh screen. This procedure assures good initial dispersion of these formulation types.

Spray Tank pH: A spray tank pH between 5.0 and 9.0 is suggested to achieve maximum performance of GF-3052 WG. If the water source is outside of this pH range, or tank mixing other pesticides, adjuvants, or foliar nutrients will cause the pH to fall outside this range, consider adjusting the spray tank pH to be between 5.0 and 9.0 before adding GF-3052 WG. To do this, add all other tank mix components first, then check the spray tank pH and adjust if desired, and then add GF-3052 WG. If you require additional information on how to adjust spray tank pH, contact your Dow AgroSciences representative.

Use of Adjuvants: The addition of agricultural adjuvants to sprays of GF-3052 WG may improve initial spray deposits, plant coverage, penetration into waxy leaf surfaces, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions.

- Use only adjuvant products labeled for agricultural use and follow the manufacturer's label directions. A nominal concentration of 1 to 2 quarts per 100 gallons (0.25 to 0.5% v/v) is generally sufficient.
- For dipterous leafminers and thrips, emulsified crop oils or methylated crop oil plus organosilicone combination products are recommended.

- For lepidopterous leafminers, thrips, and psyllids, citrus oils or horticultural oils may improve control.
- When using adjuvants, always conduct a jar test to determine the compatibility of the various components in the spray mixture. Crop safety should be evaluated in a small area of the crop whenever there is a significant change in spray mixture ingredients or source of water for the spray mixture.
- Do not use diesel fuel or pure mineral oil.
- When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Proper application techniques help ensure thorough spray coverage and correct dosage for optimum insect control. The following directions are provided for ground and aerial application of GF-3052 WG. Attention should be given to sprayer speed and calibration, wind speed, and foliar canopy density to ensure adequate spray coverage.

SPRAY DRIFT MANAGEMENT

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph. Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASABE Standard S-572 definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size. Exceptions may be indicated for specific crop groups.

Ground Row Crop Application

Use calibrated power-operated ground spray equipment capable of providing uniform coverage of the target crop. Orient the boom and nozzles to obtain uniform crop coverage. A minimum of 5 to 10 gallons per acre should be utilized, increasing volume with crop size and/or pest pressure. Use hollow cone, twin jet flat fan nozzles or other atomizer suitable for insecticide spraying to provide a fine to coarse spray quality (per ASABE S-572, see nozzle catalogs). Under certain conditions, drop nozzles may be required to obtain complete coverage of plant surfaces. Follow manufacturer's specifications for ideal nozzle spacing and spray pressure. Minimize boom height to optimize uniformity of coverage and optimize deposition (on-target deposition) to reduce drift.

Tree and Vine Crop Application

Dilute Spray Application: This application method is based upon the premise that all plant parts are thoroughly wetted, to the point of runoff, with spray solution. To determine the number of gallons of dilute spray required per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance. Use of tree row volume is appropriate.

Apply GF-3052 WG in a manner that achieves uniform coverage of the entire crop canopy but not past the point of runoff. For optimum control of target pests, complete and uniform spray coverage is essential. The spray volume required to achieve complete and uniform coverage will depend upon tree size and shape, leaf size, and density, and the application equipment used. To determine the required spray

volume per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance. Use of tree row volume is appropriate.

Groundboom Application

For groundboom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy. For ground boom and airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top of the crop canopy, spray must be directed into the canopy. Calibrate airblast application equipment and operate in a manner that achieves full displacement of the air within the crop canopy with air containing spray droplets.

Aerial Application

Apply in spray volume of 3 to 5 gallons or more per acre (10 gallons or more per acre for trees, vines, grove, or orchard crops). Nozzle configuration should provide a medium to fine droplet size per ASABE S-572 standard (see USDA-ARS or NAAA handbook). Guidance for ASABE S-572 nozzle configuration can be found at the following web site: <http://apmru.usda.gov/downloads/downloads.htm>. Boom length must be less than 75% of wing or 85% of rotor span and swath adjustment (offset) to compensate for crosswinds. Observe minimum safe application height (maximum 12 feet for agricultural canopies). Use GPS equipment, swath markers or flagging to ensure proper application to the target area. Configure the boom nozzle used (e.g., at NAAA/ Operation Safe Fly-In) for both crosswind and near parallel winds. If application is made parallel to the wind direction, adjust swath width downward. Use swath adjustment (offset) to compensate for crosswinds. Do not apply under completely calm wind conditions. It is best to apply when wind speed is between 2 to 10 mph. Under conditions of low humidity and high temperatures, adjust spray volume and droplet size upward to compensate for evaporation of spray droplets. In tree crops, insect control by aerial application may be less than control by ground application because of the reduced coverage.

Additional Requirements for Aerial Applications: Mount the spray boom on the aircraft to minimize drift caused by wingtip or rotor vortices. Use the minimum practical boom length and do not exceed 75% of the wing span or 80% rotor diameter. Flight speed and nozzle orientation must be considered in determining droplet size. Spray must be released at the lowest height consistent with the pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Chemigation Application

GF-3052 WG may be applied through properly equipped chemigation systems for insect control in potatoes. Follow use directions for these crops in the Uses section of this label. Do not apply GF-3052 WG by chemigation to other labeled crops except as specified in Dow AgroSciences state-specific 24(c) labeling.

Directions for Sprinkler Chemigation: Apply this product only through sprinkler irrigation systems including: center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move. Do not apply this product through any other type of irrigation system. Sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units are not recommended.

For continuously moving systems, the mixture containing GF-3052 WG must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. If continuously moving irrigation equipment is used, apply in no more than 0.25 inch of water. For irrigation systems that do not move during operation, apply in no more than 0.25 inch of irrigation immediately before the end of the irrigation cycle.

Chemigation Equipment Preparation: The following use directions are to be followed when this product is applied through irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of GF-3052 WG needed to cover

the desired acreage. Mix according to instructions in the Mixing Directions section above using a dilution concentrate matching your injector system requirements. Continually agitate the mixture during mixing and application.

Chemigation Equipment Calibration: In order to calibrate the irrigation system and injector to apply the mixture containing GF-3052 WG, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Calculate the amount of product required and premix. 3) Determine the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 4) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes (minus time to flush out) to cover the treatment area. This value equals the gallons per minute output that the injector or eductor must deliver. Convert the gallons per minute to milliliters or ounces per minute if needed. 5) Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the injector pump/system be calibrated at least twice before operation, and the system should be monitored during operation.

Chemigation Equipment Requirements:

- The system must contain an air gap, an approved backflow prevention device, a functional check valve, vacuum relief valve (including inspection port), and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- To insure uniform mixing of the insecticide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all back-flow prevention devices on the water line.
- The tank holding the insecticide mixture should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector point.

Chemigation Operation: Start the water pump and irrigation system, and let the system achieve the desired pressure and speed before starting the injector. Check for leaks and uniformity and make repairs before any chemigation takes place. Start the injection system and calibrate according to manufacturer's specifications. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

Chemigation Precautions:

- Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers or other experts.

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.

Chemigation Restrictions:

- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply and must be followed.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application, if they irrigate nontarget areas.
- Do not allow irrigation water to collect or runoff and pose a hazard to livestock, wells, or adjoining crops.
- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn.
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

Rotational Crop Restrictions

Crops labeled for GF-3052, plus turf and ornamental crops, may be immediately rotated to a treated field. All other crops may be rotated 30 days following last application.

Use Directions

Barley, Triticale and Wheat

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
Aphids (except Russian wheat aphid) Greenbug	1.25 – 2.0
cereal leaf beetle	2.5

Russian wheat aphid armyworms ¹ corn earworm (headworm) grasshoppers (suppression) southwestern corn borer webworms	2.5
---	-----

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

Application Timing: Scout for **armyworms** with enough regularity to monitor egg laying and egg hatch and treat when thresholds are reached. Time applications of GF-3052 WG to coincide with peak egg hatch and/or small larval stage of each generation. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified for the target pest. Use a higher rate in the rate range for heavy infestations, advanced growth stages of target pests, or difficult spray coverage situations.

Resistance Management: Do not make more than two consecutive applications of Group 4C (sulfoxaflor) or 5 (spinetoram and spinosad) insecticides. If additional treatments are required after two consecutive applications of Group 4C or 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 21 days of grain or straw harvest or within 7 days of grazing, or forage, fodder, or hay harvest.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not make more than 2 applications per crop. Do not make more than 2 applications of any product containing sulfoxaflor per crop, or 3 applications of any product containing spinetoram, per year.
- Do not apply more than a total of 5.0 oz of GF-3052 WG (0.09 lb ai sulfoxaflor and 0.03 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.09 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.141 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Brassica (Cole) Leafy Vegetables (Crop Group 5)¹

¹Brassica (cole) leafy vegetables (crop group 5) broccoli, broccoli raab, Brussels sprouts, cabbage, cauliflower, cavalo, Chinese broccoli, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens

In the state of Georgia, do not apply GF-3052 WG to broccoli raab, Chinese cabbage (bok choy), collards, kale, mizuna, mustard greens, mustard spinach, rape greens

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
aphids	1.5-2.0
armyworms ¹ cabbage looper ² diamondback moth ² imported cabbageworm ² light brown apple moth	4.5

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of lepidopterous larvae, leafminers, and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Heavy infestations may require repeat applications not to exceed 14 oz of GF-3052 WG per acre per year, but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pest. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Two applications may be required for optimum control of whiteflies.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. For diamondback moth and thrips, if additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 2 applications. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Make treatment decisions for the entire farm and consider area wide programs if other growers are in close proximity. Do not make more than 6 applications of GF-3052 WG per calendar year for diamondback moth over an entire farm (an area of abutting or nearby fields).

Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 6 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor or spinetoram that total more than 0.266 lb ai per acre per year.
- Do not apply to seedling cole crops grown for transplant within a greenhouse, shade house, or field plot.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Bulb Vegetables (Crop Group 3)¹

¹Bulb vegetables (crop group 3) dry bulb onion, garlic, great-headed (elephant) garlic, green onion, leek, shallot, Welsh onion

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
armyworms ^{1,2} European corn borer ² loopers ²	4.5

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of lepidopterous larvae may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing Directions. Control of thrips requires thorough coverage of the crop. Coverage can be increased by using higher spray pressure and dual directed nozzles.

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for larger larvae or heavier infestations.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. For thrips, if additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 2 applications. Do not make more than 2 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 5 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.234 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Citrus (Crop Group 10)¹

¹Citrus (crop group 10) grapefruit, lemons, limes, oranges, tangerines

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
Aphids	1.5-2.5

Asian citrus psyllid ¹ citrus leafminer ¹ citrus orangedog katydids ² lepidopterous larvae, including: avocado leafroller citrus peelminer cutworms fruit tree leafroller light brown apple moth orange tortrix western tussock moth citrus snow scale mealybugs citrus thrips, <i>Scirtothrips citri</i> ¹ citricola scale Florida red scale California red scale (suppression only)	4.5
---	-----

¹Control of leafminers, thrips, and psylla may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

²Katydid: Control of nymphs only; suppression of adults.

Advisory Pollinator Statement: Notifying known beekeepers within 1 mile of the treatment area 48 hours before the product is applied will allow them to take additional steps to protect their bees. Also, limiting application to times when managed bees and native pollinators are least active, e.g., before 7 am or after 7 pm local time or when the temperature is below 55°F at the site of application, will minimize risk to bees.

Application Timing: Treat when pests appear or in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area. Time application for scales to the crawler stage.

Application Rate: The amount of GF-3052 WG applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or smaller trees and a higher rate in the rate range for heavy infestations and/or larger trees.

To achieve good control, uniform coverage and proper equipment calibration are required. Apply only with air-assisted equipment. Low volume applications may not provide the same level of efficacy that can be obtained with an airblast sprayer at higher application volumes.

Reduce the potential for drift when making low volume applications. Factors that may impact drift include wind speed and temperature inversion. Only apply this product if the wind direction favors on-target deposition. Do not apply when wind velocity exceeds 15 mph. See the Application Directions section for additional information on reducing spray drift.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. For citrus thrips, rotate to another class of effective products for the next 2 applications after using 2 applications of GF-3052 WG within a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not make more than 3 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 3 applications of any product containing spinetoram, per year.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.188 lb ai per acre per year.
- Do not apply to citrus nurseries or citrus in greenhouses.
- Only one application is allowed between 3 days before bloom and until after petal fall per year.

Cotton**Pests and Application Rates:**

Pests	GF-3052 WG (oz/acre)
thrips (early season) ²	2.0-3.5
cotton aphid	1.5-2.0
cotton bollworm (pre-bloom) cotton leafperforator European corn borer tobacco budworm tarnished plant bug western tarnished plant bug silverleaf whitefly sweetpotato whitefly	3.75

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of lepidopterous larvae, leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Advisory Pollinator Statement: Notifying known beekeepers within 1 mile of the treatment area 48 hours before the product is applied will allow them to take additional steps to protect their bees.

Application Timing: Treat when pests appear or in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

For **cotton bollworm**, use a lower rate in the rate range at pre-bloom timings and a higher rate in the rate range at post-bloom timings. For **tobacco budworm** and/or **cotton bollworm**, scout fields twice per week and apply GF-3052 WG when the majority of the population is within blackhead egg stage to 1/8-inch larval length. The following table illustrates the size of development of worms in relation to age and stage of development (instar) as a guide to timing treatments for optimum control:

Age (Days)	Average Size (Inch)	Instar
Hatch	1/16	1st
3	¼	2nd
5	½	3rd
8	7/8	4th
10	1	5th

Note: A scouting schedule of only once per week is risky since hatching worms will have grown to 3rd instar before the next scouting observation has determined the need to spray.

Beet Armyworm: Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply GF-3052 WG when field scouting reveals three occurrences or more of egg hatch or larval feeding per 100 feet of row.

Loopers: Economic thresholds vary with local conditions and sampling methods. The following is an example of one such method: apply GF-3052 WG when field scouting reveals 4 larvae per 1 foot of row or 25% defoliation.

Application Rate: Use a higher rate of GF-3052 WG in the rate range and higher spray volume when one or more of the following is true: tobacco budworms or bollworms are more than 1/4 inch in length; target pest population is 2X above state threshold level; or foliage canopy is tall/dense and worms are present in the lower part of the canopy. Heavy infestations may require repeat applications, but follow resistance management guidelines and do not exceed 14.0 oz of GF-3052 WG per acre per year. Two applications may be required for optimum whitefly or plant bug control under high pest pressure or heavy immigration of plant bugs from other crops.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 5 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor or spinetoram that total more than 0.266 lb ai per acre per year.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 6 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.

Cucurbit Vegetables (Crop Group 9)¹

¹Cucurbit vegetables (crop group 9) cucumber, edible gourds, muskmelons (cantaloupe, honeydew, etc.), pumpkin, summer squash, watermelon, winter squash

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
aphids	1.5-2.0
armyworms ¹ cabbage looper melonworm pickleworm rindworms	3.75

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of lepidopterous larvae, leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Advisory Pollinator Statement: Notifying known beekeepers within 1 mile of the treatment area 48 hours before the product is applied will allow them to take additional steps to protect their bees. Also, limiting application to times when managed bees and native pollinators are least active, e.g., before 7 am or after 7 pm local time or when the temperature is below 55°F at the site of application, will minimize risk

to bees.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Heavy infestations may require repeat applications, but follow resistance management guidelines and do not exceed 14.0 oz of GF-3052 WG per acre per year. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Two applications may be required for optimum whitefly control.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest for all cucurbit crops except cucumbers. Do not apply within 1 day of harvest for cucumbers.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor or spinetoram that total more than 0.266 lb ai per acre per year.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year. See Resistance Management regarding number of applications for specific pests.

Fruiting Vegetables (Crop Group 8)¹ and Okra

¹Fruiting vegetables (crop group 8) eggplant, ground cherry, pepino, pepper (except black), tomatillo, tomato

Pests and Application Rates:

Pests	TwingGuard (oz/acre)
aphids	1.5-2.0
armyworms ¹	3.75
Colorado potato beetle	
European corn borer	
hornworms	
light brown apple moth	
loopers	
tomato fruitworm, <i>Helicoverpa zea</i>	
tomato pinworm	

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Scout weekly throughout the season to monitor populations of **leafminers** and **thrips** to determine when economic thresholds are exceeded. Scout

weekly throughout the season to monitor beneficial populations. For **lepidopterous larvae**, scout with enough regularity to monitor the population size of each of the labeled pests. Time applications of GF-3052 WG to coincide with peak egg hatch in species without overlapping generations.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Two applications may be required for optimum whitefly control.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. For **thrips**, if additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 2 applications. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Do not apply Group 5 insecticides to consecutive generations of Colorado potato beetle and do not make more than 2 applications per single generation of Colorado potato beetle.

Restrictions:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- **Minimum Re-Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per calendar year. Do not make applications of any product containing sulfoxaflor or spinetoram that total more than 0.266 lb ai per acre per year.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 6 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.
- Do not apply to seedling fruiting vegetables and okra grown for transplant within a greenhouse, shade house, or outdoor transplant bed.

Grape

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
cutworm	4.5
European grapevine moth	
grape berry moth	
grape leaffolder	
light brown apple moth	
omnivorous leafroller	
orange tortrix	
redbanded leafroller	
thrips ¹	
western grape leaf skeletonizer	
grape leafhopper	
mealybugs	

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Heavy infestations may require repeat applications not to exceed 14.0 oz of GF-3052 WG per acre per year, but follow resistance management

guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Carefully adjust equipment and spray volume to assure thorough uniform coverage. Use a higher rate of GF-3052 WG in the rate range for larger larvae or moderate to severe infestations and/or larger plant volume.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.305 lb ai per acre per year.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 5 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Leafy Vegetables (Except *Brassica*) (Crop Group 4)¹, Leaves of Root and Tuber Vegetables (Crop Group 2)² and Watercress

¹Leafy vegetables (except *Brassica*) (crop group 4) amaranth, arugula, cardoon, celery, celtuce, chervil, Chinese celery, Chinese spinach, corn salad, dandelion, dock, edible-leaved chrysanthemum, endive (escarole), Florence fennel, garden cress, garden purslane, garland chrysanthemum, head lettuce, leaf lettuce, leafy amaranth, New Zealand spinach, orach, parsley, radicchio (red chicory), rhubarb, spinach, Swiss chard, tampala, upland cress, vine spinach, winter cress, winter purslane, yellow rocket

²Leaves of root and tuber vegetables (crop group 2) bitter cassava, black salsify, carrot, celeriac (celery root), chicory, dasheen (taro), edible burdock, garden beet, oriental radish (daikon), parsnip, radish, rutabaga, sugar beet, sweet cassava, sweet potato, tanier, true yam, turnip, turnip-rooted chervil

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
aphids	1.5-2.0
armyworms ¹ cabbage looper corn earworm, <i>Helicoverpa zea</i> diamondback moth imported cabbageworm light brown apple moth	4.5

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Scout at least weekly and consider the impact of both pests and beneficials. Treat

when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Heavy infestations may require repeat applications not to exceed 14.0 oz of GF 3052 WG per acre per year, but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Two applications may be required for optimum whitefly control.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:**
Leafy vegetables (including watercress): Do not apply within 7 days of harvest.
- **Leaves of root and tuber vegetables:** Do not apply within 7 days of harvest. **Note:** Root and tuber vegetables treated as described may only be harvested for the foliage, not for the root or tuber.
- **Minimum Treatment Interval:**
Leafy vegetables (including watercress): Do not make applications less than 7 days apart.
Leaves of root or tuber vegetables: Do not make applications less than 7 days apart.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 6 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.
- Do not apply to seedling leafy crops grown for transplant within a greenhouse, shade house, or outdoor transplant bed.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor or spinetoram that total more than 0.266 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Root and Tuber Vegetables (Crop Group 1)¹

¹Root and tuber vegetables (crop group 1) arracacha, arrowroot, bitter cassava, black salsify, carrot, celeriac, chayote root, chicory, Chinese artichoke, chufa, dasheen, edible burdock, edible canna, garden beet, ginger, ginseng, horseradish, Jerusalem artichoke, leren, oriental radish, parsnip, potato, radish, rutabaga, salsify, skirret, Spanish salsify, sugar beet, sweet cassava, sweet potato, tanier, true yam, tumeric, turnip, turnip-rooted chervil, turnip-rooted parsley, yam bean

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
Aphids	1.5-2.0
Colorado potato beetle	3.5

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of leafminers, psyllids, and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Consult your Dow

AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Heavy infestations may require repeat applications not to exceed 14.0 oz of GF-3052 WG per acre per year, but follow resistance management guidelines. Two applications may be required for optimum whitefly control.

Chemigation: GF-3052 WG may be applied to potatoes by chemigation at labeled rates. Refer to the Chemigation Application section.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not apply Group 5 insecticides to consecutive generations of Colorado potato beetle and do not make more than 2 applications of Group 5 insecticides per single generation of Colorado potato beetle. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

Garden beet and sugar beet:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.250 lb ai per acre per year.
- See Resistance Management regarding number of applications for specific pests.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Black salsify, carrot, chicory, ginseng, horseradish, parsnip, salsify, skirret, Spanish salsify, turnip-rooted chervil, turnip-rooted parsley:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.219 lb ai per acre per year.
- See Resistance Management regarding number of applications for specific pests.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Arracacha, arrowroot, bitter cassava, chayote root, Chinese artichoke, chufa, dasheen, edible canna, ginger, Jerusalem artichoke, leren, potato, sweet cassava, sweet potato, taniel, true yam, turmeric, yam bean:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year.

- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.250 lb ai per acre per year.
- See Resistance Management regarding number of applications for specific pests.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Celeriac, edible burdock, oriental radish, radish, rutabaga, turnip and other root vegetables not specifically listed:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not make more than 3 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 3 applications of any product containing spinetoram, per year.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.188 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Soybean

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
soybean aphid	1.5-2.0
armyworms ¹ cabbage looper corn earworm (podworm) green cloverworm saltmarsh caterpillar soybean looper true armyworm velvetbean caterpillar	3.0-3.75

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae.

Application Rate: Use a higher rate in the rate range for heavy infestations and/or difficult spray coverage situations.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 4C or 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.109 lb ai per acre per year.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product

- containing sulfoxaflor per crop or spinetoram per year.
- Do not make more than two applications to soybean forage.

Stone Fruits (Crop Group 12)¹

¹Stone fruits (crop group 12) apricots, cherries, nectarines, peaches, plums, plumcots, prunes

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
aphids	1.5-2.0
peach twig borer (dormant spray)	4.5

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear and in accordance with local economic thresholds.. Closely follow regional spray recommendations for oriental fruit moth applications based upon biofix dates, egg hatch, and/or pheromone trap catches. Oriental fruit moth larvae must be controlled before they penetrate the fruit. GF-3052 WG is a larvicide; begin applications shortly before egg hatch. For oriental fruit moth and thrips, repeat applications may be needed to maintain control; but follow resistance management guidelines. Peach twig borer applications can be made as dormant, delayed dormant, or May sprays. For cherry fruit fly and western cherry fruit fly, maintain protective sprays at 7-day intervals while adults are present and fruit is susceptible to attack. Time application for San Jose scale to the crawler stage. For all pests, consult with your Dow AgroSciences representative, state agricultural experiment station, certified pest control advisor or extension specialist for specific application timings in your area.

Application Rate: The amount of GF-3052 WG applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations or smaller trees and a higher rate in the rate range for heavy infestations or larger trees.

Resistance Management: Do not make more than 2 consecutive applications of a Group 4C insecticide (sulfoxaflor) or 3 consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after 2 consecutive applications of Group 4C insecticide or 3 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not treat consecutive generations of oriental fruit moth and leafrollers with Group 5 insecticides. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- Preharvest Interval:** Do not apply within 14 days of harvest for apricots or 7 days of harvest for other stone fruit crops.
- Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year.
- See Resistance Management regarding number of applications for specific pests.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.438 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Strawberry

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
armyworms ¹ leafrollers light brown apple moth thrips ² plant bugs	3.75

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Advisory Pollinator Statement: Notifying known beekeepers within 1 mile of the treatment area 48 hours before the product is applied will allow them to take additional steps to protect their bees. Also, limiting application to times when managed bees and native pollinators are least active, e.g., before 7 a.m. or after 7 p.m. local time or when the temperature is below 55°F at the site of application, will minimize risk to bees.

Application Timing: Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate specified to control target pests. Use a higher rate in the range for larger larvae or moderate to severe pest infestations.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 4C or Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. For thrips, if additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 2 applications. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.234 lb ai per acre per year.
- Do not make more than 3 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 3 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.

Succulent, Edible Podded and Dry Beans¹

¹Succulent, edible podded, and dry beans including adzuki bean, asparagus bean, bean, blackeyed pea, broad bean, chickpea, Chinese longbean, cowpea, fava bean, field bean, garbanzo bean, grain lupine, green lima bean, jackbean, kidney bean, lablab bean, lima bean, moth bean, mung bean, navy bean, pinto bean, rice bean, runner bean, snap bean, soybean (immature seed), sweet lupine, sword bean, tepary bean, wax bean, white lupine, white sweet lupine, yardlong bean.

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
Aphids	1.5-2.0
European corn borer (eggs & larvae) armyworms ¹ corn earworm, <i>Helicoverpa zea</i> light brown apple moth loopers	3.75

¹With the exception of yellowstriped armyworm and western yellowstriped armyworm.

²Control of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Heavy infestations may require repeat applications, but follow resistance management guidelines and do not exceed 14.0 oz of GF-3052 WG per acre per year. Treat when pests appear and in accordance with local economic thresholds. When treating for worms, target eggs at hatch or small larvae. For **European corn borer**, treat when moth flights first appear and use the lower end of the rate range to control eggs and larvae every three days before they enter the plant. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations for your area.

Application Rate: Apply GF-3052 WG as a foliar spray at the rate specified to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

Resistance Management: Do not make more than 2 consecutive applications of Group 4C (sulfoxaflor) or Group 5 (spinetoram and spinosad) insecticides. If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Do not make more than 3 applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area

Restrictions:

- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- Do not make more than 4 applications per crop. Do not make more than 4 applications of any product containing sulfoxaflor per crop, or 6 applications of any product containing spinetoram, per year. See Resistance Management regarding number of applications for specific pests.

Succulent Beans:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.219 lb ai per acre per year.

Dried Beans:

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- Do not apply more than a total of 14 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.094 lb ai per acre per year.

Tree Nuts (Crop Group 14)¹ and Pistachios

¹Tree nuts (crop group 14) almonds, cashew, chestnut, filbert (hazelnut), macadamia nut, pecan walnut

Pests and Application Rates:

Pests	GF-3052 WG (oz/acre)
aphids	1.5-2.0
peach twig borer (dormant spray)	4.0

Application Timing: Apply GF-3052 WG as either a dormant or as a foliar spray in accordance with local economic thresholds. Closely follow regional spray recommendations based upon biofix dates, egg hatch, and/or pheromone trap catches. Lepidopterous larvae must be controlled before they penetrate the nuts or shoots. GF-3052 WG is a larvicide; begin applications shortly before egg hatch. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Use of Crop Oils: Crop oils labeled for agricultural use may be added to the dormant spray solution for suppression of overwintering mites and scale insects. Consult specific oil labels and University or Extension recommendations for precautions and restrictions regarding the use of oils in nut and fruit trees.

Application Rate: The amount of GF-3052 WG applied per acre will depend upon tree size, volume of foliage present and pest pressure. Use a higher rate in the rate range for larger trees or heavy infestations.

Resistance Management: Do not make more than 2 consecutive applications of a Group 4C insecticide (sulfoxaflor) or 3 consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after 2 consecutive applications of a Group 4C or 3 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 1 application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not apply treatments less than 7 days apart.
- Do not make more than 4 applications per year. Do not make more than 4 applications of any product containing sulfoxaflor per crop or spinetoram per year.
- Do not apply more than a total of 14.0 oz of GF-3052 WG (0.266 lb ai sulfoxaflor and 0.088 lb ai spinetoram) per acre per year. Do not make applications of any product containing sulfoxaflor that total more than 0.266 lb ai per acre per year. Do not make applications of any product containing spinetoram that total more than 0.438 lb ai per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

EPA accepted __/__/__