



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 25, 2020

Maria Pavlakos
Global Regulatory Manager - Crop Protection
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Amendment to a Supplemental Label – Extended expiration date, and fixed typos
Product Name: MEZAVUE
EPA Registration Number: 62719-717
Application Date: November 5, 2019
Decision Number: 557565

Dear Ms. Pavlakos:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval is only for a supplemental label that is an addendum to the master label. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

This supplemental labeling contains some new and/or revised uses and/or directions which may be additional to the uses and/or directions found on the label on or attached to the container, but this supplemental labeling does not by itself constitute the complete set of use directions. The complete set of use directions is set forth on the container label as combined with this supplemental labeling.

A stamped copy of your labeling is enclosed for your records. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Curtis Hildebrandt at 703-347-8198 or by email at hildebrandt.curtis@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

RESTRICTED USE PESTICIDE

May Injure (Phytotoxic) Susceptible, Non-Target Plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

Supplemental Labeling

Dow AgroSciences, 9330 Zionsville Road, Indianapolis, IN 46268-1054 USA

MezaVue™

EPA Reg. No. 62719-717

For Distribution and Use Only in the Following States:
AL, AR, AZ, CO, GA, ID, KS, LA, MO, MS, MT, ND, NE, NV, NM, OK, SD, TN, TX, UT, WY

**For Use on Grass Harvested for Hay
Intended for Distribution or Sale Off the Farm or Ranch**

**For Use On Grass Harvested for Silage, Haylage, Baylage, or Green Chop
Intended for Use On the Farm or Ranch**


This supplemental label expires on May 30, 2023, and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for MezaVue herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of MezaVue according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for MezaVue.

Use Precautions and Restrictions


Consult with a Dow AgroSciences representative if you do not understand the Use Precautions and Restrictions. Call 800-258-3033 for more information.



IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- It is mandatory to follow the **Use Precautions and Restrictions** section of this product label.
- Manure and urine from animals consuming treated grass or forage may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
- The Applicator must provide the land manager with a copy of the Dow AgroSciences Stewardship instructions regarding uses of forage from areas treated with MezaVue.

Forage and Manure Management



- **Do not use grasses treated with MezaVue in the preceding 18 months for hay intended for export outside the United States.**
- **Do not use hay or straw from areas treated with MezaVue within the preceding 18 months, or manure from animals feeding on hay treated with MezaVue, in compost.**
- **Do not use grasses treated within the preceding 18 months for seed production.**
- **Do not use on Timothy hay or other cool-season grasses grown for hay.**
- Do not overseed ryegrass for 4 months after treatment.
- **MezaVue is highly active against many broadleaf plant species.** Do not use this product on areas where loss of desirable broadleaf forage plants, including legumes, cannot be tolerated.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if MezaVue concentration remaining in the soil will adversely affect the legume establishment.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or grass hay harvest following application of MezaVue at labeled rates for non-lactating dairy animals or other livestock. Cutting hay too soon after spraying weeds will reduce weed control. For optimum uptake and translocation of MezaVue, wait 14 days after herbicide application to cut grass hay to allow herbicide to work. Do not transfer grazing animals from areas treated with MezaVue to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough MezaVue to cause injury to sensitive broadleaf plants.

- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.
- **Transfer of Animals Feeding on MezaVue-Treated Forage:** Do not transfer animals grazing or feeding on hay from areas treated with MezaVue to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough MezaVue to cause injury to sensitive broadleaf plants.
- **Restrictions in Hay or Manure Use:**
 - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18 months in compost, mulch, or mushroom spawn.
 - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch, or mushroom spawn.
 - Do not spread manure from animals that have grazed or consumed forage or hay from treated areas within the previous 3 days on land used for growing broadleaf crops.
 - Manure from animals that have grazed forage or eaten hay harvested from MezaVue-treated areas within the previous 3 days may only be used on pasture grasses, grass grown for seed, and wheat.
 - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from MezaVue-treated areas until an adequately sensitive field bioassay is conducted to determine that the MezaVue residues in the soil is at level that is not injurious to the crop to be planted.
 - To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of MezaVue in plant residues or manure is more rapid under warm, moist soil conditions and may be accelerated by supplemental irrigation.
- **Crop Rotation:** Do not rotate non-cropland to cropland for one year following an application of MezaVue. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of MezaVue present in the soil will not adversely affect that broadleaf crop.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern or drainage. The field bioassay can be initiated at any time between harvest of the treated crop and the planting of the intended rotational crop. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, forage grasses, native grasses or grasses grown for hay.
- Trees adjacent to or in a treated area can occasionally be affected by root uptake of MezaVue through movement into the soil. Do not apply MezaVue within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses, and leguminous trees such as locusts, redbud, mimosa, and caragana.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R429-002

EPA accepted ___/___/___

Replaces R429-001