



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

62719-754

Date of Issuance:

5/29/25

NOTICE OF PESTICIDE:

☒ Registration

☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

GF-3840 AG

Name and Address of Registrant (include ZIP Code):

Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

Signature of Approving Official:

Nathan Mellor, Chief  
Fungicide and Herbicide Branch,  
Registration Division (7505T)

Date:

5/29/25

2. Registration Term for Implementing Additional Mitigations from the Services

If, following formal consultation with FWS and NMFS, additional modifications are identified in any applicable Biological Opinion, EPA will notify Corteva in writing within 45 calendar days of the issuance of the Biological Opinion of any necessary required changes. Within 30 calendar days of receiving EPA's notice, Corteva must submit an amendment application incorporating the necessary changes, including amended labels. Alternatively, Corteva may respond by submitting a request for voluntary cancellation of this product. If Corteva fails to comply with this term, Corteva has agreed in prior written acceptance of the terms that EPA may cancel the registration under an expedited process under FIFRA 6(e).

3. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, "EPA Reg. No. 62719-754."

4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 5/12/2025

If you have any questions, please contact Lydia Crawford at 202-566-2575 or at [crawford.lydia@epa.gov](mailto:crawford.lydia@epa.gov).

Enclosure

(Base label):

FLORYLPICOXAMID	GROUP	21	FUNGICIDE
-----------------	-------	----	-----------

# GF-3840 AG

## FUNGICIDE

Adavelt™ Active

**For disease control in the following crops: barley, canola, dry peas and beans, sugar beet, and wheat.**

**For use in the contiguous United States and Hawaii only.**

### Active Ingredient:

Florylpicoxamid\*:

(2S)-1,1-bis(4-fluorophenyl)propan-2-yl N-[[3-(acetyloxy)-4-methoxypyridin-2-yl]carbonyl]


-L-alaninate ..... 9.9%

**Other Ingredients** ..... 90.1%

**Total** ..... 100.0%

\*CAS # 1961312-55-9

GF-3840 AG is formulated as an emulsifiable concentrate (EC) containing 0.83 lb ai/gal of florylpicoxamid.



### Keep Out of Reach of Children

## WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes</b>	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
<b>If swallowed</b>	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
<b>If on skin or clothing</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 -20 minutes. Call a poison control center or doctor for treatment advice.
<b>NOTE:</b> Have the product container or label with you when calling the poison control center or doctor, or going for treatment. For medical emergencies, call the poison control center at 800-222-1222. For general information on this product, call 800-441-3637.	

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

**Causes Substantial But Temporary Eye Injury** Do not get in eyes, on skin, or on clothing. Wear appropriate protective eye wear (goggles, face shield or safety glasses). • **Harmful If Swallowed or if Absorbed Through Skin.** Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, shoes plus socks and appropriate chemical-resistant/waterproof gloves.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Protective eyewear (goggles, faceshield, or safety glasses).
- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant/waterproof (such as nitrile or butyl) gloves

### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

---

## DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See inside of booklet for precautionary statements and complete Directions for Use.

Mix or shake well before use.

### (Storage and Disposal for rigid containers 5 gal or less)

#### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an

approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### (Storage and Disposal for refillable rigid containers greater than 5 gal)

##### **Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### (Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

##### **Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local

authorities.

**Notice:** Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-TLU

Batch / Lot (see container) *(Editor's note for artwork: required for non-refillable pkg)*

EPA Est. \_\_\_\_\_

[CD02-xxx-xxx]

<sup>TM</sup>®Trademarks of Corteva Agriscience and its affiliated companies

**[Produced for] [Distributed by] [Manufactured for]**  
**Corteva Agriscience LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET CONTENTS \_\_\_\_\_**

(Booklet cover):. {Note – text from the booklet cover may be used as the base label for containers too small to hold the full text from the base label page above, e.g., 1-gallon container.}

FLORYLPICOXAMID	GROUP	21	FUNGICIDE
-----------------	-------	----	-----------

## GF-3840 AG

### FUNGICIDE

Adavelt™ Active

**For disease control in the following crops: barley, canola, dry peas and beans, sugar beet, and wheat.**

**For use in the contiguous United States and Hawaii only.**

#### Active Ingredient:

Florylpicoxamid\*:

(2S)-1,1-bis(4-fluorophenyl)propan-2-yl N-[[3-(acetyloxy)-4-methoxypyridin-2-yl]carbonyl]

-L-alaninate ..... 9.9%

**Other Ingredients** ..... 90.1%

**Total** ..... 100.0%

\*CAS # 1961312-55-9

GF-3840 AG is formulated as an emulsifiable concentrate (EC) containing 0.83 lb ai/gal of florylpicoxamid.

#### Keep Out of Reach of Children

### WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

See inside of booklet for First Aid, precautionary statements, and complete Directions for Use.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.  
Mix or shake well before use

EPA Reg. No. 62719-TLU

Batch / Lot (see container) (Editor's note for artwork: required for non-refillable pkg)

EPA Est. \_\_\_\_\_

[CD02-xxx-xxx]

™®Trademarks of Corteva Agriscience and its affiliated companies

[Produced for] [Distributed by] [Manufactured for]

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268

**NET CONTENTS** \_\_\_\_\_

[Booklet Page 1 through end]]

## WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes</b>	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
<b>If swallowed</b>	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
<b>If on skin or clothing</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 -20 minutes. Call a poison control center or doctor for treatment advice.
<b>NOTE:</b> Have the product container or label with you when calling the poison control center or doctor, or going for treatment. For medical emergencies, call the poison control center at 800-222-1222. For general information on this product, call 800-441-3637.	

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**Causes Substantial But Temporary Eye Injury** Do not get in eyes, on skin, or on clothing. Wear appropriate protective eye wear (goggles, face shield or safety glasses). • **Harmful If Swallowed or if Absorbed Through Skin.** Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, shoes plus socks and appropriate chemical-resistant/waterproof gloves.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Protective eyewear (goggles, faceshield, or safety glasses)
- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant/waterproof (such as nitrile or butyl) gloves

### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.



**User Safety Recommendations**

Users should:

- Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours

**DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS:** Before using this product, you must obtain any applicable Endangered Species Protection Bulletins (Bulletins) within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins>. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of Federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov)."

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Protective eyewear (goggles, faceshield, or safety glasses)

- Coveralls
- Chemical resistant/waterproof (such as nitrile or butyl) gloves
- Shoes plus socks

## PRODUCT INFORMATION

GF-3840 AG (Adavelt) is a broad-spectrum fungicide with preventive and curative properties registered for the control of key diseases in the specified crops. The active ingredient in GF-3840 AG is Florylpicoxamid, which belongs to the Quinone Inside Inhibitors (Qil) class, target site of action Group 21. This fungicide may be applied as a foliar treatment, solo or in tank mixes with other crop protection products. Optimum disease control is achieved when this product is applied in alternating, regularly scheduled preventative spray programs, following the labeled rates and use directions, in a rotation program with other registered fungicide groups for the specified crops and targets.

## USE PRECAUTIONS

Carefully read, understand and follow label use rates and restrictions. For proper application, determine the size of the area to be treated, the specified label use rate and the gallonage to be applied to the area. Use maximum label rates and shortened spray schedules for severe or threatening disease conditions. Under low disease conditions, minimum label use rates per application can be used. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is advised prior to use. Under certain weather conditions (particularly high temperatures), this fungicide in combination with high rates of non-ionic surfactant or oil containing (petroleum or crop) additives or adjuvants may cause injury.

## USE RESTRICTIONS

- **DO NOT** exceed the maximum use rate of product per acre per year, the maximum rate per application, and the total number of applications of GF-3840 AG per year as stated in Specific Uses and Directions, below. Follow the preharvest interval (PHI) restrictions as stated.
- When used alone, or in combination with other products that contain florylpicoxamid, **DO NOT** apply more than 0.268 pounds ai/A/yr.
- **DO NOT** tank mix with copper based fungicides.
- **For use in the contiguous United States and Hawaii only.**

## RESISTANCE MANAGEMENT

### Integrated Pest Management

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when disease forecasting models reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

### Resistance

For resistance management, GF-3840 AG contains florylpicoxamid, a Group 21 fungicide. Any fungal population may contain individuals naturally resistant to GF-3840 AG and other Group 21 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields without rotation with other fungicides from a different Group. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of GF-3840 AG or other Group 21 fungicides within a growing season sequence with different groups that control the same pathogens. Avoid consecutive applications of GF-3840 AG or other fungicides in the same group in a season.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest

- when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
  - Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
  - Monitor treated fungal populations for resistance development.
  - Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
  - For further information or to report suspected resistance contact your local company representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

### **MANDATORY SPRAY DRIFT MANAGEMENT MITIGATIONS**

#### **For Aerial and Ground Boom Applications:**

- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Select nozzle and pressure that deliver medium or coarser spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with American Society of Agricultural & Biological Engineers standards 572.1 and 641 (ASABE S572 and S641).
- During application, the Sustained Wind Speed, as defined by the National Weather Service (standard averaging period of 2 minutes) must register between 3 and 15 miles per hour.
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- Do not apply during temperature inversions.

#### **For Aerial Application:**

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft to minimize drift caused by wing tip or rotor blade vortices.
- Wind speed and direction must be measured on location using a windsock, an anemometer (including systems to measure wind speed or velocity on an aircraft), or an aircraft smoke system.
- When the wind speed is between 11-15 miles per hour, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- When the wind speed is between 11-15 miles per hour, applicators must use a minimum of  $\frac{3}{4}$  swath displacement upwind at the downwind edge of the field. Otherwise, applicators must use a minimum of  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field.
- Do not release spray at a height greater than 10 ft above the crop canopy unless a greater application height is required for pilot safety.

#### **For Ground Boom Application:**

- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but do not exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions.
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure wind speed or velocity using application equipment).

#### **ADDITIONAL SPRAY DRIFT INFORMATION:**

This section is intended to provide additional information for applicators to assist in implementing the mandatory spray drift mitigations above. THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. Be aware of nearby non-target sites and environmental conditions.

#### **Importance of Droplet Size**

An effective way to reduce spray drift is to apply large droplets. Consider the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size — Aircraft**

- **Adjust Nozzles** - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **Release Height – Aircraft**

- Higher release heights increase the potential for spray drift.

### **Release Height - Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.

### **Controlling Droplet Size — Ground Boom**

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Shielded Sprayers**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **Temperature and Humidity**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **Wind**

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

**Measuring wind speed and wind direction:** Applicators should check and acquire the predicted wind speed and direction for the application site within 12 hours prior to conducting applications to determine the time periods wind speed is likely to fall outside the applicable thresholds.

Applicators should reassess wind speed and direction at the application site every 15 minutes while applications are in progress. Measuring wind speed and direction can be done by:

- Relying on equipment on the application equipment that measures wind speed (e.g., aerial equipment).
- Using a tower anemometer with telemetry or handheld anemometer. Users should read user manual on how to calibrate, operate and interpret the output from an anemometer. Ground applicators should stop every 15 minutes to take a reading with a tower anemometer with telemetry or handheld anemometer. Some anemometers may have software that would allow users to view wind measurements in real time while making an application, and, in those cases, applicators would not have to stop to take measurements.
- Using a windsock. Wind can be estimated with a windsock using the strips on a windsock. The applicator should consult the user manual for the windsock on wind speed estimation and direction of wind. Applicators should look at the sock at least every 15 minutes to estimate wind speed and direction. The windsock should be pointed in the opposite direction of the windbreak and the non-managed area.

- Using an aircraft smoke system. Laying down several puffs of smoke along different lines using an aircraft smoke system can provide an accurate view of what the wind speed and direction for the application.
- Checking behind the spray rig at least every 15 minutes to see if the spray has changed direction from when the application started.

## COMPATIBILITY

GF-3840 AG is compatible with most commonly used fungicides (except copper based fungicides), insecticides, growth regulators, micronutrients and spray additives and adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialist prior to use. When an additive and adjuvant is to be used with this product, it is advised to use a Chemical Producers and Distributors Association certified additive and adjuvant. To determine the physical compatibility of GF-3840 AG with other products, use a jar test. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions and in the correct mixing order as defined below. Invert the jar containing the mixture several times after adding each product and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible, and the tank mix combination should not be used. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

## SPRAY ADDITIVES

Under some conditions, the use of additives or adjuvants may improve the performance of GF-3840 AG. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which Corteva has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing GF-3840 AG with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

A spreading/penetrating type adjuvant such as a non-ionic surfactant, crop oil concentrate, or blend may be added at the manufacturer's recommended rates. **DO NOT** include an additive including latex, as efficacy may be reduced. Consult a Corteva representative for more information concerning additives or adjuvants.

## MIXING DIRECTIONS

**DO NOT** prepare more spray mixture than is required for the immediate operation. **DO NOT** tank mix with copper-based fungicides. Thoroughly clean spray equipment before and after using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use. Rinsate may be disposed of on site, according to label use directions, or at an approved waste disposal facility.

### GF-3840 AG – Alone

Add 1/2-2/3 of the required amount of water to the spray or mixing tank.

- With the agitator running, add GF-3840 AG to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after the product has completely dispersed into the mix water.
- Maintain agitation until all the mixture has been sprayed.

### GF-3840 AG – Tank Mix

#### Mixing Order for Tank Mixes:

1. Fill the spray tank to at least 1/4 to 1/3 of the required spray volume.
2. Start agitation.
3. Add different formulation types in the following order, allowing time for complete mixing and dispersion after addition of each while following label directions for each material added to the tank.

- (a) Water soluble bags
  - (b) Dry flowables
  - (c) Wettable powders
  - (d) Suspensions and liquid flowables
  - (e) GF-3840 AG and other emulsifiable concentrates and any solutions
  - (f) Adjuvants
4. Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be re-suspended before spraying is resumed.

Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and restrictions which appear on the tank-mix product label. Label dosage rate must not be exceeded, and the most restrictive label precautions and restrictions must be followed. This product must not be mixed with any product which prohibits such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### **Clean-Out Procedures for Spray Equipment:**

1. Drain any remaining spray mixture from the application equipment, then wash out tank, boom, and hoses with clean water. Drain again.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add commercial tank cleaner at manufacturers recommended rates, or at a rate of 1 gallon per 100 gallons of water. Recirculate for 10 - 20 minutes and spray out the mixture through the boom.
4. Remove all spray nozzles and screens and clean separately.
5. Thoroughly clean exterior surfaces of spray equipment.

#### **CROP ROTATION RESTRICTIONS**

- Crops found on this, or other florylpicoxamid labels can be planted immediately following the last application of GF-3840 AG.
- All other crops intended for food or feed may be planted 30 days following the last application of GF-3840 AG when applying no more than a total of 2 applications at 11 fl oz (0.07 lb ai/A) or 60 days following the last application of GF-3840 AG when applying no more than a total of 2 applications at 20.5 fl oz (0.134 lb ai/A).

#### **APPLICATION INSTRUCTIONS**

Apply this product in conjunction with management practices for the crop uses that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, accredited laboratories, or other identification methods when developing disease control strategies.

Optimum disease control is achieved when GF-3840 AG is applied as a preventative disease control spray program at a rate according to the specified use and following the use restrictions. Apply GF-3840 AG with enough water volume to ensure thorough coverage. For ground application, use at least the minimum water volume indicated. For best results, allow GF-3840 AG to dry at least 1 hour before any rainfall or field irrigation. Ensure thorough coverage of the crop for good disease control avoiding spray overlap.

Under conditions favorable for high disease development, reduce the spray interval between applications of GF-3840 AG. Under light to moderate disease pressure, apply GF-3840 AG at the low use rate and/or longer application interval. When disease pressure is high or when used as a curative treatment, use

higher rates of GF-3840 AG and/or shorter application interval unless otherwise specified. See the Specific Uses and Directions section below for specific application rates for various diseases.

**Ground Application:**

Apply GF-3840 AG in sufficient water to ensure thorough coverage of the canopy for optimum disease control. Refer to specific use directions for additional information.

**Aerial Application:**

Unless otherwise specified on this label or in supplemental labeling, use no less than 5 gallons of spray solution per acre. **DO NOT** apply when conditions favor drift from target area. Refer to spray drift section for additional information and instructions

**Chemigation**

Apply GF-3840 AG only through sprinkler irrigation systems (such as center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems).

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Specific Instructions for Public Water Systems:**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Specific Instructions for Sprinkler Irrigation Systems:**

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to



- prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
  4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
  5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
  6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
  7. Do not apply when wind speed favors drift beyond the area intended for treatment.
  8. Good agitation is required in the injection tank.
  9. In moving systems, apply specified dosage of GF-3840 AG as a continuous injection. In nonmoving systems inject GF-3840 AG for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
  10. Mix the amount of GF-3840 AG needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.
  11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all GF-3840 AG is flushed from system.



## SPECIFIC USES AND DIRECTIONS

### Barley

Disease Common Name	Disease Scientific Name	Product Use Rate per application (fl oz/A)
• Net blotch	• <i>Pyrenophora teres</i>	11 (0.07 lb ai/A)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 11 fluid ounces per acre (0.07 lb ai/acre) of GF-3840 AG per application.</li> <li>• <b>DO NOT</b> apply more than 22 fluid ounces per acre ( 0.14 lb ai/acre) of GF-3840 AG per year.</li> <li>• <b>DO NOT</b> apply more than 2 applications of GF-3840 AG per acre per year.</li> <li>• <b>DO NOT</b> apply more than 1 application of GF-3840 AG by air.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days</li> <li>• Apply no later than end of head emergence (Zadoks 59, Feekes 10.5).</li> <li>• For use in the contiguous US and Hawaii only.</li> </ul> <b>Use Instructions</b> <ul style="list-style-type: none"> <li>• Begin application of GF-3840 AG before disease development.</li> <li>• Apply the product after flag leaf emergence and before 50% head emergence.</li> <li>• Avoid consecutive applications of florylpicoxamid. GF-3840 AG should always be applied in mixture with another fungicide from a different FRAC group which provides robust control of net blotch.</li> <li>• Apply at a volume of 9 to 32 gal/A of water, ensuring thorough coverage of the canopy.</li> </ul>		

### Canola

Disease Common Name	Disease Scientific Name	Product Use Rate per application (fl oz/A)
<ul style="list-style-type: none"> <li>• Blackleg</li> <li>• White mold/Stem rot (suppression)</li> <li>• Alternaria Black spot (suppression)</li> </ul>	<ul style="list-style-type: none"> <li>• <i>Leptosphaeria maculans</i></li> <li>• <i>Sclerotinia sclerotiorum</i> (suppression)</li> <li>• <i>Alternaria</i> spp. (suppression)</li> </ul>	13.7 to 20.5 (0.09 to 0.134 lb ai/A)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 20.5 fluid ounces per acre (0.134 lb ai/acre) of GF-3840 AG per application.</li> <li>• <b>DO NOT</b> apply more than 41 fluid ounces per acre (0.268 lb ai/acre) of GF-3840 AG per year.</li> <li>• <b>DO NOT</b> apply more than 2 applications of GF-3840 AG per acre per year.</li> <li>• <b>DO NOT</b> apply more than 1 application of GF-3840 AG by air.</li> <li>• <b>Minimum Retreatment Interval:</b> 7 days</li> <li>• <b>Preharvest Interval (PHI):</b> 21 days</li> <li>• For use in the contiguous US and Hawaii only.</li> </ul> <b>Use Instructions:</b> <ul style="list-style-type: none"> <li>• For blackleg apply GF-3840 AG at 2-4 leaf stage.</li> <li>• For white mold, begin application at early flowering (20-40%) or prior to the onset of disease. Use the higher rate if conditions are conducive to disease development.</li> <li>• Avoid consecutive applications of florylpicoxamid.</li> <li>• For Alternaria Blackspot apply at early pod development.</li> <li>• For white mold and Alternaria black spot use the highest labeled rate.</li> <li>• Apply at a volume of 15 to 32 gal/A of water, ensuring thorough coverage of the canopy.</li> </ul>		

### Pulses, Dried Shelled Bean, Except Soybean, and Pulses, Dried Shelled Pea (Subgroup 6-22E<sup>1</sup> and 6-22F<sup>2</sup>)

<sup>1</sup>African yam bean; American potato bean; Bean (*Lupinus* spp., dry seed, including, but not limited to Andean lupin, blue lupin, grain lupin, sweet lupin, white lupin, white sweet lupin, and yellow lupin); Bean

(*Phaseolus* spp., dry seed, including, but not limited to black bean, cranberry bean, dry bean, field bean, French bean, garden bean, great northern bean, green bean, kidney bean, lima bean, navy bean, pink bean, pinto bean, red bean, scarlet runner bean, tepary bean, and yellow bean); Bean (*Vigna* spp., dry seed, including, but not limited to adzuki bean, asparagus bean, blackeyed pea, catjang bean, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean); broad bean (fava bean) dry seed; guar bean, dry seed; goa bean, dry seed; horse gram; jackbean, dry seed; lablab bean, dry seed; morama bean; sword bean, dry seed; winged pea, dry seed; velvetbean, dry seed; cultivars, varieties, and/or hybrids of these commodities.

<sup>2</sup>Dry seed of any of the following: Pea (*Pisum* spp.; including, but not limited to dry pea, field pea, green pea, yellow pea, wrinkled pea, marrowfat pea, and garden pea); chickpea; grass pea; lentil; pigeon pea; cultivars, varieties, and/or hybrids of these commodities.

Disease Common Name	Disease Scientific Name	Product Use Rate per application (fl oz/A)
<ul style="list-style-type: none"> <li>• Anthracnose</li> <li>• Ascochyta blight</li> <li>• Mycosphaerella blight</li> </ul>	<ul style="list-style-type: none"> <li>• <i>Colletotrichum</i> spp.</li> <li>• <i>Phoma exigua</i>, <i>Ascochyta</i> spp.</li> <li>• <i>Mycosphaerella</i> spp</li> </ul>	20.5 (0.134 lb ai/A)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 20.5 fluid ounces per acre or 0.13 lb ai/acre of GF-3840 AG per application.</li> <li>• <b>DO NOT</b> apply more than 41 fluid ounces per acre (or 0.26 lb ai/acre) of GF-3840 AG per year.</li> <li>• <b>DO NOT</b> apply more than 2 applications of GF-3840 AG per acre per year.</li> <li>• <b>DO NOT</b> apply more than 1 application of GF-3840 AG by air.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days</li> <li>• <b>Preharvest Interval (PHI): 14 days</b></li> <li>• For use in the contiguous US and Hawaii only.</li> </ul> <b>Use Instructions:</b> <ul style="list-style-type: none"> <li>• Make the first application at the beginning of flowering or prior to onset of disease.</li> <li>• Avoid consecutive applications of florylpicoxamid.</li> <li>• Apply at a volume of 15 to 32 gal/A of water, ensuring thorough coverage of the canopy.</li> </ul>		

## Sugar Beet

Disease Common Name	Disease Scientific Name	Product Use Rate per application (fl oz/A)
<ul style="list-style-type: none"> <li>• Cercospora leafspot</li> <li>• Powdery mildew (suppression)</li> </ul>	<ul style="list-style-type: none"> <li>• <i>Cercospora beticola</i></li> <li>• <i>Erysiphe betae</i> (suppression)</li> </ul>	13.7 to 20.5 (0.09 to 0.134 lb ai/A)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 20.5 fluid ounces per acre (0.134 lb ai/acre) of GF-3840 AG per application.</li> <li>• <b>DO NOT</b> apply more than 41 fluid ounces per acre (0.268 lb ai/acre) of GF-3840 AG per year.</li> <li>• <b>DO NOT</b> apply more than 2 applications of GF-3840 AG per acre per year.</li> <li>• <b>Minimum Retreatment Interval:</b> 10 days</li> <li>• <b>Preharvest Interval (PHI): 21 days</b></li> <li>• For use in the contiguous US and Hawaii only.</li> </ul> <b>Use Instructions</b> <ul style="list-style-type: none"> <li>• Apply preventively when conditions are favorable for disease development or based on a forecasting system.</li> <li>• For powdery mildew, apply at the first sign of disease at the highest labeled rate.</li> <li>• Avoid consecutive applications of florylpicoxamid.</li> <li>• Apply at a volume of 15 to 32 gal/A of water, ensuring thorough coverage of the canopy.</li> </ul>		

## Wheat

Disease Common Name	Disease Scientific Name	Product Use Rate per application (fl oz/A)
• Leaf Blight	<i>Septoria tritici</i>	6.8 (0.045 lb ai/A)
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Maximum Application Rate: <b>DO NOT</b> apply more than 6.8 fluid ounces per acre (0.045 lb ai/acre) of GF-3840 AG per application.</li> <li>• <b>DO NOT</b> apply more than 13.7 fluid ounces per acre (0.09 lb ai/acre) of GF-3840 AG per year.</li> <li>• <b>DO NOT</b> apply more than 2 applications of GF-3840 AG per acre per year.</li> <li>• <b>DO NOT</b> apply more than 1 application of GF-3840 AG by air.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days</li> <li>• For use in the contiguous US and Hawaii only.</li> <li>• Apply the last application no later than end of flowering (Zadoks 69, Feekes 10.53).</li> </ul> <b>Use Instructions</b> <ul style="list-style-type: none"> <li>• Begin application of GF-3840 AG before disease development. Apply the product after flag leaf emergence and before end of heading.</li> <li>• Avoid consecutive applications of florypicoxamid. GF-3840 AG should always be applied in mixture with another fungicide from a different FRAC group which provides robust control of <i>Septoria tritici</i>.</li> <li>• Apply at a volume of 9 to 32 gal/A of water, ensuring thorough coverage of the canopy.</li> </ul>		

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### Nonrefillable rigid containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Refillable rigid containers larger than 5 gal:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two

minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable rigid containers larger than 5 gal:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

---

**Terms and Conditions of Use**

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent consistent with applicable law use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

**Warranty Disclaimer**

---

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions for use, subject to the inherent risks set forth below. To the extent consistent with applicable law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

**Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent consistent with applicable law Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent consistent with applicable law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

---

**Limitation of Remedies**

---

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Corteva Agriscience be liable for consequential, incidental, or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

<sup>TM</sup>®Trademarks of Corteva Agriscience and its affiliated companies

EPA accepted \_\_/\_\_/\_\_