



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Registration Division (7505T)

1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

EPA Reg. Number:

62719-780

Date of Issuance:

4/2/25

NOTICE OF PESTICIDE:

☒ Registration

☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Hulk CA

Name and Address of Registrant (include ZIP Code):

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the final printed label for the record before you release the product for shipment.

*Continues page 2*

Signature of Approving Official:

Mindy Ondish, Product Manager 23

Herbicide Branch, Registration Division (7505T)

Date:

4/2/25

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSFs:

- Basic CSF dated 9/28/2023
- Alternate CSF 1 dated 9/28/2023

If you have any questions, please contact Endia Blunt at [blunt.endia@epa.gov](mailto:blunt.endia@epa.gov) or at 202-566-2505.

Enclosure

(Base label for bulk or tote):

**FLORPYRAUXIFEN-BENZYL****GROUP****4****HERBICIDE****Hulk™ CA****HERBICIDE****with Rinskor™ active**

Active Ingredient:

%w/w

florpyrauxifen-benzyl: 2-pyridinecarboxylic acid,

4-amino-3-chloro-6-(4-chloro-2-fluoro-3-

methoxy-phenyl)-

5-fluoro-, phenyl methyl ester.....2.7%

Other Ingredients ..... 97.3%

Total..... 100.0%

**ACCEPTED****04/02/2025**Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 62719-780

Contains 0.21 lb florpyrauxifen-benzyl per gallon.

**Keep Out of Reach of Children****CAUTION****First Aid****If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.**Note to Physician:** Have the product container or label with you when calling a poison control center at 1-800-222-1222 or doctor, or going for treatment. You may also contact Corteva at 1-800-992-5994 day or night, for emergency treatment information.**Precautionary Statements****Hazards to Humans and Domestic Animals****Causes Moderate Eye Irritation****Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.****Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:** When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d-e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from applications is likely to result in damage to sensitive aquatic organisms in water bodies adjacent to the treatment area. Do not contaminate water when disposing of equipment wash waters or rinsate.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Nonrefillable containers 5 gallons or less:****Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers larger than 5 gallons:****Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full

with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable containers larger than 5 gallons:**

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refer to the inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**[Mix or Shake Well Before Use]**

**[Avoid Freezing]**

EPA Reg. No. 62719-780

EPA Est. \_\_\_\_\_

™®Trademarks of Corteva Agriscience and its affiliated companies

**Produced for  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET CONTENTS \_\_\_\_\_**

(Box / Booklet cover / small container base label):

<b>FLORPYRAUXIFEN-BENZYL</b>	<b>GROUP</b>	<b>4</b>	<b>HERBICIDE</b>
------------------------------	--------------	----------	------------------

# Hulk™ CA

## HERBICIDE

with Rinskor™ active

Active Ingredient:	%w/w
florpyrauxifen-benzyl: 2-pyridinecarboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)- 5-fluoro-, phenyl methyl ester.....	2.7%
Other Ingredients .....	97.3%
Total.....	100.0%

Contains 0.21 lb florpyrauxifen-benzyl per gallon.

**Keep Out of Reach of Children**

## CAUTION

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to the inside of label booklet for additional precautionary information including First Aid and Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**[Mix or Shake Well Before Use]**

**[Avoid Freezing]**

EPA Reg. No. 62719-780

EPA Est. \_\_\_\_\_

™® Trademarks of Corteva Agriscience and its affiliated companies

**Produced for  
Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

**NET CONTENTS** \_\_\_\_\_

(Booklet page 1 through end):

**First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**Note to Physician:** Have the product container or label with you when calling a poison control center at 1-800-222-1222 or doctor, or going for treatment. You may also contact Corteva at 1-800-992-5994 day or night, for emergency treatment information.

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION****Causes Moderate Eye Irritation**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.**

**Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:** When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d-e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from applications is likely to result in damage to sensitive aquatic organisms in water bodies adjacent to the treatment area. Do not contaminate water when disposing of equipment wash waters or rinsate.

---

**Directions for Use**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

#### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### Refillable containers larger than 5 gallons:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.



**Nonrefillable containers larger than 5 gallons:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

---

## Product Information

---

Hulk™ CA herbicide is a postemergence herbicide for selective control of susceptible grass, sedge, and broadleaf weeds.

## Use Restrictions

- Do not make more than 2 applications per year.
- Do not use organosilicone surfactants in spray mixtures of this product.
- Do not apply Hulk CA directly to, or otherwise permit Hulk CA to come into direct contact with, carrots, cotton, soybeans, grapes, tobacco, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious injury may occur. Do not permit spray mists containing Hulk CA to drift onto desirable broadleaf plants.
- Do not rotate treated land to highly sensitive crops for 3 months following application.
- Do not apply where runoff or irrigation water may flow directly onto agricultural land to be used for growing highly sensitive crops.
- Do not allow tank mixes of Hulk CA to sit overnight prior to application. See additional tank mix restrictions below.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Do not compost any plant material from treated area.
- Do not tank mix with malathion or methyl parathion. Do not make an application of malathion or methyl parathion within 7 days of an application of Hulk CA. See additional tank mix restrictions below.
- Do not mix with products that contain propanil.
- Do not apply when wind is blowing toward adjacent cotton, carrots, soybeans, corn, grain sorghum, wheat, grapes, tobacco, flowers, ornamental shrubs or trees, or other desirable broadleaf plants.
- Do not apply with aerial application equipment.

## Mixing Instructions

### Use of Adjuvants

Use of an agriculturally approved methylated seed oil adjuvant at a rate of 0.5 pints per acre is allowed to be added to Hulk CA. Do not use pure organosilicone surfactants in spray mixtures of this product. Read and follow all use directions and precautions on methylated seed oil labels.

### Hulk CA – Alone

Fill spray tank to one-half full with water. Start agitation. Add correct quantity of Hulk CA and recommended adjuvant. Continue agitation while filling spray tank to required volume and during application.

### Hulk CA – Tank Mixes

DO NOT TANK MIX ANY PESTICIDE PRODUCT WITH THIS PRODUCT without first referring to the following website for the specific product: [www.hulkcatankmix.com](http://www.hulkcatankmix.com). This website contains a list of active ingredients that are currently prohibited from use in tank mixture with this product.

Continuous agitation is required for tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks.

**Tank Mixing Restrictions**

Only use products in tank mixture with this product that: 1) are registered for the intended use site, application method and timing; 2) are not prohibited for tank mixing by the label of the tank mix product; and 3) do not contain one of the prohibited active ingredients listed on [www.hulkcatankmix.com](http://www.hulkcatankmix.com) website.

Applicators and other handlers (mixers) must access the website within one week prior to application in order to comply with the most up-to-date information on tank mix partners.

Do not exceed specified application rates for respective products or maximum allowable application rates for any active ingredient in the tank mix.

Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels. It is the pesticide user's responsibility to ensure that all products in the mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

When mixing with products that recommend additional adjuvant the total adjuvant should not exceed 0.5 pints of methylated seed oil.

**Tank Mix Compatibility Testing:** When tank mixing Hulk CA with other permitted materials including adjuvants that will be utilized, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately one-half (1/2) hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

**Mixing Order:** Fill the tank one-third (1/3) full with water. Start the agitation. Different formulation types should be added in the following order: dry flowables (DF), wettable powders (WP), aqueous suspensions (AS), flowables (F), or liquids (L). Allow each product type to completely disperse before adding another. Continue agitation and fill tank to three-fourths (3/4) full, add the correct quantity of Hulk CA and mix thoroughly. Finally, add any solution (S) formulations or surfactant, agitate and finish filling. Maintain agitation during filling and during application. If spraying and agitation must be stopped before the tank is empty, suspended materials may settle to the bottom. It is important to re-suspend all of the settled material before continuing application. A sparger agitator is particularly useful for this purpose. Do not allow tank mixes to set overnight.

Carefully follow all mixing instructions for each material added to the tank. Initial dispersion of dry or flowable formulations can be improved by mixing with a small amount of water (slurrying) and pouring the slurry through a 20 to 35 mesh wetting screen in the top of the spray tank. Line screens in the tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

**Clean-Out Procedures for Spray Equipment**

1. Drain any remaining spray mixture from the application equipment, then wash out tank, boom, and hoses with clear water. Drain again.
2. Hose down the interior surfaces of the tank while filling the tank 1/2 full of water.
3. Add commercial tank cleaner, such as household ammonia, at a rate of 1 gallon per 100 gallons of water. Re-circulate for 10 to 20 minutes and spray out the mixture through the boom.
4. Remove all spray nozzles and screens and clean separately.
5. If spray equipment will be used for pesticide application to crops sensitive to Hulk CA, repeat steps 1 through 3.
6. Thoroughly clean exterior surfaces of spray equipment.

Rinsate may be disposed of onsite according to label use directions or at an approved waste disposal facility. Reduced results may occur if water containing soil is used, such as visibly muddy water or water from ponds

and ditches that is not clear.

## **Spray Drift Management**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

### **Ground Application**

- To minimize spray drift from ground application, apply Hulk CA with a nozzle class that ensures coarse or coarser spray (according to ASABE S572).
- Use the minimum boom height based upon the nozzle manufacturer's specifications.
- Apply with the nozzle height recommended by the manufacturer, but no more than 36 inches above the ground or crop canopy.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

## **Spray Drift Advisories**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Ground Boom**

- **Volume:** Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure:** Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle:** Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND

CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### Susceptible Plants

Do not apply under circumstances where spray drift may occur to off-target food or forage, or other plantings. Spray drift may damage or render crops unfit for sale, use, or consumption. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants and tree crops. **Before making an application, please refer to your state's sensitive crop registry (if available) to identify any commercial specialty or certified organic crops that may be located nearby.**

**Do not apply when wind is blowing toward adjacent cotton, carrots, soybeans, grapes, tobacco, vegetable crops, sensitive ornamental plants, or other desirable broadleaf plants.**

Where states have more stringent regulations, they must be observed.

### Buffer Zone

Buffer zones are defined as the minimum downwind distance between the application site and the non-target broadleaf crop. The buffer zones listed below must be followed for applications of Hulk CA:

#### Downwind Buffer to All Non-Target Broadleaf Crops:

Application rate, fl oz/acre	Ground Buffer Zone to Non-Target Broadleaf Crop, ft
5 or less	40
>5-10	85
>10-16	145
>16-21	200

## Application Instructions

### Environmental Conditions and Herbicidal Activity of Hulk CA

Factors for effective weed control with Hulk CA include proper application rate, weed size, daytime and nighttime temperatures, soil moisture prior to and following application, and use of adjuvants. Best weed control results are obtained when Hulk CA is applied to actively growing weeds, when daytime and nighttime temperatures are warm (60 degrees Fahrenheit or more), and soil moisture is adequate to support active weed growth prior to and following application.

- Hulk CA is rainfast in 2 hours.
- Applications made immediately prior to, during, or immediately following periods of large day/night temperature fluctuations or where daytime and nighttime temperatures do not exceed 60 degrees Fahrenheit may decrease weed control.
- Poor weed control and crop injury may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, or hail damage, prior herbicide applications or soils with high salt content.

### Ground Application

Apply in a spray volume of 10 gpa or more when applying by ground. Use coarse or coarser nozzle spray quality per S-572 ASABE standard; see USDA literature or nozzle manufacturer guidelines. Follow nozzle manufacturer's recommendations for nozzle pressure, spacing and boom height to provide a uniform spray pattern. Follow appropriate Spray Drift Management information where drift potential is a concern.

## Resistance Management

This product contains the active ingredient florasulfuron-benzyl, a Group 4 herbicide, based on the mode of action (MOA) classification system of the Weed Science Society of America. The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.

- Start with a clean field, using either a burndown herbicide application or tillage.
- If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.
- Apply full rates of this product for the most difficult to control weed in the field at the specified time to minimize weed escapes.
- Scout fields before and after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent.
- Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective modes of action for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 4 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
  - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - A spreading patch of non-controlled plants of a particular weed species; and
  - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-Group 4 herbicides.
- Avoid making more than two sequential applications of this product and any other Group 4 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, including mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields to reduce weed seed production.

### **Brassica (Cole) Head and Stem Vegetables (Crop Group 5-16)**

Broccoli; Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; cultivars, varieties, and hybrids of these commodities

Hulk CA may be applied as a pre-plant burndown, early postemergence, or fallow (post-harvest) spray.

<b>Weed Control</b>	<b>Rate (fl oz/acre)</b>	<b>Specific Use Directions</b>
Pre-plant burndown	3 to 16	– Application before planting for total burndown control.
Early Postemergence	3 to 16	– Early spring.
Fallow (post-harvest)	3 to 16	– Application post harvesting for total non-selective weed control.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>• Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b>		

- Do not make more than 1 application per year (maximum of 16 fl oz per application).
- Do not apply more than 16 fl oz (0.026 lbs AI) per acre per year.
- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.

At a rate of 3 to 16 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium album</i>
Common mustard	<i>Brassica rapa</i>
Cutleaf evening-primrose	<i>Oenothera laciniata</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed	<i>Conyza canadensis</i>
Mallow	<i>Malva</i> spp.
Nightshade	<i>Solanum</i> spp.
Pigweed	<i>Amaranthus</i> spp.
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Willowherb	<i>Epilobium</i> spp.

### **Brassica (Cole) Leafy Greens (Crop Subgroup 4-16B)**

Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; watercress; cultivars, varieties, and hybrids of these commodities

Hulk CA may be applied as a pre-plant burndown, early postemergence, or fallow (post-harvest) spray.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Pre-plant burndown	3 to 16	– Application before planting for total burndown control.
Early Postemergence	3 to 16	– Early spring.
Fallow (post-harvest)	3 to 16	– Application post harvesting for total non-selective weed control.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>• Do not make more than 1 application per year (maximum of 16 fl oz per application).</li> <li>• Do not apply more than 16 fl oz (0.026 lbs AI) per acre per year.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>• Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> </ul>		

At a rate of 3 to 16 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium album</i>
Common mustard	<i>Brassica rapa</i>
Cutleaf evening-primrose	<i>Oenothera laciniata</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed	<i>Conyza canadensis</i>
Mallow	<i>Malva</i> spp.
Nightshade	<i>Solanum</i> spp.
Pigweed	<i>Amaranthus</i> spp.

Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Willowherb	<i>Epilobium</i> spp.

### **Bulb Vegetables (Crop Group 3-07)**

Chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritillaria, bulb; fritillaria, leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these

Hulk CA may be applied as a pre-plant burndown, postemergence, or fallow (post-harvest) spray.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Pre-plant burndown	3 to 16	– Application before planting for total burndown control.
Early Postemergence	3 to 16	– Early spring.
Fallow (post-harvest)	3 to 16	– Application post harvesting for total non-selective weed control.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>Do not make more than 1 application per year (maximum of 16 fl oz per application).</li> <li>Do not apply more than 16 fl oz (0.026 lbs AI) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> </ul>		

At a rate of 3 to 16 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium album</i>
Common mustard	<i>Brassica rapa</i>
Cutleaf evening-primrose	<i>Oenothera laciniata</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed	<i>Conyza canadensis</i>
Mallow	<i>Malva</i> spp.
Nightshade	<i>Solanum</i> spp.
Pigweed	<i>Amaranthus</i> spp.
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Willowherb	<i>Epilobium</i> spp.

### **Cereals (Barley, Millet, Oats, Rye, Teff, Triticale, and Wheat)**

Hulk CA may be applied as a pre-plant burndown, postemergence (3 leaf to flag leaf emergence), or fallow (post-harvest) spray.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	5 to 16 5 to 21	Post emergence Early spring (3 leaf to flagleaf emergence) Pre-plant burndown and fallow post-harvest
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from</li> </ul>		

abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.

**Crop-Specific Directions:**

- Do not make more than 2 applications per year (maximum of 21 fl oz per application).
- Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.
- **Minimum Retreatment Interval:** 14 days.
- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.

At a rate of 5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium alba</i>
Horseweed	<i>Conyza canadensis</i>
Kochia	<i>Kochia scoparia</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Russian thistle	<i>Salsola tragus</i>
Tall waterhemp	<i>Amaranthus tuberculatus</i>
Wild buckwheat	<i>polygonum convolvulus</i>

### **Citrus (Crop Group 10-10)**

Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uni fruit; cultivars, varieties, and/or hybrids of these

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Refer to application timing for directions.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>• To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>• Do not make more than 2 applications per year (maximum of 21 fl oz per application).</li> <li>• Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>• Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>• Do not apply by air.</li> <li>• Do not spray on top of young trees or expose tree foliage directly to product spray as this could cause undesirable injury.</li> </ul>		

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Annual sowthistle	<i>Sonchus oleraceus</i>
Broadleaf plantain	<i>Plantago major</i>
Burning nettle	<i>Urtica urens</i>



California burclover	<i>Medicago polymorpha</i>
Canada thistle	<i>Cirsium arvense</i>
Coast fiddleneck	<i>Amsinckia menziesii</i>
Common lambsquarters	<i>Chenopodium album</i>
Curly dock	<i>Rumex crispus</i>
Field bindweed	<i>Convolvulus arvensis</i>
Filaree – redstem	<i>Erodium cicutarium</i>
Filaree – whitestem	<i>Erodium moschatum</i>
Geranium - small flower	<i>Geranium pusillum</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy fleabane	<i>Conyza bonariensis</i>
Hoary cress	<i>Cardaria draba</i>
Horseweed	<i>Conyza canadensis</i>
Italian ryegrass	<i>Lolium multiflorum</i>
Knotweed - prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lady's thumb	<i>Persicaria maculosa</i>
London rocket	<i>Sisymbrium irio</i> L.
Mallow - common	<i>Malva neglecta</i>
Mallow - little (cheeseweed)	<i>Malva parviflora</i>
Mustard - black	<i>Brassica nigra</i>
Mustard - common	<i>Brassica rapa</i>
Mustard - wild	<i>Brassica arvensis</i>
Narrow leaf plantain	<i>Plantago lanceolata</i>
Panicle willowherb	<i>Epilobium brachycarpum</i>
Prickly lettuce	<i>Lactuca serriola</i>
Redmaids rockpurslane	<i>Calandrinia umbellata</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Swinecress	<i>Coronopus didymus</i>
Vetch - purple	<i>Vicia americana</i>
Virginia pepperweed	<i>Lepidium virginicum</i>

## Coffee

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Early spring
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>Do not make more than 2 applications per year (maximum of 21 fl oz per application).</li> <li>Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li><b>Minimum Retreatment Interval:</b> 14 days.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>Do not apply by air.</li> <li>Do not spray on top of young trees or expose tree foliage directly to product spray as this could</li> </ul>		

cause undesirable injury.
---------------------------

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Benghal dayflower	<i>Commelina benghalensis</i>
Galinsoga	<i>Galinsoga parviflora</i>
Hairy beggarticks	<i>Bidens pilosa</i>
Lilac tasselflower	<i>Emilia sonchifolia</i>
Morningglory	<i>Ipomoea grandifolia</i>
Radish	<i>Raphanus raphanistrum</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>

### **Corn (Field Corn, Sweetcorn, and Popcorn)**

Hulk CA may be applied as a pre-plant burndown, postemergence (V3 to V6 growth stage) or fallow (post-harvest) spray.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Pre-plant burndown	5 to 16	– Application before planting for total burndown control.
Postemergence	3 to 16	– Early post through 30 inches corn.
Fallow (post-harvest)	5 to 16	– Application post harvesting for total non-selective weed control.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> <li>Crop injury could result from applications &gt;V6 corn stage and yield could be compromised.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>Do not make more than 1 application per year (maximum of 16 fl oz per application).</li> <li>Do not apply more than 16 fl oz (0.026 lbs AI) per acre per year.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>Applications should be made in post emergence at V3–V6 corn stage to avoid potential crop injury.</li> </ul>		

At a rate of 5 to 16 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium album</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>
Giant ragweed	<i>Ambrosia trifida</i>
Hemp sesbania	<i>Sesbania herbacea</i>
Horseweed	<i>Conyza canadensis</i>
Palmer amaranth	<i>Amaranthus palmeri</i>
Pitted morningglory	<i>Ipomoea lacunosa</i>
Prickly sida	<i>Sida spinosa</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Tall waterhemp	<i>Amaranthus tuberculatus</i>
Velvetleaf	<i>Abutilon theophrasti</i>

### **Cotton**

**Application Methods and Equipment:** Hulk CA may be applied as a pre-plant burndown, post-direct (minimum of 6 to 8 inches tall), or fallow (post-harvest) spray.

For post-directed sprays, care must be exercised to avoid spray contact with the cotton leaves. Use rigid precision ground spray equipment and spray shields to prevent spray contact with cotton foliage. Use branch lifters or shields, as necessary, to avoid contact of directed sprays with cotton plant.

**Tank Mixing:** For control of additional broadleaf and grass weeds, Hulk CA may be applied as a postemergence directed spray in tank mix combination with other herbicides registered for postemergence use in cotton. See Mixing Instructions section for more information.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	5 to 16 5 to 16	Apply as a post-directed spray. See Mixing Directions for surfactant recommendations. Pre-plant burndown and fallow post-harvest
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> <li>For post-direct sprays, care must be exercised to avoid spray contact with the cotton leaves.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>Do not make more than 2 applications per year (maximum of 16 fl oz per application).</li> <li>Do not apply more than 32 fl oz (0.052 lbs AI) per acre per year.</li> <li><b>Minimum Retreatment Interval:</b> 14 days.</li> <li>Do not apply to cotton less than 6 inches tall or severe crop injury will result.</li> <li>Exercise care to avoid spray contact with cotton leaves. Leaves accidentally sprayed will exhibit necrotic (dead) spots and may be dropped from the plant.</li> </ul>		

At a rate of 5 to 16 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Hemp sesbania	<i>Sesbania herbacea</i>
Horseweed	<i>Conyza canadensis</i>
Palmer amaranth	<i>Amaranthus palmeri</i>
Pitted morningglory	<i>Ipomoea lacunose</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>

## **Grain Sorghum (Milo)**

Hulk CA may be applied as a pre-plant burndown, postemergence (3 leaf through 7 leaf broadcast post emergence or 8 leaf to boot with drop nozzles) or fallow (post-harvest) spray.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	5 to 16 5 to 21	Post emergence Early spring Pre-plant burndown and fallow post-harvest
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>Do not make more than 1 application per year (maximum of 21 fl oz per application).</li> <li>Do not apply more than 21 fl oz (0.035 lbs AI) per acre per year.</li> </ul>		

- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Do not spray in mid/postemergence applications (> V6 sorghum stage) as it could affect crop yield.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.
- Applications to sorghum should be made at V3-V6 stage to avoid crop injury and potential yield reduction.

At a rate of 5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Common lambsquarters	<i>Chenopodium album</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>
Giant ragweed	<i>Ambrosia trifida</i>
Horseweed	<i>Conyza canadensis</i>
Kochia	<i>Kochia scoparia</i>
Palmer amaranth	<i>Amaranthus palmeri</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>

## Papaya

Hulk CA may be applied as a broadcast spray for early post control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Early spring
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>• To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>• Do not make more than 2 applications per year (maximum of 21 fl oz per application).</li> <li>• Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>• Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>• Do not apply by air.</li> <li>• Hulk CA is phytotoxic to plant foliage. Do not allow direct or indirect applications of Hulk CA to contact any green foliage or injury will occur.</li> </ul>		

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Hairy beggarticks	<i>Bidens pilosa</i>

## Pineapple

Hulk CA may be applied as a postemergence spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Early spring
<b>Precautions:</b>		

- Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.

**Crop-Specific Directions:**

- To enhance the spectrum of control a broad spectrum postemergence herbicide is required.
- Do not make more than 2 applications per year (maximum of 21 fl oz per application).
- Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.
- Minimum Retreatment Interval:** 14 days.
- Preharvest Interval:** Do not apply within 60 days of harvest.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.
- Do not apply by air.
- Hulk CA is phytotoxic to plant foliage. Do not allow direct or indirect applications of Hulk CA to contact any green foliage or injury will occur.

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Hiriyali	
Nutsedge	<i>Cyperus</i> spp.

### **Pome Fruit (Crop Group 11-10)**

Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Refer to application timing for directions.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b>		
<ul style="list-style-type: none"> <li>To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>Do not make more than 2 applications per year (maximum of 21 fl oz per acre per application).</li> <li>Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li><b>Minimum Retreatment Interval:</b> 14 days.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>Do not apply by air.</li> <li>Do not spray on top of young trees or expose tree foliage directly to product spray as this could cause undesirable injury.</li> </ul>		

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Annual sowthistle	<i>Sonchus oleraceus</i>
Broadleaf plantain	<i>Plantago major</i>
Burning nettle	<i>Urtica urens</i>
California burclover	<i>Medicago polymorpha</i>
Canada thistle	<i>Cirsium arvense</i>
Coast fiddleneck	<i>Amsinckia menziesii</i>
Common lambsquarters	<i>Chenopodium album</i>
Curly dock	<i>Rumex crispus</i>

Field bindweed	<i>Convolvulus arvensis</i>
Filaree – redstem	<i>Erodium cicutarium</i>
Filaree – whitestem	<i>Erodium moschatum</i>
Geranium – small flower	<i>Geranium pusillum</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy fleabane	<i>Conyza bonariensis</i>
Hoary cress	<i>Cardaria draba</i>
Horseweed	<i>Conyza canadensis</i>
Italian ryegrass	<i>Lolium multiflorum</i>
Knotweed - prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lady's thumb	<i>Persicaria maculosa</i>
London rocket	<i>Sisymbrium irio</i> L.
Mallow - common	<i>Malva neglecta</i>
Mallow - little (cheeseweed)	<i>Malva parviflora</i>
Mustard - black	<i>Brassica nigra</i>
Mustard – common	<i>Brassica rapa</i>
Mustard - wild	<i>Brassica arvensis</i>
Narrow leaf plantain	<i>Plantago lanceolata</i>
Panicle willowherb	<i>Epilobium brachycarpum</i>
Prickly lettuce	<i>Lactuca serriola</i>
Redmaids rockpurslane	<i>Calandrinia umbellata</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Swinecress	<i>Coronopus didymus</i>
Vetch – purple	<i>Vicia americana</i>
Virginia pepperweed	<i>Lepidium virginicum</i>

### Stone Fruit (Crop Group 12-12)

Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Refer to application timing for directions.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>Do not make more than 2 applications per year (maximum of 21 fl oz per acre per application).</li> <li>Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li><b>Minimum Retreatment Interval:</b> 14 days.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>Do not apply by air.</li> <li>Do not spray on top of young trees or expose tree foliage directly to product spray as this could cause undesirable injury.</li> </ul>		

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Annual sowthistle	<i>Sonchus oleraceus</i>
Broadleaf plantain	<i>Plantago major</i>
Burning nettle	<i>Urtica urens</i>
California burclover	<i>Medicago polymorpha</i>
Canada thistle	<i>Cirsium arvense</i>
Coast fiddleneck	<i>Amsinckia menziesii</i>
Common lambsquarters	<i>Chenopodium album</i>
Curly dock	<i>Rumex crispus</i>
Field bindweed	<i>Convolvulus arvensis</i>
Filaree – redstem	<i>Erodium cicutarium</i>
Filaree – whitestem	<i>Erodium moschatum</i>
Geranium – small flower	<i>Geranium pusillum</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy fleabane	<i>Conyza bonariensis</i>
Hoary cress	<i>Cardaria draba</i>
Horseweed	<i>Conyza canadensis</i>
Italian ryegrass	<i>Lolium multiflorum</i>
Knotweed - prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lady's thumb	<i>Persicaria maculosa</i>
London rocket	<i>Sisymbrium irio L.</i>
Mallow - common	<i>Malva neglecta</i>
Mallow - little (cheeseweed)	<i>Malva parviflora</i>
Mustard - black	<i>Brassica nigra</i>
Mustard – common	<i>Brassica rapa</i>
Mustard - wild	<i>Brassica arvensis</i>
Narrow leaf plantain	<i>Plantago lanceolata</i>
Panicle willowherb	<i>Epilobium brachycarpum</i>
Prickly lettuce	<i>Lactuca serriola</i>
Redmaids rockpurslane	<i>Calandrinia umbellata</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Swinecress	<i>Coronopus didymus</i>
Vetch – purple	<i>Vicia americana</i>
Virginia pepperweed	<i>Lepidium virginicum</i>

## Sugarcane

Hulk CA may be applied as a broadcasted spray for control of emerged weeds or as a ripener applied prior to sugarcane harvest.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	5 to 21	Early spring to early summer  For ripener, apply at a rate of 1 to 3 fl oz/acre at late summer prior to sugarcane harvest.
<b>Precautions:</b>		

- Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.

**Crop-Specific Directions:**

- Do not make more than 2 applications per year (maximum of 21 fl oz per application).
- Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.
- Minimum Retreatment Interval:** 14 days.
- Preharvest Interval:** Do not apply within 60 days of harvest.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.
- Apply at least 14 days prior to sugarcane harvest as a ripener.

At a rate of 5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Black nightshade	<i>Solanum nigrum</i>
Itchgrass	<i>Rottboellia cochinchinensis</i>
Purple nutsedge	<i>Cyperus rotundus</i>
Red morningglory	<i>Ipomoea coccinea</i>
Yellow nutsedge	<i>Cyperus esculentus</i>

### Tree Nuts (Crop Group 14-12)

African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	10.5 to 21	Refer to application timing for directions.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b>		
<ul style="list-style-type: none"> <li>To enhance the spectrum of control a broad spectrum postemergence herbicide is required.</li> <li>Do not make more than 2 applications per year (maximum of 21 fl oz per application).</li> <li>Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li><b>Minimum Retreatment Interval:</b> 14 days.</li> <li><b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>Do not apply by air.</li> <li>Do not spray on top of young trees or expose tree foliage directly to product spray as this could cause undesirable injury.</li> </ul>		

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Annual sowthistle	<i>Sonchus oleraceus</i>
Broadleaf plantain	<i>Plantago major</i>
Burning nettle	<i>Urtica urens</i>
California burclover	<i>Medicago polymorpha</i>



Canada thistle	<i>Cirsium arvense</i>
Coast fiddleneck	<i>Amsinckia menziesii</i>
Common lambsquarters	<i>Chenopodium album</i>
Curly dock	<i>Rumex crispus</i>
Field bindweed	<i>Convolvulus arvensis</i>
Filaree – redstem	<i>Erodium cicutarium</i>
Filaree – whitestem	<i>Erodium moschatum</i>
Geranium - small flower	<i>Geranium pusillum</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy fleabane	<i>Conyza bonariensis</i>
Hoary cress	<i>Cardaria draba</i>
Horseweed	<i>Conyza canadensis</i>
Italian ryegrass	<i>Lolium multiflorum</i>
Knotweed - prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lady's thumb	<i>Persicaria maculosa</i>
London rocket	<i>Sisymbrium irio</i> L.
Mallow - common	<i>Malva neglecta</i>
Mallow - little (cheeseweed)	<i>Malva parviflora</i>
Mustard - black	<i>Brassica nigra</i>
Mustard - common	<i>Brassica rapa</i>
Mustard - wild	<i>Brassica arvensis</i>
Narrow leaf plantain	<i>Plantago lanceolata</i>
Panicle willowherb	<i>Epilobium brachycarpum</i>
Prickly lettuce	<i>Lactuca serriola</i>
Redmaids rockpurslane	<i>Calandrinia umbellata</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Swinecress	<i>Coronopus didymus</i>
Vetch - purple	<i>Vicia americana</i>
Virginia pepperweed	<i>Lepidium virginicum</i>

### **Tropical and Subtropical, Small fruit, edible peel (Crop Subgroup 23A) including olives**

Acerola; African plum; agritos; almondette; appleberry; arbutus berry; bayberry, red; bignay; breadnut; cabeluda; carandas-plum; Ceylon iron wood; Ceylon olive; cherry-of-the-Rio-Grande; Chinese olive, black; Chinese olive, white; chirauli-nut; cocoplum; desert-date; false sandalwood; fragrant manjack; gooseberry, Abyssinian; gooseberry, Ceylon; gooseberry, otaheite; governor's plum; grumichama; guabiroba; guava berry; guava, Brazilian; guava, Costa Rican; guayabillo; illawarra plum; Indian-plum; Jamaica-cherry; jambolan; kaffir-plum; kakadu plum; kapundung; karanda; lemon aspen; mombin, yellow; monos plum; mountain cherry; olive; persimmon, black; pitomba; plum-of-Martinique; rukam; rumberry; sea grape; sete-capotes; silver aspen; water apple; water pear; water berry; wax jambu; cultivars, varieties, and hybrids of these commodities

Hulk CA may be applied as a broadcasted spray for control of emerged weeds.

<b>Weed Control</b>	<b>Rate (fl oz/acre)</b>	<b>Specific Use Directions</b>
Postemergence	10.5 to 21	Early spring – summer (Feb - August)
<b>Precautions:</b> <ul style="list-style-type: none"> <li>Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		

**Crop-Specific Directions:**

- To enhance the spectrum of control a broad spectrum postemergence herbicide is required.
- Do not make more than 2 applications per year (maximum of 21 fl oz per application).
- Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.
- **Minimum Retreatment Interval:** 14 days.
- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.
- Do not apply by air.
- Do not spray on top of young trees or expose tree foliage directly to product spray as this could cause undesirable injury.

At a rate of 10.5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Annual sowthistle	<i>Sonchus oleraceus</i>
Broadleaf plantain	<i>Plantago major</i>
Burning nettle	<i>Urtica urens</i>
California burclover	<i>Medicago polymorpha</i>
Canada thistle	<i>Cirsium arvense</i>
Coast fiddleneck	<i>Amsinckia menziesii</i>
Common lambsquarters	<i>Chenopodium album</i>
Curly dock	<i>Rumex crispus</i>
Field bindweed	<i>Convolvulus arvensis</i>
Filaree – redstem	<i>Erodium cicutarium</i>
Filaree – whitestem	<i>Erodium moschatum</i>
Geranium - small flower	<i>Geranium pusillum</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy fleabane	<i>Conyza bonariensis</i>
Hoary cress	<i>Cardaria draba</i>
Horseweed	<i>Conyza canadensis</i>
Italian ryegrass	<i>Lolium multiflorum</i>
Knotweed - prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lady's thumb	<i>Persicaria maculosa</i>
London rocket	<i>Sisymbrium irio L.</i>
Mallow - common	<i>Malva neglecta</i>
Mallow - little (cheeseweed)	<i>Malva parviflora</i>
Mustard - black	<i>Brassica nigra</i>
Mustard - common	<i>Brassica rapa</i>
Mustard - wild	<i>Brassica arvensis</i>
Narrow leaf plantain	<i>Plantago lanceolata</i>
Panicle willowherb	<i>Epilobium brachycarpum</i>
Prickly lettuce	<i>Lactuca serriola</i>
Redmaids rockpurslane	<i>Calandrinia umbellate</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Swinecress	<i>Coronopus didymus</i>
Vetch - purple	<i>Vicia americana</i>
Virginia pepperweed	<i>Lepidium virginicum</i>

---

**Pre-Plant Burndown and Fallow Cropland Application**


---

**Application Information**

See individual crop sections of this label for additional directions, precautions, and restrictions.

**Use Rate**

Apply 5 to 21 fluid ounces of Hulk CA per acre. Unless otherwise specified, sequential treatment of Hulk CA may also be made ahead of certain crops as specified below, provided the total amount of Hulk CA during the crop season does not exceed 36 oz per acre.

**Application Timing**

Hulk CA may be used as a pre-plant burndown or fallow treatment in the spring, summer or fall when the majority of weeds have emerged and are actively growing.

**Tank Mixtures**

Hulk CA may be tank mixed with one or more registered products according to the specific tank mixing instructions in this label and respective product labels. Read and follow all manufacturers' label recommendations for the companion herbicide. Always follow the most restrictive label use directions.

**Crop Specific Information**

Apply Hulk CA as a pre-plant burndown treatment prior to planting the crops listed below.

**Corn (Field, Popcorn, and Sweetcorn)**

Hulk CA may be used as a pre-plant burndown treatment at least 14 days prior to planting at a rate of 5 to 16 ounces of product per acre. Sequential treatments of Hulk CA may also be applied during one cropland season provided the total amount does not exceed 32 ounces per acre per crop season; for example, 16 ounces in the fall followed by 16 ounces in the spring. Separate sequential applications by at least 14 days.

**Cotton**

Hulk CA may be used as a pre-plant burndown treatment at least 14 days prior to planting at a rate of 5 to 16 ounces of product per acre. Sequential treatments of Hulk CA may also be applied during one cropland season provided the amount does not exceed 32 ounces per acre per pre-plant crop season; for example, 16 ounces in the fall followed by 16 ounces in the spring. Separate sequential applications by at least 14 days.

**Soybean**

Hulk CA may be used as a pre-plant burndown treatment at least 14 days prior to planting at a rate of 5 to 10.5 ounces of product per acre.

**Barley, Triticale, and Wheat (including spring, winter, and durum)**

Apply 5 to 16 fl oz of Hulk CA per acre as a burndown treatment to wheat (including durum), barley, and triticale to control emerged weeds prior to or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.

Applications should be made to the main flush of actively growing weeds. Include a methylated seed oil or crop oil concentrate at 1% v/v. If another herbicide is tank mixed with Hulk CA to increase the broadleaf or grass weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from application made to seedling weeds.

Warm, moist growing conditions promote active weed growth and enhance the activity of Hulk CA by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

**Tank Mixes for Corn (Field, Popcorn, and Sweetcorn), Cotton, Sorghum, Soybean, and Sugarcane**

Hulk CA may be used as a pre-plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product, such as glyphosate, glufosinate, or 2,4-D. See the Mixing Instructions section of this label for additional information. Read and follow all manufacturers' label recommendations for the companion herbicide.

**Use Restrictions**

- Do not apply within 14 days prior to planting of listed crops.
- Do not apply more than 42 oz of Hulk CA per acre per growing season.

---

### Non-Cropland Areas

---

Non-food producing, non-cultivated agricultural, or non-agricultural areas including highway and utility rights-of-way, industrial sites, tank farms, storage areas, airports, fencerows not adjacent to food/feed crop fields and farmsteads

Weed Control	Rate (fl oz/acre)	Specific Use Directions
Postemergence	5 to 21	Apply when weeds are actively growing (early spring through late summer)
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Poor weed control may result from application of Hulk CA made to plants under stress from abnormally hot or cold weather; environmental conditions such as drought, hail damage, or prior herbicide applications.</li> </ul>		
<b>Crop-Specific Directions:</b> <ul style="list-style-type: none"> <li>• Do not make more than 2 applications per year (maximum of 21 fl oz per application).</li> <li>• Do not apply more than 42 fl oz (0.07 lbs AI) per acre per year.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days.</li> <li>• Make applications in a minimum of 10 gallons per acre (gpa) for ground applications.</li> <li>• Do not apply Hulk CA in any residential setting.</li> <li>• Allow 30 days or more between treatments.</li> </ul>		

At a rate of 5 to 21 fl oz/acre the following weeds are either controlled or suppressed:

Common Name	Scientific Name
Alligatorweed	<i>Alternanthera philoxeroides</i>
Ammannia (redstem)	<i>Ammannia coccinea</i>
Arrowhead / bulltongue / grassy arrowhead	<i>Sagittaria</i> spp.
Ducksalad	<i>Heteranthera limosa</i>
Eclipta	<i>Eclipta prostrate</i>
Falsepimpernel, low	<i>Lindernia dubia</i>
Horseweed	<i>Conyza</i> spp.
Roundleaf mudplantain	<i>Heteranthera reniformis</i>
Spreading dayflower	<i>Commelina diffusa</i>
Wild carrot	<i>Daucus carota</i>
Wild plantain	<i>Plantago</i> spp.

---

### Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

### Warranty Disclaimer

---

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions for use, subject to the inherent risks set forth below. To the extent consistent with applicable law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent consistent with applicable law, Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent consistent with applicable law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

---

**Limitation of Remedies**

---

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Corteva Agriscience be liable for consequential, incidental, or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

<sup>TM</sup>®Trademarks of Corteva Agriscience and its affiliated companies

EPA accepted \_\_/\_\_/\_\_