

DRAFT LABEL

FICSAN® TABLETS

SELECTIVE INJECTION HERBICIDE TABLETS

by Forestry Injection Company FIC, AB

Tablets containing water soluble herbicide for insertion into broadleaf trees and stumps

Kills undesirable trees and stumps -- roots and all.

ACCEPTED
NOV 13 1990
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62802-E

DMC
2001-2
12-1-90

Active Ingredient:	Hexazinone; [3-Cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione]	72%
Inert Ingredients:	28%
TOTAL	100%

Contents: 300 tablets*
Each tablet contains 360 mg of the active ingredient

Read the entire label before using this product.

Use only according to label instructions.

EPA Reg. No. 62802-E

EPA Est. No. 62957-SE-001

**FOR PROFESSIONAL USE ONLY
REFORMULATION OR REPACKAGING IS PROHIBITED**

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Causes eye injury. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with soap and water. Get medical attention.

* Alternate label to read 100 tablets.

ENVIRONMENTAL HAZARDS

Do not contaminate water or wetland (swamps, bogs, marshes or potholes).

STORAGE & DISPOSAL

Store product in a dry place in original container only, away from other pesticides, fertilizer, food or feed. Do not contaminate water, food or feed by storage or disposal. Dispose of according to local regulations.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FICSAN TABLETS are recommended for killing trees in forests, parks, fence rows; in highway and railroad rights-of-ways; and in industrial and residential sites. It may be used for killing trees standing in water and in wetlands.

FICSAN TABLETS will kill the following trees:

- | | |
|----------------|-----------|
| Ash | Oak |
| Aspen, quaking | Sourwood |
| Birch | Sumac |
| Dogwood | Sweet gum |
| Elm | Willow |
| Hawthorn | |

For effective results, FICSAN TABLETS must be inserted such that the content of the active ingredient is in direct contact with the tree's phloem tissue.

Time Required for Effective Killing of Trees and Sprout Control:

At least one growing season is required for killing of trees. For root sprout control, 2 - 3 growing seasons are required.

BEST AVAILABLE COPY

Use Precautions:

- 1) Do not apply to trees when their sap is frozen.
- 2) Do not apply to trees that may be later standing in water above the point of insertion.
- 3) Apply to birch trees only when in full foliage.
- 4) If used for sprout control of stumps, apply within three weeks of felling the trees.

Method of Application:

Insert FICSAN® TABLETS into trees and stumps using the FIC® hollow punch to open a hole in the tree or stump and the FIC® Injector to insert the tablet into the hole. Holes should angle slightly downward (about 30° below horizontal) and should extend to a depth of 0.8 - 1.2 inches past the cambium (the first living tissue inside the bark). On thick-barked species, it may be necessary to remove excess bark with an axe to permit proper hole depth. To remove the FIC® hollow punch from the tree or stump, turn it downward and then push outward and upward.

For trees 10 inches or less in diameter at breast height, insert one (1) tablet every 4 inches of tree circumference. For trees greater than 10 inches in diameter at breast height, insert one (1) tablet every 2.5 inches of tree circumference.

Registrant and Manufacturer: Forestry Injection Company FIC AB
Kungsgatan 18 - 632 21, Eskilstuna, SWEDEN
Telephone +46-16-11 47 00

Sold by: Forestry Injection Company FIC Inc.
291 Hwy. 51-C-7, P.O. Box 1336
Ridgeland, Mississippi 39157
Telephone: (601) 856-6835

NOTICE OF WARRANTY

Forestry Injection Company FIC, AB warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of FIC. In no case shall FIC be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. FIC MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

BEST AVAILABLE COPY