

PM 22 63239-15 pslf



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
63239-15

Date of Issuance:
JUL 31 1995

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
63239-15

Name and Address of Registrant (include ZIP Code):

APC Holdings Corporation
2215 West Street - 2nd Floor
Germantown, TN 38138

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

- a. Add the phrase, "EPA Reg. No. 63239-15".
- b. In the Environmental Hazards Section delete "For terrestrial uses,". Refer to PR Notice 93-8.

Signature of Approving Official:

Date:

JUL 31 1995

19/9

page 2
EPA Reg. No. 63239-15

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

James M. Stone
Acting Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

APC HOLDINGS
DSMA LIQUID

DSMA Plus Surfactant
For Selective Postemergence Weed Control

ACTIVE INGREDIENT:	
Disodium Methanearsonate Anhydrous*	21.8%
INERT INGREDIENTS:	78.2%
TOTAL:	100.0%

Total Arsenic (as elemental)
all in water soluble form.....8.9%

*Product contains 3.6 lbs. Disodium Methanearsonate
Hexahydrate equivalent per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements.

Net Contents:

ACCEPTED
with COMMENTS
in EPA Letter Dated:

the Federal Insecticide
and Fungicide Act
and for the purpose
of the EPA Reg. No.

63239-15

MANUFACTURED FOR:
APC HOLDINGS
P.O. Box 13347
Memphis, TN 38113-0347

EPA Reg. No. 63239-
EPA Est. No. 19713-MS-1
March 31, 1995

4/1

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Never give anything by mouth to an unconscious person. Seek medical attention.

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Seek medical attention.

IF ON SKIN: Immediately wash skin with plenty of soap and water. If irritation develops, seek medical attention.

IF INHALED: Remove victim to fresh air.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, and chemical-resistant apron when cleaning equipment, mixing or loading.

For exposures in enclosed areas: A respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

For exposure outdoors: Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not feed treated foliage to livestock or graze treated areas. Do not spray or allow drift onto vegetables, ornamentals or other desirable plants. Do not apply when weather conditions favor drift from target areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION

DSMA LIQUID is a postemergence herbicide for use in cotton, turf and non-crop areas. DSMA LIQUID is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Local conditions and recommendations vary; consult local agricultural experiment station or extension service weed specialists for recommendations in your area.

MIXING INSTRUCTIONS

DSMA LIQUID is completely water soluble. Fill the spray tank with half the required amount of water and, with pump or agitator operating, add the recommended amount of DSMA LIQUID and continue filling tank with balance of water. Do not store spray solution in tank overnight. Clean application equipment thoroughly after use. Although DSMA LIQUID is only moderately corrosive, do not use in galvanized or aluminum equipment.

APPLICATION METHODS:

DSMA LIQUID should be applied with a low-volume, properly calibrated sprayer having satisfactory pumping and bypass action. ADJUST NOZZLES TO KEEP SPRAY OFF COTTON FOLIAGE AND IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS. Adequate coverage is very important for effective control. Apply during warm weather when weeds are in an active stage of growth.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation equipment.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralls

Waterproof gloves

Chemical-resistant footwear plus socks

Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard of agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and domestic animals off treated areas until this material has been washed into the soil.

COTTON:

DSMA LIQUID is useful as a directed postemergence spray for controlling weeds in cotton such as barnyardgrass, cocklebur, smooth and large crabgrass, dallisgrass, goosegrass, johnsongrass, nutsedge, puncturevine, and sandbur. DO NOT feed treated foliage to livestock or graze treated areas.

Preplant or postplant up to cracking application on cotton:

A single ground or aircraft application of DSMA LIQUID can be made to prepared cotton seedbeds when planting has been delayed and weeds have emerged, or as a post-plant treatment, but no later than initial cracking of soil in field before emergence of cotton.

Planting of cotton may immediately follow the preplant application. Mix at a rate of 1 gallon of DSMA LIQUID in 40 gallons of water for broadcast ground application or in 5 to 10 gallons of water for aircraft application and apply to one acre using a properly calibrated sprayer.

Postemergent, over-the-top application on cotton:

A single ground or aircraft application may be made as an over-the-top broadcast spray to actively growing cotton as a salvage operation when cotton is 3 to 6 inches high or up to early first square stage, whichever occurs first. Mix DSMA LIQUID at a rate of 3 to 4 pints in 40 gallons of water for ground application, or in 5 to 10 gallons of water for aircraft application and apply to one acre using a properly calibrated sprayer. Apply as an over-the-top broadcast spray only when cotton is 3 to 6 inches high or up to early first square stage, whichever occurs first.

DO NOT apply as an over-the-top broadcast spray after cotton is more than 6 inches high or after early first square stage. Slight burning or reddish discoloration of cotton foliage may occasionally be seen following recommended treatment.

Directed application:

Apply as a directed spray at the rate of 40 gallons per acre of a solution containing 1 gallon of DSMA LIQUID in 40 gallons of water when weeds are small. For band applications, apply 1 gallon of above diluted spray per acre for each 1 inch band width to be treated of cotton grown on 40 inch row spacing. A second or repeat application, if needed, should be timed about 1 to 3 weeks after the first application. Keep spray off cotton foliage.

Apply Only When Cotton is 3 Inches High to First Bloom. Do Not Apply After First Bloom.

Slight burning or reddish discoloration of cotton foliage may occasionally be seen following recommended treatment; however, cotton plants will develop normally and yield will not be affected.

LAWN AND ORNAMENTAL TURF:

DSMA LIQUID can be used for selective control of bahiagrass, barnyardgrass, chickweed, smooth and large crabgrass, dallisgrass, nutsedge, sandbur and wood sorrel with little or no injury to well established, actively growing turfgrasses. Mow turfgrass to height of 1 to 1 1/2 inches before treatment. Mix 4 fluid ounces (8 tablespoons) of DSMA LIQUID in 2 1/2 gallons of water for application to an area of 1,000 square feet. Application should be uniform and thorough to adequately wet all undesirable plants. Two or more repeat treatments at 14 day intervals may be necessary. Make applications during warm weather when the temperature is between 80 and 90°F. DO NOT water turf for at least 24 hours after application. Turfgrasses may be temporarily discolored. Bermudagrass, bluegrass and zoysiagrass have shown tolerance to properly applied DSMA LIQUID. Injury may result if applied to bentgrass and fescues. DO NOT apply to St. Augustinegrass, carpetgrass, centipedegrass or to Dichondra. DO NOT reseed until 2 weeks after last application. DO NOT apply with hose-end applicators.

8/13

AGRICULTURAL PLANTINGS:

DSMA LIQUID is useful as a directed application in non-bearing vineyards and in non-bearing deciduous fruit, nut and citrus orchards such as almond, apple, apricot, cherry, grapefruit, lemon, lime, orange, peach, pear, plum, prune and tangerine orchards. It should be applied at the rate of 3 1/2 to 7 quarts per acre. Mix at the rate of 3 1/2 quarts of DSMA LIQUID to 50 gallons of water. Apply as a directed spray in interspaces and around base of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required. However, do not exceed 3 applications per year.

DO NOT allow spray solution to contact leaves, stem or bark. Use a shield, if necessary, for nursery plantings or young trees. DO NOT apply around trees or vines from which fruit will be harvested within one year of treatment.

NON-CROP:

DSMA LIQUID is effective in controlling barnyardgrass, cocklebur, smooth and large crabgrass, dallisgrass, goosegrass, johnsongrass, nutsedge and many similar weeds on drainage ditchbanks, fence rows, rights-of-way, storage yards and similar non-crop areas. Application should be made when weeds are small and conditions are favorable for good weed growth. Mix at a rate of 1 gallon of DSMA LIQUID in 40 gallons of water.

Spray undesirable vegetation thoroughly to point of run-off. Adequate coverage and complete wetting of foliage is important for effective control. Repeat applications may be necessary if regrowth occurs.

DO NOT contaminate waters used for domestic purposes, by animals, including wildlife and aquatic life, or for irrigation purposes.

6

349

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS:

Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not stack over 2 pallets high. Move containers by handles or in cases. Do not move containers from one area to another unless they are securely sealed. Keep containers tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizers to avoid contamination. This material is incompatible with Aluminum and zinc. Do not expose this concentrate or the spray solution to aluminum or galvanized metal. Store in original containers only. If the contents are leaking or material is spilled, follow these steps while wearing Personal Protective Equipment.

- 1) Contain spill, absorb with material such as sawdust, clay granules or dirt.
- 2) Collect and place in suitable containers for disposal.
- 3) Wash area with water and soap to remove remaining pesticide.
- 4) Follow washing with clean water rinse.
- 5) Do not allow run-off to enter sewer or contaminate water supplies.
- 6) Dispose of wastes as indicated below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage, and handling of this material not in strict accordance with directions given herewith.

In no case shall APC Holdings or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by APC Holdings and is accepted as such by the Buyer.

9