



## For Crabgrass Control

ACTIVE INGREDIENT:

Calcium Acid Methanearsonate	10.3%
OTHER INGREDIENTS:	89.7%
TOTAL:	100.0%
Total Arsenic as elemental, all in water soluble tom, is 5.09	<b>6</b> .

One gallon contains 1 pound of the active ingrediem CANNA.

# KEEP OUT OF REACH OF CHILDREN CAUTION

# See Below for Practical Treatment (First Lid) SHAKE WELL BEFORE USING

EPA Reg. No. 63239-15

EPA Est. No. 19713-MS-1

Net Contents:

### PRACTICAL TREATMENT (FIRST AID)

IF SWALLOWED: Call a consistion immediately. Tork it or 2 gasses of water and induce vomiting by touching back of Index with Index. Do not induce vomiting or give anything by mouth to at inconscious betson. IF ON SKIN: Wash with plenty of water. Get medical attention if initiation persists.

IF INHALED: Remove victim to fresh air. Apply artifical respirator, preferably mouth to mouth, if indicated.

IF IN EYES: Flush with plenty of water. Get medical attention if initiation persists.

#### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animats

CAUTION: Hamful if swallowed or inhaled. Avoid treathing stray mist. (Coarse sprays are less likely to drift.) Avoid contact with eves, skin or clothing. Causes eye initation. Wear protective potting when handling or applying this product, including long pans, long steave shirt and impermeable gloves and boots. Wash thoroughly with steap and water after handling. Remove contaminated clothing and wast before reuse. Keep children and domestic animals of: reatted areas until washed into soil.

#### ENVIRONMENTAL HAZARDS

Do not contaminate waters used by wildlife, aquate life, domestic animals or waters used by imigation. Do not apply directly to water, or to areas where surface water is present or to interlead areas zelow the mean high water mark. Do not contaminate water by disanime of sequement or disposal of wastes.

#### PRODUCT INFORMATION

CALAR (Calcium Acid Methanearsonate) is a post-energence ungrass herbicide. Used in the control of Crabgrass, Barmyarograss, Witchgrass and Foxtail. May be used on Bermuda, merion blue, Kenacky blue, red top, fescue and zoysia grasses. Do not use on St. Augustime grass.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product n a manner inconsistent with its labeling. Not for use on turf beeing gower flor sate or other commercial use as sod, or for commercial seed production, or for research purposes. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not use on grass grown for seed. Do not allow livestock to graze on treated areas. Do not use drainage water from turf treated with this product for crop irrigation.

This product is most effective when applied to weeds during hot summer weather, but has worked effectively at temperatures as low as 70°F on days with high humidity. Apply to weeds when actively growing.

Rate of Application at 80° F and Above		
Rate	Amount of Water	Area Covered
8 fluid ounces	5 to 10 Gallons	1.000 spare feet
21/2 Gallons	50 to 100 Gallons	1 Acre

This product may be used on bentgrass greens at one-half ( $\frac{1}{2}$ ) the rate as recommended above to treat the same area. For temperatures below 80° F, as in Spring and Fall, use twice the rate as specified in the table above to treat grasses other than bent. The amount of water to cover a given area remains unchanged regardless of the quantity of chemicals used. Thoroughly mix by agitation or recirculating solution in spray tank prior to application. To decrease runoff and increase effectiveness of chemicals through greater leaf absorption, add to the mixture one quant (1 quart) of a suitable surfactant to each 50 galions of water used.

- Be sure soil is moist before treatment. During dry spells, water thoroughly prior to application.
- (2) Use a fixed boom power sprayer property calibrated to a uniform speed and rate of delivery.
- (3) Normal rainfall following application will not adversely affect results. Resume keeping soil moist day after application and especially during hot, dry weather.
- (4) Two applications, 5 to 7 days apart, are usually required for complete killing of weeds. New turt should not be treated until the third mowing. If more than the recommended rate is used, most grasses including zoysia will show temporary discoloration. Normal color will return in a few days.

(5) Do not apply with hose-end applicators.

Do not spray or allow to drift onto desirable flowers, trees, shrubs, vegetables, or into lakes or streams. Coarse sprays are less likely to drift. Do not store near fertilizer or seeds.

Clean equipment after use by flushing with water.

# STORAGE AND DISPOSAL

STORAGE: Do not store near feed or food products. Do not contaminate water, food or feed by storage and disposal. PESTICIDE DISPOSAL: Pesticide wastes are toxic, improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use excryding to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Thoractization recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and Local authoration by burning. If burned, stay out of smoke.

\*\*\*\* \*\*\*\* T.


Manufactured For: APC Holdings P.D. BDs 13947 • MEMPHIS, TN 38113-0347

# WARRANTY-CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

BEFORE USING ANY HERBICIDE, STOP AND READ THE LABEL.



ż

1

92

0