

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE: <u> x </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number:	Date of Issuance:
	63281-9	10-9-97
	Term of Issuance: Conditional	
		Name of Pesticide Product:
		BC II Phenolic
Name and Address of Registrant (include ZIP Code).		
Huntington Laboratories, Inc. 970 East Tipton Street Huntington, IN 46750 AttN: Ms. Sally Hayes - Manager, Regulatory Affairs		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 63281-9". 3. Submit two copies of the revised final printed label for the record. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.</p> <p>A stamped copy of the label is enclosed for your records.</p>		
Signature of Approving Official:		Date:
		OCT 10 1997

-2-

If you have any questions, please contact LaVerne Dobbins at (703) 308-6378.

Sincerely,



Portia M. Jenkins
Acting Product Manager (34)
Regulatory Management Branch II
Antimicrobials Division (7510W)

Enclosure:

376

BC II PHENOLIC
Germicidal Detergent

Cleans, Disinfects, Deodorizes
Pleasant Fragrance

PSEUDOMONACIDAL STAPHYLOCIDAL TUBERCULOCIDAL *VIRUCIDAL FUNGICIDAL

BC II PHENOLIC is a multi-purpose germicidal detergent proven effective the AOAC Use-Dilution Method in 400 ppm hard water (calculated as CaCO₃) in the presence of 5% organic bioload (Fetal Bovine Serum). BC II PHENOLIC's formula disinfects, cleans and deodorizes in one labor-saving step. BC II PHENOLIC is effective against the following pathogenic organisms:

- | | |
|---------------------------|------------------------------------------------|
| Pseudomonas aeruginosa | Shigella flexneri |
| Salmonella choleraesuis | Streptococcus pyogenes |
| {Staphylococcus aureus | Enterococcus faecalis |
| Enterobacter aerogenes | **Enterococcus faecalis (vancomycin resistant) |
| {Enterobacter cloacae | |
| **Enterococcus sp. | *HIV-1 (AIDS Virus) |
| Escherichia coli | *Rubella (German measles) |
| Klebsiella pneumoniae | *Adenovirus Type 2 |
| Mycobacterium bovis | *Herpes Simplex Type 1 |
| {Proteus mirabilis | *Human rhinovirus Type 38 |
| Proteus vulgaris | *Influenza Type A/Michigan |
| Providencia rettgeri | *Polio 1 virus |
| Pseudomonas cepacia | *Vaccinia |
| **Pseudomonas maltophilia | *Newcastle Disease Virus |
| Salmonella typhi | |
| Salmonella typhimurium | Candida albicans |
| Serratia marcescens | Trichophyton mentagrophytes |

*Tested against antibiotic resistant strain only
{Tested against standard ATCC strains and antibiotic resistant strains.

ACTIVE INGREDIENTS:

o-phenylphenol.....	3.20%
o-benzyl-p-chlorophenol.....	5.00%
p-tert-amyphenol.....	1.60%
INERT INGREDIENTS:.....	90.20%
TOTAL.....	100.00%

KEEP OUT OF REACH OF CHILDREN
DANGER

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 10 1997

(See side panel for additional precautionary and practical treatment statements)

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

Net Contents: 1 U. S. Gal./3.78 L
5 U.S Gal./18.9 L

632819

EPA Reg. No. 63281-O
EPA Est. 303-IN-1(L), 1677-IL-2(J), 1677-NJ-1(W), 1677-TX-1(D), 1677-GA-1(M), 1677-MN-1(P), 1677-CA-1(S), 1677-PR-1(B), 1677-OH-1(H)
The superscript refers to the first letter of the date code.

Huntington Professional Products, A Service of Ecolab Inc.
370 Wabasha Street
St. Paul, MN 55102

466

STATEMENT OF PRACTICAL TREATMENT

In case of skin contact, wash thoroughly with soap and water. In case of eye contact, immediately flush eyes with water for 15 minutes and get prompt medical attention. If swallowed, promptly drink a large quantity of water. Avoid alcohol. Call a physician immediately.

FOR EMERGENCY MEDICAL INFORMATION, CALL 1-800-328-0026

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: CORROSIVE: Concentrate causes eye and skin damage. May be absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Harmful if swallowed. Avoid contamination of food, water or feed.

ENVIRONMENTAL HAZARDS: (for 5 gallon and larger only)

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: (1 gal) Do not reuse empty containers. Wrap container and put in trash. (5 gal plastic) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 10 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended for the pesticide,
registered under EPA Reg. No.

63281-9

DILUTION: 1 TO 128 (one ounce per gallon)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

GENERAL USE DIRECTIONS:

USES: For use on hard, non-porous surfaces such as floors, walls, metal surfaces, painted surfaces, exterior bowl surfaces, empty basins, showers, conductive flooring and lavatory fixtures. For use in hospitals, nursing homes, schools, colleges, medical and dental offices, farms, hog farms, equine farms, poultry and turkey farms, federally inspected meat and poultry plants, tack shops, kennels, pet shops, veterinary clinics, and animal life science laboratories. This includes barns, calf pens, laying operations, breeder operations, hatcheries, agricultural buildings and premises, live haul equipment, trucks, foot baths, and meat and poultry processing plants.

APPLICATIONS: Use 1 ounce of BC II PHENOLIC per gallon of water for a minimum contact time of 10 minutes in a single application. For disinfecting, remove gross filth and heavy soil deposits, then thoroughly wet surfaces. BC II PHENOLIC is a extremely versatile and can be applied with a mop, sponge, or cloth, as well as soaking or used as a holding solution. Thoroughly wet surfaces with the recommended use solution for at least 10 minutes. The recommended use solution is used once and discarded on a daily basis. Rinsing is not necessary on floor surfaces unless floors are to be waxed or polished. Tuberculocidal effectiveness is in 10 minutes at 20°C.

HATCHERY: Can be used for cleaning, disinfecting and deodorizing floors, walls and equipment throughout the hatchery. This includes incubators, setters, and racks. Add 1 ounce per gallon of water. Hot or cold water may be used. Remove loose waste, gross filth, and heavy soil deposits prior to disinfecting. Repeat with fresh solution to disinfect.

USDA: For use in federally inspected meat and poultry plants on all surfaces in inedible product processing areas, nonprocessing areas, and/or exterior areas.

***KILLS HIV (AIDS VIRUS) ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for the transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures. **CLEANING PROCEDURE:**

Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of BC II PHENOLIC. This cleaning may be accomplished with any cleaning solution, including this product.

CONTACT TIME: HIV-1 (AIDS virus) is inactivated after five (5) minutes at 25°C. Use at a 10 minute contact time at 20°C for other viruses and bacteria claimed.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 10 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.
03281-9

4/6

DIRECTIONS FOR USE ON FARM PREMISES:

1. Do not use in milking stalls, milking parlors, or milk houses.
2. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
3. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
4. Empty all troughs, racks, and other feeding and watering appliances.
5. Thoroughly clean all surfaces with soap or detergent and rinse with water.
6. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
7. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
8. Ventilate buildings, coops, cars, boats, and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set, or dried.
9. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

(Refer to label for precautionary and practical treatment statements.)

EPA Reg. No. 63281-

[Can be used as an insert or hang tag or place on label.]

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 10 1997

Under the Federal Insecticide
Fungicide and Rodenticide Act as
amended for the pesticide,
registered under EPA Reg. No.

63281-9