

63376-1

06/01/2012

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460-0001

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 1, 2012

Ms. Kim Davis  
Consultant/Agent for  
Family Products Sdn. Berhad  
c/o RegWest Company LLC  
8203 West 20<sup>th</sup> Street, Suite A  
Greeley, CO 80634-4396

**Subject: Amended Labeling; Response to EPA Letter Dated September 8, 2011  
Family Mosquito Coils, EPA Reg. No. 63376-1  
Your Submission Dated March 2, 2012**

Dear Ms. Davis:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the following label revision provision(s):

First Aid section comment(s):

- To facilitate inclusion of required label language from the acute toxicity that was most recently completed in support of the reregistration of the subject product, you must make the following changes:
  - Insert the following text as a new 3<sup>rd</sup> bulleted statement within the "If Inhaled:" subsection which reads "Call a poison control center or doctor for further treatment advice."
  - Insert the following text as a new 3<sup>rd</sup> bulleted statement within the "If on Skin or Clothing:" subsection which reads "Call a poison control center or doctor for further treatment advice."
  - Revise the "HOT LINE NUMBER" subsection to read "Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-558-5252, Monday through Friday, 8:30 a.m. to 4:30 p.m. EST. After 5:00 p.m., call your poison control center."

A copy of your label stamped "Accepted with Comments" is enclosed for your records. Per 40 CFR §156.10(a)(6), submit one (1) copy of your final printed labeling, incorporating the above changes, before you release the product for shipment. As defined in 40 CFR §152.3, "final printed labeling" means the "label or labeling of the product when distributed or sold." Clearly legible reproductions or photo reductions will be accepted for unusual labels. Note that a clean copy of the master label in most cases does not meet the definition of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with section 6(e) of FIFRA. Your release for shipment of the product bearing amended labeling constitutes accepting of these conditions.

A copy of the acute toxicity review that was most recently completed in support of the reregistration of this product is also enclosed for your records. If you have any questions about this label review, please contact Mr. Carmen J. Rodia, Jr. at (703) 306-0327 or via e-mail at [Rodia.Carmen@epa.gov](mailto:Rodia.Carmen@epa.gov).

Sincerely Yours,

*For*

Richard J. Gebken  
Product Manager (10)  
Insecticide Branch  
Registration Division (7504P)

Enclosure:: Label stamped "Accepted"  
Acute toxicity review, dated May 29, 2012  
063376-00001 D424941 & D462015

# Family Mosquito Coils

{Box Front Panel}

For Outdoor Use

[[Select marketing claims from *Marketing Claims* section below]]

**Active Ingredient:**

d/l-allethrolone d-trans chrysanthemate (CAS No. 584-79-2) .....	0.15%
<b>Other Ingredients</b> .....	<b>99.85%</b>
<b>Total</b> .....	<b>100.00%</b>

**Keep Out of Reach of Children**

### CAUTION

See back {or other appropriate descriptive term} panel for First Aid and additional Precautionary Statements.

EPA Reg. No. 63376-1      EPA Est. \_\_\_\_\_

Manufacturing Date: \_\_\_\_\_

Net Contents: \_\_\_\_ oz. (\_\_\_\_ gm.)  
{0.8 to 3.5 oz.}

(\_\_\_\_ Mosquito Coils)  
{2 to 10}  
[One Reusable Pot]

**ACCEPTED  
With COMMENTS  
In EPA Letter Dated:  
June 1, 2012**  
Under the Federal Insecticide, Fungicide  
and Rodenticide Act, As amended, for the  
pesticide Registered under EPA Reg. No:  
**63376-1**

{Box Back Panel}

### First Aid

<b>If Inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Immediately rinse skin with plenty of water for 15-20 minutes.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p>Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also contact {appropriate toll-free 24/7 number} for emergency medical treatment information.]</p>	

*{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}*

### PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

**CAUTION:** Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapors. Thoroughly wash with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**Physical or Chemical Hazards**

Use with caution to avoid fire hazard.

**Environmental Hazards**

This product is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain do not blow or wash pesticide from the treatment area.


**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Family Mosquito Coils**

Use outdoors on patios, porches and campsites. Keeps area free from mosquitoes for up to [2] [4] [6] [8] hours [in still air].


Loosen coils until ends move freely. Grasp each comma shaped end in the center between thumb and forefinger of each hand and separate the two coils by gently pressing them apart. Hold one comma shaped end and lightly shake or gently pry to separate the coils. Insert top of coil stand into slot in comma shaped end of coil. Light the outer end of the coil and let burn (flame) approximately 1/4 inch or less. Blow out the flame and allow the coil to smolder until extinguished. For shorter periods of use break the coil into desired length and place in notched cut out of holder. For best results, do not use in wind. Coil burns for up to [2] [4] [6] [8] hours in still air.



Separate Coil



Light Coil



Place in Stand

{When marketed with ceramic pot:}

Insert the coil into metal stand in pot. Light the outer end of the coil and let burn (flame) approximately 1/4 inch or less. **Blow out the flame**, place the lid on the pot and allow the coil to smolder until extinguished. For shorter periods of use, carefully extinguish coil and re-light when needed. For best results, do not use in wind. Coil burns up to [2] [4] [6] [8] hours in still air.

{Graphic}

Mount Coil

{Graphic}

Light Coil

{Graphic}

Blow out

Flame

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Do not store near sources of heat or open flame. Store in original container in a cool, dry (preferably locked) area inaccessible to children or pets. **Pesticide/Container Disposal: If empty:** Non-refillable container; do not reuse or refill this container. Offer for recycling if available. **If partly filled or for unused coils:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

{Per PR Notice 2007-4 the batch code/lot number will appear on the label or container.}

**Family Products Sdn. Berhad**

Lot 34, Kampong Sentosa Industrial Estate  
31100 Sg. Siput (U), Perak, Malaysia

[Questions? Comments? Call {appropriate number} weekdays, or write {Person's Name}, Family Products Sdn. Berhad, Perak, Malaysia ]

© {year} All Rights Reserved.

{Coil Wrapper Front Panel}

**FAMILY MOSQUITO COILS**

[[Select marketing claims from *Marketing Claims* section below]]

For Outdoor Use

**Active Ingredient:**

d/l-allethrolone d-trans chrysanthemate (CAS No. 584-79-2) .....	0.15%
<b>Other Ingredients</b> .....	<b>99.85%</b>
<b>Total</b> .....	<b>100.00%</b>

**Keep Out of Reach of Children**

**CAUTION**

See back {or other appropriate descriptive term}  
panel for First Aid and additional Precautionary Statements.

Net Contents: \_\_\_\_ oz. (\_\_\_\_ gm.)  
{0.8 to 3.5 oz.}

(\_\_\_\_ Mosquito Coils)  
{2 to 10}

EPA Reg. No. 63376-1      EPA Est. \_\_\_\_\_

**Family Products Sdn. Berhad**

Lot 34, Kampong Sentosa Industrial Estate  
31100 Sg. Siput (U), Perak, Malaysia

{Pot Label}

## Family Mosquito Coils

For Best Results Use Only Family Mosquito Coil Refills

Only for outdoor use. Place on a flat heat-resistant surface away from anything flammable and out of reach of children and pets. Keep pot free of matches or other foreign matter. Do not use smoldering coil to light other items. To extinguish coil, smother lit coil by carefully lifting coil out of pot and pressing the glowing end against the base of the pot. Do not leave a burning coil unattended. Coils other than Family Mosquito Coils will not work in this device. Family Products Sdn. Berhad will not be held responsible for damage, injury or poor performance caused by use of other mosquito coil refills.

**Important Note:** To reduce fire hazard and to avoid overheating of the pot surface, blow out flame after lighting coil!

[Questions? Comments? Call {appropriate number} weekdays, or write {Person's Name}, Family Products Sdn. Berhad, Perak, Malaysia ]

© {year} All Rights Reserved.

{Marketing Claims}

[Refills]

Use on Patios, Campsites, Cabanas and Other Outdoor Areas  
To Keep the Area Free of Mosquitoes for Hours

Each Coil Burns Two to Four [Six] [Eight] Hours

For Best Results Use Only Family Mosquito Coil Refills

{End of Marketing Claims}

[ ] Denotes alternate/optional language

{ } Denotes language that does not appear on the market labeling

## **PLEASE NOTE**

**This image contains more than one label  
approved for this product on this date.**

**PRIA DUE DATE: 8/2/12****# 100-RUUE**

Previously 100-OEEN

**Note to: File**

**Re:** New Product Registration, R310  
Product Name: Lumax EZ Herbicide  
EPA Registration Number: 100-1442  
Submission Date: January 10, 2012  
Decision Number: 459863

**New Product Unconditional Registration**

- New Restricted Use (S-metolachlor, Atrazine, Mesotrione) product for use on corn and sorghum. Citing 100-1356 as a similar product.

**Chemistry Review**

- Not substantially similar due to a difference in the nominal concentration for one of the actives.
- Group A & B data is acceptable.
- Basic CSF dated November 29, 2011 is acceptable.
- Storage Stability and Corrosion Characteristics data is required within 15 months as a term of registration.
- Physical and Chemical Hazards Statement – Shyman Mathur/Chemistry Team Leader confirmed that the “Do not use or store near heat or open flame.” is not required for this product since the flashpoint is above 150 degrees.

**TRB Acute Toxicity Review**

- Substantially similar to the cited product; data may be bridged.
- Precautionary Statements – the Notice of Registration for this product follows TRB’s recommendations for the Precautionary Statements, including protective eyewear. Other PPE requirements are addressed later in this document.
- Protective Eyewear – The Metolachlor RED and Label Review Manual do not require protective eyewear for this (Primary eye irritation - Tox III) product. The Acute Tox review recommends that protective eyewear be included in the Precautionary Statements, so it is included in the Notice of Registration.
- First Aid box requirements are less than on the proposed label. The registrant may remove the optional statements.
- Skin Sensitizer Statement – The Metolachlor Reregistration Eligibility Document (RED) requires a dermal sensitizer statement. The TRB Acute Tox review determined that the product is negative for dermal sensitization. John Redden/Acting Technical Review Branch Chief stated that a RED is a snapshot in time, and confirmed that a skin sensitizer statement is not needed for this product.

**Personal Protective Equipment (PPE)**

- The PPE for this product is based on the Metolachlor Reregistration Eligibility Document (RED) requirements, not on the Acute Toxicity Review. Please note that the RED PPE requirements are greater than those outlined in the Acute Toxicity Review.

**SEE NEXT PAGE**

- The new product requirements conflict with the Notice of Registration for the cited product where the registrant is instructed to reduce the RED required PPE to long-sleeved shirts and long pants, chemical resistant gloves – Category C, and shoes plus socks per the Acute Toxicity Review for that product.
- Background - When contacted regarding this action, Syngenta/Jerry Wells explained that Syngenta submitted a revised S-metolachlor product label reducing the PPE requirement believing that the cancer classification for metolachlor/S-metolachlor had changed. He also mentioned that he believed the RED initially required the extensive PPE listed on the proposed label because of potential cancer risk. The Syngenta label with reduced PPE was approved. Through SLITS, the State of Maryland contacted RD inquiring as to why the requirements for that S-metolachlor product were less than those outlined in the RED. RD later determined that any changes in PPE would require HED review before registrants could revise any product labels. Knowing that HED review is required before the PPE can be reduced, Syngenta incorporated all of the RED PPE requirements on the proposed label, which are acceptable as presented. Additionally, Kathryn Montague/PM23 and John Redden/Acting TRB Chief agreed that the RED PPE requirements, which are more protective than those recommended by the TRB Acute Toxicity Review, are appropriate for this action.

**Use Rates**

The percentage of active ingredients in this product is slightly less than the percentage contained in the cited product. Use rates were adjusted upward to compensate for the reduced percentage of active ingredients. The use rates are comparable for each of the products. There are differences in the use rates, although they are negligible. For example, the Atrazine use rates for this triple active product are as follows:


Cited Product Maximum Annual Per Acre	3.00 qts. (.75 lb. atrazine)
This Product - Maximum Annual Per Acre	3.25 qts. (.7596 lb. atrazine)
Cited Product Max Rate for Corn	3.00 qts. (.75 lb. atrazine)
This Product - Max Rate for Corn	3.25 qts. (.7596 lb. atrazine)
Cited Product Max Rate for Grain Sorghum	2.50 qts. (.625 lb. atrazine)
This Product – Max Rate for Grain Sorghum	2.70 qts. (.6311 lb. atrazine)

**Safener & Active Ingredient Tolerances**

- Dan Kenny/Herbicide Branch Chief confirmed that the incorporation of Benoxacor as a safener in this product, which contains atrazine and mesotrione in addition to S-metolachlor as active ingredients, is acceptable.
- Corn and sorghum tolerances are established for each of the active ingredients in this product.
- See the next two pages for additional information regarding tolerances.



Page 3 of 4  
New Product Registration, R310  
Product Name: Lumax EZ Herbicide  
EPA Registration Number: 100-1442  
Submission Date: January 10, 2012  
Decision Number: 459863



**e-CFR Data is current as of July 27, 2012**

**Title 40: Protection of Environment**

**PART 180—TOLERANCES AND EXEMPTIONS FOR PESTICIDE CHEMICAL RESIDUES IN FOOD**

**Subpart C—Specific Tolerances**

**§ 180.460 Benoxacor; tolerances for residues.**

(a) *General*. Tolerances are established for residues of the inert ingredient (safener) benoxacor (4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1, 4-benzoxazine) at 0.01 parts per million (ppm) when used in pesticide formulations containing metolachlor or S-metolachlor in or on raw agricultural commodities for which tolerances have been established for metolachlor or S-metolachlor.

(b) Section 18 emergency exemptions. [Reserved]

(c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. [Reserved]

[63 FR 7305, Feb. 13, 1998, as amended at 70 FR 21631, Apr. 27, 2005]

**§ 180.368 S-metolachlor; tolerances for residues.**

(2) Tolerances are established for residues of S-metolachlor, including its metabolites and degradates, in or on the commodity(s), as defined. Compliance with the tolerance levels specified in the following table below is to be determined by measuring only the sum of free and bound S-metolachlor, S-2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl)acetamide, its R-enantiomer, and its metabolites, determined as the derivatives, 2-(2-ethyl-6-methylphenyl)amino-1-propanol and 4-(2-ethyl-6-methylphenyl)-2-hydroxy-5-methyl-3-morpholinone, calculated as the stoichiometric equivalent of S-metolachlor, in or on the commodity.

Commodity	Parts per million
Corn, field, grain	0.10
Corn, field, forage	6.0
Corn, field, stover	6.0
Corn, pop, grain	0.10
Corn, pop, stover	6.0
Corn, sweet, forage	6.0
Corn, sweet, kernel plus cob with husks removed	0.10
Corn, sweet, stover	6.0
Sorghum, grain, forage	1.0
Sorghum, grain, grain	0.3
Sorghum, grain, stover	4.0
Sorghum, sweet, stalk	4.0

**SEE NEXT PAGE**

4/4

**§ 180.220 Atrazine; tolerances for residues.** (a) *General*. Tolerances are established for the combined residues of the herbicide atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine) and its chlorinated metabolites 2-amino-4-chloro-6-isopropylamino-s-triazine, 2-amino-4-chloro-6-ethylamino-s-triazine, and 2,4-diamino-6-chloro-s-triazine, in or on the following food commodities:

Commodity	Parts per million
Corn, field, forage	1.5
Corn, field, grain	0.20
Corn, field, stover	0.5
Corn, pop, forage	1.5
Corn, pop, grain	0.20
Corn, pop, stover	0.5
Corn, sweet, forage	15
Corn, sweet, kernel plus cob with husks removed	0.20
Corn, sweet, stover	2.0
Sorghum, forage, forage	0.25
Sorghum, grain, forage	0.25
Sorghum, grain, grain	0.20
Sorghum, grain, stover	0.50

**§ 180.571 Mesotrione; tolerances for residues.** (a) *General*. Tolerances are established for residues of the herbicide mesotrione, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only mesotrione, 2-[4-(methylsulfonyl)-2-nitrobenzoyl]-1,3-cyclohexanedione, in or on the following raw agricultural commodities:

Commodity	Parts per million
Corn, field, forage	0.01
Corn, field, grain	0.01
Corn, field, stover	0.01
Corn, pop, grain	0.01
Corn, pop, stover	0.01
Corn, sweet, forage	0.5
Corn, sweet, kernel plus cob with husks removed	0.01
Corn, sweet, stover	1.5
Sorghum, grain, forage	0.01
Sorghum, grain, grain	0.01
Sorghum, grain, stover	0.01
Sorghum, sweet	0.01

**NEXT**

**LABEL**

100-1442

8/2/2012

1/40



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

A Reg. Number: 100-1442  
Date of Issuance: AUG - 2 2012

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:  
Unconditional  
Name of Pesticide Product:  
Lumax-EZ Herbicide

Name and Address of Registrant (include ZIP Code):  
Syngenta Crop Protection LLC  
PO Box 18300  
Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5) provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data. If required, failure to submit acceptable data to fulfill these requirements may result in registration cancellation in accordance with FIFRA section 6(e).
- 2) Replace the phrase "EPA Reg. No. 100-XXXX" with "EPA Reg. No. 100-1442" and assure that the EPA Establishment Number and Net Contents are also on the label.
- 3) Generate one-year storage stability (830.6317) and corrosion characteristics (830.6320) data on the product. The observations should be made at 0, 3, 6, 9, and 12 month intervals. The results must be submitted to the Agency in electronic and hard copy format within 15 months of the date on this notice.
- 4) Per the Acute Toxicity Review, revise the Precautionary Statement appearing under the signal word to read as follows: "Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear [protective eyewear.]\*" Please note that the eyewear may be specified.
- 5) Per the Acute Toxicity Review, the "If on skin or clothing", "If swallowed", and "If inhaled" statements appearing in the First Aid box are not required. These optional statements may remain on the label as additional safety information for product users or they may be removed.

SEE NEXT PAGE FOR ADDITIONAL COMMENTS

Signature of Approving Official:  
Kathryn V. Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date:  
AUG - 2 2012

- 6) NOTE Per the chemistry review, the Basic Confidential Statements of Formula dated November 29, 2011 is acceptable
- 7) Per the Chemistry Review, the Physical and Chemical Hazards statement appearing on page 5 is not required You may retain the statement or you may remove it from the label
- 8) The "USE PRECAUTIONS" header appearing in the middle of page 8 must be changed to read "USE RESTRICTIONS AND PRECAUTIONS"
- 9) The term "restrictions" must be added to the bullet near the middle of page 9 that reads "Observe all precautions and limitations on the label of each product used in tank mixtures " for it to read "Observe all restrictions, precautions, and limitations on the label of each product used in tank mixtures "
- 10) In the ROTATIONAL CROPS section near the top of page 16, the last bullet that reads "Do not rotate to food or feed crops other than those listed on this label " conflicts with the previous bullets and must be removed from the label
- 11) Aerial application is prohibited for this product The reference to aircraft smoke generators in the paragraph at the top of page 19 is confusing and must be removed from the label The sentence must be changed from "Their presence can be indicated by ground fog, however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator " to read "Their presence can be indicated by ground fog, however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source "
- 12) In the CROP USE DIRECTIONS under the CORN header on page 22, the preemergence surface application restriction reads "Do not exceed 3 0 qts of Lumax EZ per season " This rate conflicts with the 3 25 qts per acre application instructions in the "Lumax EZ Use Rate" portion of this page Revise the restriction to read "Do not exceed 3 25 qts of Lumax EZ per season "
- 13) In the GRAIN SORGHUM use directions on page 27 the restriction reads "Do not apply more than 2 5 quarts of Lumax EZ per growing season " This rate conflicts with the 2 7 qts per acre application rate referenced on page 26 in the grain sorghum section Revise the restriction to read "Do not apply more than 2 7 quarts of Lumax EZ per growing season "
- 14) All Precautionary Statements on the base label and all booklets must be consistent Modify the statements on pages 31 and 32 as specified above Additionally, the First Aid text appearing on page 31 must be contained in a box
- 15) NOTE While no additional data is being requested at this time, any marketing claims made on the pesticide label at any time must be substantiated by data maintained in your files If data supporting marketing claims made on the product label is not available then those claims must be removed

**SEE NEXT PAGE**

Page 3 of 3  
EPA Registration # 100 1442  
Product Lumax EZ Herbicide  
Decision Number 459863

16) NOTE Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

17) Submit one (1) copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

4/40

[Booklet]

**RESTRICTED USE PESTICIDE  
(GROUND AND SURFACE WATER CONCERNS)**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER

**For Ground Application Only**

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited

**GROUP 5 15 27 HERBICIDES**

**Lumax® EZ**

**Herbicide**

A Preemergence Herbicide for Control of Annual Grass and Broadleaf Weeds in Field Corn, Field Production Seed Corn, Field Silage Corn, Sweet Corn, Yellow Popcorn, and Grain Sorghum

**Active Ingredients\***

S-Metolachlor (CAS No 87392-12-9)	27.10%
Atrazine** (CAS No 1912-24-9)	10.20%
Mesotrione (CAS No 104206-82-8)	2.71%
Other Ingredients	59.99%
<b>Total</b>	<b>100.00%</b>

\*Active ingredients per gallon: S-metolachlor 2.49 pounds, Mesotrione 0.249 pounds, and Atrazine 0.935 pounds

\*\*Atrazine with a maximum of 0.27% related triazines

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See additional precautionary statements and directions for use inside booklet

EPA Reg No 100-XXXX  
EPA Est

 **Callisto Plant Technology®**

2.5 gallons  
Net Contents

120 gallons  
Net Contents

220 gallons  
Net Contents

330 gallons  
Net Contents

\_\_\_\_\_ gallons [bulk]  
Net Contents

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

**AUG - 2 2012**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg No

100-1442

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye</li> <li>• Call a Poison Control Center or doctor for treatment advice</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a Poison Control Center or doctor for treatment advice</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a Poison Control Center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth if possible</li> <li>• Call a Poison Control Center or doctor for further treatment advice</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment	
<b>HOT LINE NUMBER</b>	
For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call <b>1-800-888-8372</b>	

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION**

Cases moderate eye irritation Avoid contact with eyes skin or clothing Wear protective eyewear Harmful if swallowed Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below If you want more options, follow the instructions for Category A on an EPA Chemical Resistance Category Selection Chart



**Mixers, Loaders, Applicators and other handlers must wear**

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves - Category A (e.g. barrier laminate butyl rubber  $\geq 14$  mils nitrile rubber  $\geq 14$  mils neoprene rubber  $\geq 14$  mils natural rubber  $\geq 14$  mils polyethylene, polyvinyl chloride (PVC)  $\geq 14$  mils or Viton®  $\geq 14$  mils)
- Chemical-resistant footwear plus socks
- Chemical-resistant apron when mixing/loading, cleaning up spills or cleaning equipment or otherwise exposed to the concentrate
- Chemical-resistant headgear for overhead exposure

See engineering controls for additional requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Aerial application is prohibited

**Engineering Control Statements**

When applicators use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)] the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

**Users should**

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. This pesticide contains atrazine, which has been shown to be toxic to aquatic invertebrates. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from treated areas.

**Ground Water Advisory**

Lumax EZ contains the active ingredients S-metolachlor atrazine and mesotrione

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water Atrazine has been found in ground water Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable i.e., well-drained Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water

S-metolachlor has the potential to leach through soil into ground water under certain conditions as a result of agricultural use Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination

**Surface Water Advisory**

The active ingredients in this product have the potential to contaminate surface water through ground spray drift Under some conditions the active ingredients may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application These include poorly drained or wet soils with readily visible slopes toward adjacent surface waters frequently flooded areas areas overlaying extremely shallow ground water areas with in-field canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water

A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams and springs will reduce the potential for contamination of water from runoff Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours Sound erosion control practices will reduce this product's contribution to surface water contamination

**Mixing/Loading Instructions**

Care must be taken when using this product to prevent back-siphoning into wells spills or improper disposal of excess pesticide, spray mixtures or rinsates

Check valves or antisiphoning devices must be used on mixing equipment

This product must not be mixed/loaded or used within 50 ft of wells including abandoned wells drainage wells, and sink holes Operations that involve mixing, loading, rinsing, or washing to this product into or from pesticide handling or application equipment or containers within 50 ft of any well are prohibited unless conducted on an

impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain, at a minimum, 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

Additional State imposed requirements regarding well head setbacks and operational area containment must be observed.

This product must not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be applied within 66 ft. of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 ft. from the edge of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 ft. buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

**Tile-Outletted Terraced Fields Containing Standpipes**

One of the following restrictions must be used in applying atrazine to tile-outletted terraced fields containing standpipes:

1. Do not apply this product within 66 ft. of standpipes in tile-outletted terraced fields.
2. Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2-3 inches in the entire field.
3. Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

**Physical and Chemical Hazards**

Do not use or store near heat or open flame.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA; and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

10/40

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through [www.atrazine-watershed.info](http://www.atrazine-watershed.info) or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Syngenta Crop Protection LLC for a refund.

Lumax EZ must be used only in accordance with requirements on this label or in separately published Syngenta supplemental labeling requirements for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil and water, wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves – Category A (e.g. barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils or Viton ≥14 mils)
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES**

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited

---

**PRODUCT INFORMATION**

---

Lumax EZ may be used preemergence and postemergence in the culture of field corn, field seed corn and field corn silage. Lumax EZ may also be used in the culture of sweet corn, yellow popcorn and grain sorghum but the application must be made prior to crop emergence, (i.e., preemergence) or severe crop injury may occur.

Lumax EZ is a combination of the herbicides S-metolachlor, mesotrione, and atrazine plus the safener benoxacor. Lumax EZ controls weeds by interfering with normal germination and seedling development. It is recommended for management of the weed species listed in Tables 1 and 2.

**USE PRECAUTIONS**

Read all label directions before using.

**Atrazine Herbicide Rate Limitations**

Certain states may have established rate limitations within specific geographical areas for the use of atrazine. These more restrictive/protective requirements must be followed. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

- When tank mixing or sequentially applying atrazine or products containing atrazine with Lumax EZ to corn, do not exceed an application rate of 2.0 pounds active ingredient of atrazine per acre for any single application and the total pounds of atrazine applied (lb a.i. per acre) must not exceed 2.5 pounds active ingredient per acre per year.

**Note:** For purposes of calculating total atrazine active ingredient applied, Lumax EZ contains 0.935 lb a.i. atrazine plus relateds per gallon.

- **Grazing Restriction:** To avoid possible illegal residues, do not graze or feed forage from treated areas for 45 days following application.
- **Preharvest Interval (PHI):** Field corn may be treated up to 12 inches tall. Do not harvest forage, grain, or stover within 60 days after application. Do not harvest sweet corn forage within 45 days after application.

- Do not apply this product through any type of irrigation system
- Do not apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas
- To prevent off-site movement due to runoff or wind erosion
  - Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation
  - Do not apply to impervious substrates, such as paved or highly compacted surfaces
  - Do not use tail water from the first flood or furrow irrigation of treated fields to treat nontarget crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation
- Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial weed control
- Dry weather following preemergence application of Lumax EZ or a Lumax EZ tank mixture may reduce effectiveness. Cultivate if weeds develop in conventional tillage corn
- Observe all precautions and limitations on the label of each product used in tank mixtures
- Thoroughly clean sprayer or other application device before using. See **Storage and Disposal** section for disposal instructions. Do not use a sprayer or applicator contaminated with other materials or crop damage or sprayer clogging of the application device may occur
- This product will not provide consistent control of emerged grass weeds
- Do not apply more than 3.25 qts of Lumax EZ per acre per growing season
- Applying Lumax EZ postemergence (emerged corn) to corn that has received an at-plant application of Counter® insecticide can result in severe corn injury. Temporary corn injury may occur if Lumax EZ is applied to emerged corn where organophosphate insecticides other than Counter were applied at planting
- Do not make postemergence (emerged corn) applications of Lumax EZ in a tank mix with any organophosphate or carbamate insecticide, or severe corn injury may occur
- Postemergence (emerged corn) applications of any organophosphate or carbamate insecticide within 7 days before or 7 days after a Lumax EZ application may result in severe corn injury

- Do not use Lumax EZ on any crop other than field corn (for grain, seed, or silage) sweet corn (preemergence applications only), yellow popcorn (preemergence applications only), or grain sorghum (preemergence applications only)
- Do not use Lumax EZ in the culture of white popcorn or ornamental (Indian) corn or injury may occur
- Do not contaminate irrigation water used for crops other than field corn Do not contaminate water used for domestic purposes
- Avoid drift onto adjacent crops
- Avoid spray overlap, as crop injury may result
- Do not allow Lumax EZ to contaminate feed or food
- Do not store Lumax EZ near seeds, fertilizers or foodstuffs
- All containers of Lumax EZ must be kept tightly closed when not in use
- Do not use aerial application to apply Lumax EZ
- Do not apply Lumax EZ to sweet corn or yellow popcorn after the crop has emerged or severe crop injury may occur

Applied according to directions and under normal growing conditions, Lumax EZ will not harm the treated crop During germination and early stages of growth, extended periods of unusually cold and wet or hot and dry weather insect or plant disease attack carryover pesticide residues, the use of certain soil applied systemic insecticides, improperly placed fertilizers or soil insecticides may weaken crop seedlings Lumax EZ used under these conditions could result in crop injury

**RESISTANCE MANAGEMENT**

Lumax EZ is a combination of atrazine S-metolachlor and mesotrione (Group 5, 15 and 27 Herbicides)

Naturally occurring biotypes of certain broadleaf weed species with resistance to triazines ALS PPO Glycine (glyphosate) and HPPD herbicides are known to exist If biotypes of weeds resistant to triazines, ALS, PPO and glycine inhibitors are present in the field, this herbicide should control them if they are listed in Tables 1 and 2

To reduce the risk of weeds developing resistance to HPPD inhibitors, implement a program including both preemergence and postemergence herbicides that provide effective control of all weeds using multiple modes of action Consider weed resistance management strategies that include two or more modes of action where a minimum of two modes of action are effective at controlling the target weed when either are applied alone Scout fields and eliminate weed escapes Read and follow all label recommendations



Lumax EZ contains three herbicide active ingredients and three modes of action and can be an effective component of a weed resistance management strategy.

### **INTEGRATED PEST (WEED) MANAGEMENT**

Lumax EZ may be integrated into an overall pest management strategy. Practices known to reduce weed development (tillage and crop competition) and herbicide use (weed scouting, proper application timing, banding, and rotations) should be followed wherever possible. Consult local agricultural and weed authorities for additional Integrated Pest Management strategies established for your area.

### **SOIL ORGANIC MATTER**

The organic matter of the soil on which the application is to be made must be known or determined prior to application. The use rate of Lumax EZ is based on percent soil organic matter.

### **REDUCED AND NO-TILL SYSTEMS**

Lumax EZ may be used in reduced and no-till systems. The highest levels of control will be obtained when applications are made as close to planting as possible. It is recommended that a burndown herbicide such as Gramoxone Inteon®, Touchdown® brands, Roundup® brands, or 2,4-D be tank mixed with Lumax EZ in reduced or no-till systems if weeds are present at application and the corn has not yet emerged.

### **WEEDS CONTROLLED**

Lumax EZ applied as directed in this label will control or suppress the weeds listed in Tables 1 and 2. Additional weeds may be controlled with tank mixes. See the **Lumax EZ Tank Mix Combinations** section for recommended tank-mix combinations. Always consult the tank-mix product labels for specific rates and use directions.

Table 1 Weeds Controlled or Partially Controlled by Preemergence Applications of Lumax EZ

Common Name	Weed Type <sup>1</sup>	Scientific Name	C = Control PC = Partial Control
Amaranth Palmer	B	<i>Amaranthus palmeri</i>	C
Amaranth Powell	B	<i>Amaranthus powellii</i>	C
Barnyardgrass	G	<i>Echinochloa crus-galli</i>	C
Bedstraw catchweed	B	<i>Galium aparine</i>	PC
Beggarweed Florida	B	<i>Desmodium tortuosum</i>	C
Buckwheat wild	B	<i>Polygonum convolvulus</i>	C
Buffalobur	B	<i>Solanum rostratum</i>	C
Carpetweed	B	<i>Mollugo verticillata</i>	C
Chickweed common	B	<i>Stellaria media</i>	C
Cocklebur common	B	<i>Xanthium strumarium</i>	PC
Crabgrass	G	<i>Digitaria spp</i>	C
Crowfootgrass	G	<i>Dactyloctenium aegyptium</i>	C
Cupgrass prairie	G	<i>Enochloa contracta</i>	C
Cupgrass Southwestern	G	<i>Enochloa gracilis</i>	C
Cupgrass woolly	G	<i>Enochloa villosa</i>	PC
Deadnettle purple	B	<i>Lamium purpureum</i>	C
Devil s-claw	B	<i>Proboscidea louisianica</i>	C
Foxtail giant	G	<i>Setaria faberi</i>	C
Foxtail green	G	<i>Setaria viridis</i>	C
Foxtail robust (purple white)	G	<i>Setaria spp</i>	C
Foxtail yellow	G	<i>Setaria pumila</i>	C
Galinsoga	B	<i>Galinsoga parviflora</i>	C
Goosegrass	G	<i>Eleusine indica</i>	C
Henbit	B	<i>Lamium amplexicaule</i>	C
Horseweed (marestail)	B	<i>Conyza canadensis</i>	C
Jimsonweed	B	<i>Datura stramonium</i>	C
Johnsongrass seedling	G	<i>Sorghum halepense</i>	PC
Kochia	B	<i>Kochia scoparia</i>	C
Lambsquarters common	B	<i>Chenopodium album</i>	C
Mallow Venice	B	<i>Hibiscus trionum</i>	C
Millet foxtail	G	<i>Setaria italica</i>	C
Millet wild proso	G	<i>Panicum miliaceum</i>	PC
Morningglory ivyleaf/entireleaf	B	<i>Ipomoea hederacea</i>	PC
Mustard wild	B	<i>Brassica kaber</i>	C
Nightshade black	B	<i>Solanum nigrum</i>	C
Nightshade eastern black	B	<i>Solanum ptycanthum</i>	C
Nightshade hairy	B	<i>Solanum sarrachoides</i>	C
Panicum Texas	G	<i>Panicum texanum</i>	PC
Pigweed redroot	B	<i>Amaranthus retroflexus</i>	C
Pigweed smooth	B	<i>Amaranthus hybridus</i>	C
Puncturevine	B	<i>Tribulus terrestris</i>	PC

Common Name	Weed Type <sup>1</sup>	Scientific Name	C = Control PC = Partial Control
Purslane common	B	<i>Portulaca oleracea</i>	C
Pusley Florida	B	<i>Richardia scabra</i>	C
Radish wild	B	<i>Raphanus raphanistrum</i>	C
Ragweed common	B	<i>Ambrosia artemisiifolia</i>	PC
Ragweed giant	B	<i>Ambrosia trifida</i>	PC
Rice red	G	<i>Oryza sativa</i>	C
Sandbur field	G	<i>Cenchrus incertus</i>	PC
Sesbania hemp	B	<i>Sesbania exaltata</i>	C
Shattercane	G	<i>Sorghum bicolor</i>	PC
Shepherd s-purse	B	<i>Capsella bursa-pastors</i>	C
Sida prickly	B	<i>Sida spinosa</i>	PC
Signalgrass broadleaf	G	<i>Brachiana platyphylla</i>	PC
Signalgrass narrowleaf	G	<i>Brachiana piligera</i>	C
Smartweed ladysthumb	B	<i>Polygonum persicaria</i>	C
Smartweed Pennsylvania	B	<i>Polygonum pennsylvanicum</i>	C
Sprangletop red	G	<i>Leptochloa filiformis</i>	C
Starbur bristly	G	<i>Acanthospermum hispidum</i>	C
Sunflower common	B	<i>Helianthus annus</i>	PC
Velvetleaf	B	<i>Abutilon theophrasti</i>	C
Waterhemp common	B	<i>Amaranthus rudis</i>	C
Waterhemp tall	B	<i>Amaranthus tuberculatus</i>	C
Witchgrass	G	<i>Panicum capillare</i>	C

<sup>1</sup>B = Broadleaf G = Grass S = Sedge

Thoroughly till soil or make an application of a burndown herbicide to destroy germinating and emerged weeds. Plant crop into moist soil immediately after tillage.

If a significant rainfall does not occur within 7 days after application, weed control may be decreased. If irrigation is available, apply 1/2 to 1 inch of water. If irrigation is not available, a uniform shallow cultivation is recommended as soon as weeds emerge.

**Table 2. Weeds Controlled or Partially Controlled by Early Postemergence Applications of Lumax EZ**

Common Name	Weed Type <sup>1</sup>	Scientific Name	C = Control PC = Partial Control
Amaranth, Palmer	B	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	B	<i>Amaranthus powellii</i>	C
Bedstraw, catchweed	B	<i>Galium aparine</i>	PC
Beggarweed, Florida	B	<i>Desmodium tortuosum</i>	C
Buckwheat, wild	B	<i>Polygonum convolvulus</i>	C
Buffalobur	B	<i>Solanum rostratum</i>	C
Carpetweed	B	<i>Mollugo verticillata</i>	C
Chickweed, common	B	<i>Stellaria media</i>	C
Cocklebur, common	B	<i>Xanthium strumarium</i>	C
Crabgrass, large	G	<i>Digitaria sanguinalis</i>	C <sup>2</sup>
Dandelion	B	<i>Taraxacum officinale Weber</i>	PC
Deadnettle, purple	B	<i>Lamium purpureum</i>	C
Devil's-claw	B	<i>Proboscidea louisianica</i>	C
Galinsoga	B	<i>Galinsoga parviflora</i>	C
Hemp	B	<i>Cannabis sativa</i>	C
Henbit	B	<i>Lamium amplexicaule</i>	C
Horsenettle	B	<i>Solanum carolinense</i>	C
Horseweed (marestail)	B	<i>Conyza canadensis</i>	C
Jimsonweed	B	<i>Datura stramonium</i>	C
Kochia	B	<i>Kochia scoparia</i>	C
Lambsquarters, common	B	<i>Chenopodium album</i>	C
Mallow, Venice	B	<i>Hibiscus trionum</i>	C
Marestail	B	<i>Hippuris vulgaris</i>	C
Morningglory, ivyleaf/entireleaf	B	<i>Ipomoea hederacea</i>	C
Mustard, wild	B	<i>Brassica kaber</i>	C
Nightshade, black	B	<i>Solanum nigrum</i>	C
Nightshade, eastern black	B	<i>Solanum ptycanthum</i>	C
Nightshade, hairy	B	<i>Solanum sarachoides</i>	C
Nutsedge, yellow	S	<i>Cyperus esculentus</i>	PC
Pigweed, redroot	B	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	B	<i>Amaranthus hybridus</i>	C
Pokeweed	B	<i>Phytolacca americana</i>	C
Potatoes, volunteer	B	<i>Solanum spp.</i>	C
Purslane, common	B	<i>Portulaca oleracea</i>	C
Pusley, Florida	B	<i>Richardia scabra</i>	C
Radish, wild	B	<i>Raphanus raphanistrum</i>	C
Ragweed, common	B	<i>Ambrosia artemisiifolia</i>	C
Ragweed, giant	B	<i>Ambrosia trifida</i>	C
Sesbania, hemp	B	<i>Sesbania exaltata</i>	C
Shepherd's-purse	B	<i>Capsella bursa-pastoris</i>	C
Sida, prickly	B	<i>Sida spinosa</i>	C

Common Name	Weed Type <sup>1</sup>	Scientific Name	C = Control PC = Partial Control
Signalgrass broadleaf	G	<i>Brachiana platyphylla</i>	C <sup>2</sup>
Smartweed ladysthumb	B	<i>Polygonum persicaria</i>	C
Smartweed Pennsylvania	B	<i>Polygonum pennsylvanicum</i>	C
Sunflower common	B	<i>Helianthus annus</i>	C
Thistle Canada	B	<i>Cirsium arvense</i>	C
Velvetleaf	B	<i>Abutilon theophrasti</i>	C
Waterhemp common	B	<i>Amaranthus rudis</i>	C
Waterhemp tall	B	<i>Amaranthus tuberculatus</i>	C

<sup>1</sup>B = Broadleaf G = Grass S = Sedge

<sup>2</sup>Apply before the weed exceeds 2 inches in height

Lumax EZ will not provide consistent control of emerged grass weeds

A tank mix of AAtrex® with Lumax EZ can provide control of certain emerged annual grass weeds Refer to the AAtrex label for weeds controlled and other restrictions

### ROTATIONAL CROPS

When rotating crops following an application of Lumax EZ

- Field corn field seed corn field silage corn sweet corn, yellow popcorn and grain sorghum (Concep® treated seed) may be replanted immediately if crop is lost Do not reapply Lumax EZ
- Winter wheat, barley or rye may be planted 4½ months following application
- If Lumax EZ is applied after June 1, rotating to crops other than corn (all types) or grain sorghum the next spring may result in crop injury
- The following rotational interval applies only to areas west of Highway 83 in the state of Nebraska If Lumax EZ was applied to ground that was under center pivot irrigation and the soil pH is greater than 6.5 dry beans can be planted 10 months following application
- Do not rotate to crops other than corn (all types) cotton, small grain cereals soybeans sorghum or peanuts the spring following application of Lumax EZ
- Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer if additional atrazine or atrazine-containing products are used
- In eastern parts of the Dakotas, KS, western MN, and NE, do not rotate to soybeans for 18 months following application if the combined atrazine rate applied was more than 2.0 lbs a /A, or equivalent band application rate, or soybean injury may occur
- In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required use only when corn (all types) or sorghum is to

follow field corn, or a crop of untreated corn (all types) or sorghum is to precede other rotational crops.

- For all other crops, wait 18 months.
- Do not rotate to food or feed crops other than those listed on this label.

**APPLICATION PROCEDURES**

**ADJUVANTS**

When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

Where Lumax EZ is applied after the field corn has emerged, a non-ionic surfactant at 0.25% v/v (1 qt./100 gals.) may be used. A crop oil concentrate (COC) may also be used at a rate not to exceed 1% v/v (1 gallon/100 gallons) or not more than the equivalent of 1 qt./A. The use of crop oil concentrate (COC) may result in temporary crop injury.

Do not use nitrogen based adjuvants (AMS or UAN) or methylated seed oil (MSO) with Lumax EZ when applied alone to emerged field corn, or when Lumax EZ is applied as a postemergence tank mixture with other products, unless directed for a specific tank mix on this label or as part of a supplemental Lumax EZ label.

Any of these adjuvants may be used at a preemergence or preplant timing, i.e. where the corn crop has not yet emerged to increase burndown activity on existing weeds. Do not apply Lumax EZ to emerged sweet corn, yellow popcorn or grain sorghum or severe crop injury may occur.

For Lumax EZ tank mixtures with Liberty® Herbicide or Ignite® applied to emerged field corn (LibertyLink® hybrids only), AMS may be added as directed on the Liberty or Ignite label. However, AMS must be the only adjuvant added to this tank mixture, or severe crop injury may occur.

**Sprinkler Irrigation:** Do not apply Lumax EZ by sprinkler irrigation. Use a sprinkler system only to incorporate Lumax EZ after application. After Lumax EZ has been applied, a sprinkler irrigation system set to deliver ½-1 inch of water may be used to incorporate the product. Using more than 1 inch of water could result in reduced performance. On sandy soil low in organic matter, use no more than ½ inch of water. Do not use flood irrigation to apply or incorporate Lumax EZ.

**CULTIVATION**

Should weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. If Lumax EZ was incorporated, cultivate less than half the depth of incorporation.

If cultivation is necessary due to soil crusting, compaction, or escaped weeds, adjust equipment to run shallow and minimize soil movement. This will decrease the possibility of diluting or moving the herbicide from the weed control zone.

**SPRAY EQUIPMENT**

**Ground Application**

Spray nozzles should be uniformly spaced, the same size and type, and should provide accurate and uniform application. Use spray nozzles that provide medium to coarse droplet size to avoid drift yet provide good coverage. Ensure that all in-line strainer and nozzle screens in the sprayer are 50-mesh or coarser. Use a pump that can maintain a pressure of at least 35-40 psi at the nozzles and provide proper agitation within the tank to keep the product dispersed. Lower pressures may be used with extended range or drift reduction nozzles as long as adequate coverage is maintained. Always ensure that agitation is maintained until spraying is completed, even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes, resuspend the spray solution by running on full agitation prior to spraying.

**Preemergence:** Apply in a spray volume of 10-80 gals./A.

**Early Postemergence:** Good weed coverage is essential for optimum weed control. Boom height for broadcast over-the-top applications should be based on the height of the crop - at least 15 inches above the crop canopy, but only high enough to give uniform coverage. Apply in a spray volume of 10-30 gals./A. When weed foliage is dense, use a minimum spray volume of 20 gals./A. Flat fan nozzles of 80° or 110° are recommended for optimum postemergence coverage. Do not use floodjet nozzles or controlled droplet application equipment for postemergence applications. Nozzles may be angled forward 45° to enhance penetration of the crop and provide better coverage.

**Spray Drift**

**AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.** The interaction of equipment and weather related factors determine the potential for drift. The applicator is responsible for considering these factors when making an application decision.

Do not apply when weather conditions may cause drift to non-target areas. Drift may result in injury to adjacent crops and vegetation. To avoid spray drift, DO NOT apply

when the wind speed is greater than 10 mph or during periods of temperature inversions.

### Information on Droplet Size

The most effective way to reduce spray drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions.

### Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturers recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

### Application Height

Applications should be made at the lowest height above the target area that still provides uniform coverage of the target. Making applications at the lowest yet effective height reduces exposure of droplets to wind.

### Wind

Drift potential is lowest between wind speeds 10 mph or less. However, many factors, including droplet size, pressure, and equipment type determine drift potential at any given wind speed. **Note:** Local terrain can influence wind patterns.

Leave a sufficient buffer downwind of the application to avoid drift to sensitive crops. This buffer may be untreated corn rows or field border species maintained for this purpose. The width of the buffer needed for a specific application will depend on the wind speed, distance to sensitive crops, and application equipment parameters.

### Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended



droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates, indicates good vertical air mixing.

**Sensitive Areas**

Lumax EZ must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**Cleaning Equipment After Application**

Special attention must be given to cleaning equipment before spraying a crop other than field corn. Mix only as much spray solution as needed.

1. Flush tank, hoses, boom, and nozzles with clean water.
2. Prepare a cleaning solution of 1 gal. of household ammonia per 25 gals. of water. Many commercial spray tank cleaners may be used.
3. Use a pressure washer to clean the inside of the spray tank with this solution. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
4. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
5. Dispose of rinsate from steps 1-3 in an appropriate manner.
6. Repeat steps 2-5.
7. Remove nozzles, screens, and strainers and clean separately in the ammonia solution after completing the above procedures.
8. Rinse the complete spraying system with clean water.

**MIXING PROCEDURES**

---

**CARRIER**

**Preemergence Applications** Either clean water or liquid fertilizers excluding suspension fertilizers may be used as carriers for preemergence applications. If fluid fertilizers are used a compatibility test must be done. See Compatibility Test section for compatibility testing. Even if Lumax EZ is physically compatible with a fluid fertilizer constant agitation is necessary to maintain a uniform mixture during application.

**Postemergence Applications** Use only clean water as the carrier when applying Lumax EZ after field corn emergence. Do not apply Lumax EZ to emerged sweet corn or yellow popcorn or grain sorghum.

**ADDING LUMAX EZ TO THE SPRAY TANK**

The spray tank must be clean, thoroughly rinsed and decontaminated before adding either Lumax EZ alone or with tank-mix partners. If water is used as the carrier use clean water.

**Lumax EZ Applied Alone** When Lumax EZ is used alone add the recommended amount of Lumax EZ to the spray tank when the tank is half full of the carrier then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform mixture.

**Lumax EZ Applied in Tank Mixtures** Refer to the sections on this label for recommended tank mixes. Always refer to labels of the tank-mix partners for mixing directions and precautions. Do not exceed label dosage rates nor combined maximum seasonal doses for mesotrione, S-metolachlor or atrazine. This product cannot be mixed with any product bearing a label prohibition against such mixing. If a tank mixture is used a compatibility test must be done. See Compatibility Test section for details on the procedure for such a test.

If the tank-mix partner is compatible fill the tank half full of the carrier. Start and continue agitation throughout mixing and spraying. All return lines to the spray tank must discharge below the liquid level. Prepare the components and add in the following order:

1. If a wettable powder or dry flowable formulation is used make a slurry with water and add it slowly through the screen into the tank. Agitate during the procedure.
2. If a flowable formulation is used add slowly through screen into the tank. Mixing and compatibility may be improved when a dry flowable is diluted with water before adding to the tank.
3. Add Lumax EZ.

- 4 Add any other tank-mix products next with emulsifiable concentrates added last
- 5 Add an adjuvant last, if needed
- 6 Complete filling the sprayer tank and continue agitation. Apply as soon as possible after spray mixture is prepared. Do not leave mixture in spray tank overnight without agitation or unattended

**TANK MIX COMPATIBILITY TEST**

A compatibility test is recommended before tank mixing to ensure compatibility of Lumax EZ with other pesticides. The following test assumes a spray volume of 25 gals/A. For other spray volumes, make appropriate changes in the ingredients.

**Note** Nitrogen solutions or complete liquid fertilizers, excluding suspension fertilizers, may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use**. Incompatibility of tank mixtures is more common with mixtures of fertilizer and pesticides.

**Test Procedure**

- 1 Add 1.0 pt. of carrier (fertilizer or water) to each of two 1 qt. jars with tight lids. **Note** Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
- 2 To one of the jars, add ¼ tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Compex or Unite (¼ tsp. is equivalent to 2.0 pts./100 gals. spray). Shake or stir gently to mix.
- 3 To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on recommended label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.
- 4 After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15-30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) slurry the dry pesticide(s) in water before addition, or (b) add ½ the compatibility agent to the fertilizer or water and the other ½ to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

- 5 After compatibility testing is complete dispose of any pesticide wastes in accordance with the **Storage and Disposal** section in this label

---

**CROP USE DIRECTIONS**

---

**CORN**

Lumax EZ is recommended for preemergence use for control of most annual grass and broadleaf weeds in field corn field corn seed corn, field corn silage sweet corn and yellow popcorn Lumax EZ may also be applied early postemergence for the control of broadleaf weeds in field corn, field corn seed corn and field corn silage Do not apply Lumax EZ to emerged sweet corn or yellow popcorn or severe crop injury will occur

See Table 1 and 2 for a list of weeds controlled Lumax EZ will not consistently control grasses that are emerged at the time of application

**Lumax EZ Use Rate** The soil organic matter content of the field on which Lumax EZ is to be applied must be known If soil organic matter content is less than 3% use 2.7 qts of Lumax EZ per acre If soil organic matter content is 3% or greater, use 3.25 qts of Lumax EZ per acre Do not apply more than 28 days prior to planting or to field corn higher than 12 inches tall Lumax EZ is not recommended on soils with greater than 10% organic matter or poor weed control may result

**LUMAX EZ APPLIED ALONE**

**Early Preplant** Lumax EZ may be applied up to 14 days prior to planting

**Preemergence Surface** Do not exceed 3.0 qts of Lumax EZ per season Lumax EZ may be applied to the soil surface as a broadcast or banded application

**Early Postemergence** Lumax EZ may be applied after field corn emergence See the **Adjuvants** section of this label for adjuvant recommendations Do not apply early postemergence to field corn in liquid fertilizer or severe crop injury may occur Apply this treatment to small broadleaf weeds (less than 3 inches tall) Occasional field corn leaf burn may result, but this will not affect later growth or corn yield Do not apply Lumax EZ to emerged sweet corn or yellow popcorn or severe crop injury may occur Postemergence applications to field corn must be made before crop reaches 12 inches in height

This product will not provide consistent control of emerged grass weeds For control of emerged grass weeds a grass herbicide tank mix may be required (see tank mix section of this label) Tank mixes of AAtrex can improve control of emerged annual grass and broadleaf weeds Refer to the AAtrex label for weeds controlled and use restrictions

If Bicep II Magnum® Bicep Lite II Magnum® AAtrex (atrazine), Dual Magnum® or Dual II Magnum® alone or in tank mixtures have been applied early preplant, preplant surface preplant incorporated or preemergence limit the Lumax EZ early post application to not exceed a total of 2.5 lbs of active ingredient of atrazine or 3.75 lbs of S-metolachlor active per acre or illegal residues may result

**Split Application** Lumax EZ may be applied as a split application in field corn, field corn seed corn and field corn silage. For a split application program apply one-half of the labeled rate of Lumax EZ prior to crop emergence followed by a second Lumax EZ application at one-half of the labeled rate as a post application after corn emergence. The total amount of Lumax EZ applied in the split application program cannot exceed 2.7 qts /A in soils with <3% OM and cannot exceed 3.25 qts /A in soils with  $\geq$  3% OM. Refer to the **Early Postemergence** section above for instructions on post emergence applications.

## LUMAX EZ TANK MIX COMBINATIONS

### Use of Spray Adjuvants With Tank Mixtures

When Lumax EZ is used as a preemergence herbicide, and before weeds have emerged spray adjuvants have little or no influence on performance. However in burndown situations where the weeds have emerged and the corn has not an adjuvant may be used with Lumax EZ applied alone or when applied in tank mixture with a burndown herbicide as allowed on the individual product labels. Use only those adjuvants approved for agricultural crop use. See the **Adjuvants** section under **Application Procedures** for further instructions.

### Burndown Combinations for Reduced Tillage Situations

In reduced or no-till corn and before the crop has emerged, Lumax EZ tank mixes with Gramoxone Inteon or Touchdown brands (or other glyphosate products such as Roundup brands) will burndown emerged weeds. For best results tank mixes of Lumax EZ plus Gramoxone Inteon should be applied to emerged weeds that are 1-6 inches in height. Consult the Gramoxone Inteon Touchdown brand or glyphosate product label for further information on weeds controlled and application timings.

### Preemergence Tank Mixtures Applied Before Corn Emergence

The tank-mix partners listed in Table 3 may be used in either conventional reduced or no-till systems and be applied by the same methods and at the same timings as Lumax EZ unless otherwise specified in the tank-mix product label. Follow all tank-mix product labels for use rates and restrictions. Perform a compatibility test prior to spraying the tank-mix application. Tank mixtures with 2,4-D are allowed, but should only be done with extreme care with regard to ensuring compatibility before mixing a load. 2,4-D products and even batches vary greatly with regard to compatibility and should be checked each time a water or carrier source, water or carrier temperature, product source or tank mixture recipe is changed.

**Table 3 Tank Mixtures for Preemergence Applications with Lumax EZ**

Tank Mix	Rate (Max )	Objective
AAtrex or other solo Atrazine products	0.5-1.25 lbs a.i./A	Improved broadleaf and grass weed control
Princep®	0.5-1.3 lbs a.i./A	Improved broadleaf and grass weed control
Gramoxone Inteon	See product label	Burndown existing weeds
Touchdown brands	See product label	Burndown existing weeds
Roundup or other glyphosate brands	See product label	Burndown existing weeds
Warrior®	3.84 fl. ozs /A	To control insects, such as cutworm

### Early Postemergence Tank Mixtures Applied After Corn Emergence

The tank-mix partners listed in Table 4 may be used in conventional reduced or no-till systems and can be applied by the same methods and at the same timings as Lumax EZ unless otherwise specified in the tank-mix product label. Follow all tank-mix product labels for use rates and restrictions. Perform a compatibility test prior to spraying the tank mix application. Do not apply Lumax EZ tank mixtures to emerged sweet corn or yellow popcorn.

28/40

**Table 4 Tank Mixtures for Early Postemergence Weed Control with Lumax EZ**

Tank Mix <sup>1</sup>	Rate (Max)	Objective
AAtrex or other solo Atrazine products	0.5-1.25 lbs a.i./A	Improved broadleaf and annual grass weed control
Warrior	3.84 fl oz /A	To control insects such as cutworm
Accent®	As per product label	Emerged grass control
Basis®	As per product label	Emerged grass control
Steadfast®	As per product label	Emerged grass control
Steadfast ATZ®	As per product label	Emerged grass control

<sup>1</sup>Consult the 'Adjuvant' section of this label for recommendations when applying Lumax EZ alone or in tank mixture to emerged field corn

**Lumax EZ Programs with Glyphosate in Glyphosate Tolerant Corn**

Lumax EZ may be applied early postemergence at a rate down to 2 qts /A in tank mixture with a solo glyphosate product (e.g. Touchdown or Roundup brands) that is registered for use over-the-top in glyphosate tolerant field corn (e.g. Roundup Ready or Agrisure™ GT Corn). To minimize weed competition with the crop, target the application of this mixture to weeds in the 1 to 2 inch range. Do not apply this mixture to corn that is greater than 12 inches tall. If the glyphosate product has a built-in adjuvant system (i.e. the product label does not ask for additional adjuvant), only spray-grade ammonium sulfate (AMS) at 8.5 lbs /100 gal should be added to this mixture. If the glyphosate product label calls for an adjuvant in addition to AMS, add a non-ionic surfactant (NIS) at 0.25% v/v and AMS to this spray mixture. Do not add urea ammonium nitrate (UAN), crop oil concentrate (COC) or methylated seed oil (MSO) type adjuvants to these mixtures or crop injury may occur. Follow all directions for use and restrictions on the glyphosate product label.

Alternatively, Lumax EZ may be applied preemergence at a rate down to 2 qts /A as part of a two-pass weed control system when followed by a postemergence application of a glyphosate based product in glyphosate tolerant corn (e.g. Roundup Ready or Agrisure GT Corn). When used in this way, Lumax EZ will provide reduced competition of the weeds listed in Table 1 for a period of 30 or more days, thus improving the timing flexibility and effectiveness of the glyphosate based product application. Follow all directions for use and restrictions on the glyphosate product label.

Lumax EZ may be applied preemergence at 1.5 qts /A as part of a two-pass weed control system when followed by Halex™ GT postemergence in glyphosate tolerant corn (e.g. Roundup Ready or Agrisure GT Corn). Follow all directions for use and restrictions on each product label.

**Lumax EZ Programs for LibertyLink Corn**

Lumax EZ may be applied early postemergence at a rate down to 2 qts /A in tank mixture with Liberty or Ignite and applied over-the-top in field corn designated as LibertyLink. To minimize weed competition with the crop, target the application of this mixture to weeds in the 1 to 2 inch range. Do not apply this mixture to corn that is greater than 12 inches tall. Ammonium sulfate (AMS) may be added as a spray adjuvant as directed on the Liberty or Ignite label. However, AMS should be the only adjuvant added to this tank mixture. Do not add urea ammonium nitrate (UAN), crop oil concentrate (COC), non-ionic surfactants (NIS), or methylated seed oil (MSO) type adjuvants to these mixtures or crop injury may occur. Follow all directions for use and restrictions on the Liberty or Ignite product label.

Alternatively, Lumax EZ may be applied preemergence at a rate down to 2 qts /A as part of a two-pass weed control system when followed by a postemergence application of Liberty or Ignite in field corn designated as LibertyLink. When used in this way, Lumax EZ will provide reduced competition of the weeds listed in Table 1 for a period of 30 or more days, thus improving the timing flexibility and effectiveness of the Liberty application. Follow all directions for use and restrictions on the Liberty or Ignite product label.

**GRAIN SORGHUM**

Lumax EZ can be applied preplant non-incorporated (up to 21 days before planting) up through preemergence for weed control in sorghum that was seed treated with Concep III. For a listing of weeds controlled or partially controlled, see Table 1.

Apply Lumax EZ at a rate of 2.7 qts /A as a broadcast non-incorporated spray beginning at 21 days before planting and up through planting but prior to sorghum emergence. Applying Lumax EZ less than 7 days before sorghum planting will increase the risk of crop injury, especially if irrigation or rainfall is received following the application. Injury symptoms include temporary bleaching of newly emerging sorghum leaves or, in extreme conditions, stunting or partial stand loss. Applying Lumax EZ more than 7 days (but not more than 21) prior to sorghum planting will reduce the risk of crop injury.

If Lumax EZ is applied prior to planting, minimize disturbance of the herbicide treated soil barrier during the planting process in order to lessen the potential for poor weed control in the disturbed soil zone.

Lumax EZ may also be applied as a split application to grain sorghum. For a split application program, apply one-half of the labeled rate of Lumax EZ as a non-incorporated early preplant (7-21 days before planting) followed by a second Lumax EZ application at one-half of the labeled rate as a preemergence application prior to sorghum emergence. The total amount of Lumax EZ applied in the split application program cannot exceed 2.7 qts /A.



If weeds are present at the time of application it is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v or a crop oil concentrate at a rate of 1% v/v be added to the spray solution. In addition to COC or NIS a spray grade UAN at a rate of 2.5% v/v or AMS at a rate of 8.5 lb /100 gals. of spray may be added to the solution for improved control of emerged weeds. If weeds are not emerged at the time of application no additives are recommended.

**Restrictions**

- 1 Do not apply more than 2.5 quarts of Lumax EZ per growing season
- 2 Do not apply Lumax EZ to sorghum grown on sandy soils (sand, sandy loam or loamy sand)
- 3 Do not apply Lumax EZ to emerged grain sorghum or severe injury will occur
- 4 Do not use Lumax EZ in the production of forage sorghum, sweet sorghum (sorgo), sudangrass, sorghum-sudangrass hybrids, or dual purpose sorghum
- 5 Sorghum seed must be treated with Concep III herbicide safener prior to planting or severe crop injury may occur
- 6 In the state of Texas do not apply Lumax EZ to sorghum grown South of Interstate 20 (I-20) or East of Highway 277

---

**STORAGE AND DISPOSAL**

---

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage**

Keep container tightly closed when not in use. Do not store near seeds, fertilizers, or foodstuffs. Lumax EZ can be stored at temperatures as low as -10°F. Keep away from heat and flame.

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

**Pesticide Disposal**

Open dumping is prohibited. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Rinse spray equipment. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of as described above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling [less than 5 gallons]**

Non-refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows empty the remaining contents into application equipment of a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begin to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities by burning If burned stay out of smoke

For minor spills leaks etc follow all precautions on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night If the container is damaged and leaking or material has been spilled follow these procedures

- 1 Cover spill with absorbent material
- 2 Sweep into disposal container
- 3 Wash area with detergent and water and follow with clean water rinse
- 4 Do not allow to contaminate water supplies
- 5 Dispose of according to instructions

**Container Handling[Bulk/Mini-Bulk]**

Refillable container Refill this container with Lumax EZ only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities by burning If burned, stay out of smoke

For minor spills leaks etc follow all precautions on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill, fire, or other emergency, call 1-800-8372, day or night If the container is damaged and leaking or material has been spilled follow these procedures

- 1 Cover spill with absorbent material
- 2 Sweep into disposal container
- 3 Wash area with detergent and water and follow with clean water rinse

- 4 Do not allow to contaminate water supplies
- 5 Dispose of according to instructions

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

---

AAtrex® Agrisure™ GT, Bicep II Magnum® Bicep Lite II Magnum®, Callisto Plant Technology® Concep® Dual Magnum® Dual II Magnum® Gramoxone Inteon® Halax™ GT, Lumax® EZ Princep®, Touchdown®, Warrior®, and the Syngenta logo are trademarks of a Syngenta Group Company

Accent®, Basis®, Steadfast® Steadfast ATZ® and Viton® trademarks of E I du Pont de Nemours and Company

Counter® trademark of BASF Corporation

Ignite® Liberty® and LibertyLink® trademarks of Bayer CropScience

Roundup® trademark of Monsanto Company

©201X Syngenta

For non-emergency information (e.g. current product information), call  
Syngenta Crop Protection at 1-800-334-9481

Manufactured for  
Syngenta Crop Protection LLC  
P O Box 18300  
Greensboro North Carolina 27419-8300

[Base Label]

**RESTRICTED USE PESTICIDE**  
**(GROUND AND SURFACE WATER CONCERNS)**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION. THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER.

**For Ground Application Only**

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

**GROUP 5 15 27 HERBICIDES**

**Lumax® EZ**

**Herbicide**

A Preemergence Herbicide for Control of Annual Grass and Broadleaf Weeds in Field Corn, Field Production Seed Corn, Field Silage Corn, Sweet Corn, Yellow Popcorn, and Grain Sorghum

**Active Ingredients\*:**

S-Metolachlor: (CAS No. 87392-12-9)	27.10%
Atrazine** (CAS No. 1912-24-9)	10.20%
Mesotrione: (CAS No. 104206-82-8)	2.71%
Other Ingredients:	59.99%
<b>Total:</b>	<b>100.00%</b>

\*Active ingredients per gallon: S-metolachlor 2.49 pounds, Mesotrione 0.249 pounds, and Atrazine 0.935 pound.

\*\*Atrazine with a maximum of 0.27% related triazines.

See directions for use in attached booklet.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in Directions for Use section for information about this standard.

**Callisto Plant Technology**

EPA Reg. No. 100-XXXX

EPA Est.

2.5 gallons	120 gallons	220 gallons	330 gallons	_____ gallons [bulk]
Net Contents	Net Contents	Net Contents	Net Contents	Net Contents

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**CAUTION**

Cases moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear protective eyewear. Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**FIRST AID**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOT LINE NUMBER:** For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call **1-800-888-8372**.

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA Chemical Resistance Category Selection Chart.

**Mixers, Loaders, Applicators and other handlers must wear:**

- Coveralls over short-sleeved shirt and short pants

- Chemical-resistant gloves - Category A (e.g. barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, polyethylene, polyvinyl chloride (PVC)  $\geq 14$  mils or Viton®  $\geq 14$  mils)
- Chemical-resistant footwear plus socks
- Chemical-resistant apron when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to the concentrate
- Chemical-resistant headgear for overhead exposure

See engineering controls for additional requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Aerial application is prohibited.

#### **Environmental Hazards**

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. This pesticide contains atrazine, which has been shown to be toxic to aquatic invertebrates. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from treated areas.

#### **Ground Water Advisory**

Lumax EZ contains the active ingredients S-metolachlor, mesotrione and atrazine.

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

S-metolachlor has the potential to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

#### **Surface Water Advisory**

The active ingredients in this product have the potential to contaminate surface water through ground spray drift. Under some conditions, the active ingredients may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly drained or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas,

areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

**Mixing/Loading Instructions [for 2.5 gallons]**

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check valves or antisiphoning devices must be used on mixing equipment.

This product must not be mixed/loaded or used within 50 ft. of wells, including abandoned wells, drainage wells, and sink holes.\*

\*For exceptions to these restrictions, see the **Environmental Hazards** section of the **Precautionary Statements** in attached booklet.

**Mixing/Loading Instructions [for 120, 220, 330 gallons and bulk]**

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check valves or antisiphoning devices must be used on mixing equipment.

This product must not be mixed/loaded or used within 50 ft. of wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing to this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum, 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment

capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site

Additional State imposed requirements regarding well head setbacks and operational area containment must be observed

This product must not be mixed or loaded within 50 ft of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be applied within 66 ft of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 ft around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 ft buffer or setback from runoff entry points must be planted to crop or seeded with grass or other suitable crop.

**Tile-Outletted Terraced Fields Containing Standpipes**

One of the following restrictions must be used in applying atrazine to tile-outletted terraced fields containing standpipes

- 1 Do not apply this product within 66 ft of standpipes in tile-outletted terraced fields
- 2 Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2-3 inches in the entire field
- 3 Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest

**Physical and Chemical Hazards**

Do not use or store near heat or open flame

[For 120, 220, and 330 gallons and bulk]

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller. To the extent permitted by applicable



law Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions To the extent permitted by applicable law (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA and (2) Buyer and User assume the risk of any such use To THE EXTENT PERMITTED BY APPLICABLE LAW SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL

To the extent permitted by applicable law, in no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT**

SYNGENTA and Seller offer this product and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA

---

**STORAGE AND DISPOSAL**

---

Do not contaminate water, food, or feed by storage or disposal

**Pesticide Storage**

Keep container tightly closed when not in use Do not store near seeds, fertilizers, or foodstuffs Lumax EZ can be stored at temperatures as low as -10°F Keep away from heat and flame

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material

**Pesticide Disposal**

Open dumping is prohibited. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Rinse spray equipment. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of as described above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Handling [less than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment of a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc. follow all precautions on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night. If the container is damaged and leaking or material has been spilled follow these procedures:

- 1 Cover spill with absorbent material
- 2 Sweep into disposal container
- 3 Wash area with detergent and water and follow with clean water rinse
- 4 Do not allow to contaminate water supplies
- 5 Dispose of according to instructions

**Container Handling[Bulk/Mini-Bulk]**

Refillable container. Refill this container with Lumax EZ only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc. follow all precautions on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other

emergency, call 1-800-8372, day or night. If the container is damaged and leaking or material has been spilled follow these procedures:

1. Cover spill with absorbent material.
2. Sweep into disposal container.
3. Wash area with detergent and water and follow with clean water rinse.
4. Do not allow to contaminate water supplies.
5. Dispose of according to instructions.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

---

Lumax® EZ, Callisto Plant Technology®, and the Syngenta logo are trademarks of a Syngenta Group Company

©201X Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

LUMAX EZ EZ NP 0112 - bb - 1-10-12

000100.XXXXX.20120110.LUMAX-EZ-NP-0112.pdf