

**PRECAUTIONARY STATEMENT
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. May be fatal if swallowed. Avoid breathing vapors. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling this product. Wash hands after handling. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Treated effluent cannot be discharged into lakes, streams, ponds, or public water systems a discharge permit is obtained. For guidelines, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: STRONG OXIDIZING AGENTS: Mix only with water according to label directions. Mixing this product with green film such as faces, wires, etc. or with ammonia, acids, detergents or other chemicals will release hazardous gases irritating to the eyes, nose and mucous membranes.

Sodium Hypochlorite Solution degrades with age. If used later than ten (10) days after packaging, use a chlorine test kit to insure that the solution will provide the amount of available chlorine desired.

STORAGE AND DISPOSAL:

Keep this product in a tightly closed container when not in use. Store in a cool, dry, well ventilated area away from heat or open flame. In case of decomposition, isolate container (if possible) and flood area with large amounts of water to dissolve all material before discarding this container, but place in trash collection. Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

SODIUM HYPOCHLORITE SOLUTION

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE... 12.5%
INERT INGREDIENTS.....87.5%

DISINFECTANT BACTERICIDE DEODORANT
FOR
FOOD PROCESSING PLANTS, DAIRY PLANTS,
FARMS, HOTELS, RESTAURANTS, TAVERNS
SWIMMING POOLS.

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do not give vinegar or other acids. Do not induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS
PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING.**

PRODUCED BY
TIDEWATER INDUSTRIAL CORPORATION
GREENSBORO, MARYLAND 21639
EPA REGISTRATION NUMBER: 63484-1

NOT REVIEWED
In Accordance with PR Notice 82-2.
Based on Draft Labeling Dated

BEST AVAILABLE COPY

use check
419190

DIRECTION FOR SWIMMING POOL CHLORINATION
For a new pool or for spring start-up superchlorinate with one (1) pint (16 fl. oz.) of sodium hypochlorite solution for each 2000 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.
For Pool Maintenance

- A) Adjust pool water pH to 7.0-7.2 range and maintain
- B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 1.0 ppm. One fourth pint (4 fl. oz.) of this sodium hypochlorite solution for each 4000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.8 to 1.0 ppm available chlorine will depend on temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

DIRECTIONS FOR DISINFECTION OF POTABLE WATER
Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 11 parts softened water. Mix sodium hypochlorite solution with water thoroughly and begin feeding of solution with hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 8.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations.* Check water frequently with a DPD chlorine test kit.

*Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with the disinfectant for a minimum of 2 minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of one fluid oz. (2 tablespoons) of this sodium hypochlorite solution per 5 gallons of water will provide approximately 200 ppm available chlorine by weight. When used as sanitizing for non-porous surfaces such as dishes, glasses, and utensils, clean prior to applying a 200 ppm available chlorine solution.

DISINFECTANT

For disinfection of hard, inanimate surfaces such as floors, walls, and woodwork, apply a 600 ppm available chlorine solution. Thoroughly wet surfaces with the recommended solution for at least 10 minutes. Do not rinse after treatment.

DM 32 63484-1 1/1