



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 30, 2021

Eliot Harrison
Agent for Sterilex
111 lake Front Drive
Hunt Valley, MD 21030

Subject: PRIA Label Amendment – Label Amendment with Efficacy Data to Add New Disinfection, Virucidal, and other minor label claims and changes
Product Name: Sterilex Ultra Disinfectant Cleaner Solution 1
EPA Registration Number: 63761-8
Received Date: 12/15/2020
Action Case Number: 00216864

Dear Eliot Harrison:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. Pursuant to 40 CFR 156.10(a)(6), you must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. See FIFRA section 2(p)(2). If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process, FIFRA section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Assurance.

Page 2 of 2
EPA Reg. No. 63761-8
Action Case No. 00216864

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Perri Moeller by phone at (703) 347-8618, or via email at moeller.perri@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Steven Snyderman". The signature is written in a cursive style with a horizontal line underneath the name.

Steven Snyderman, Product Manager 33
Regulatory Management Branch II
Antimicrobials Division (7510P)
Office of Pesticide Programs

Enclosure: Stamped Label

Sterilex[®] Ultra Disinfectant Cleaner Solution 1

Active Ingredients:

n-Alkyl (C ₁₂ 68%, C ₁₄ 32%) dimethylethylbenzyl ammonium chloride	3.00%
n-Alkyl (C ₁₄ 60%, C ₁₆ 30%, C ₁₂ 5%, C ₁₈ 5%) dimethylbenzyl ammonium chloride	3.00%
Hydrogen peroxide	6.30%

Inert Ingredients:

TOTAL:	<u>87.70%</u> 100.00%
--------	--------------------------

KEEP OUT OF REACH OF CHILDREN

DANGER

See [back panel] [side panel] [booklet] [box] for [additional] [directions for use] [and] precautionary statement

FIRST AID {NOTE TO REVIEWER – In accordance with 40 CFR 156.68(d), all first aid statements, as prescribed, will appear on the front panel of the label.}

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

General Information: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies, call the poison control center at 1-800-222-1222. For general information on this product, call [XXX-XXX-XXXX][hours of service], or contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 63761-8

EPA Est. No. _____ {NOTE TO REVIEWER – this may be printed on container itself}

[Net Contents: {1, 2.5, 5, 15, 30, 55, 275, 305, 330} [Gallons]] {NOTE TO REVIEWER – this may be printed on container itself}

[Net Weight: _____]

[Lot Number: _____][Batch Number: _____] {NOTE TO REVIEWER – this may be printed on container itself}

[NOT FOR SALE OR USE AFTER: _____]

[Expiration Date][Exp. Date] [Expiry Date]

Manufactured by

STERILEX

111 Lake Front Drive

Hunt Valley, MD 21030

Phone (Toll-Free): 1-800-511-1659

Phone: (443) 541-8800

Fax: 410-581-8864

ACCEPTED

08/30/2021

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 63761-8

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage and skin burns. May be fatal if absorbed by the skin. Harmful if swallowed or inhaled. Do not get in eyes, on skin or on clothing. Avoid breathing dust, vapor, or spray mist. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Wear coveralls over long-sleeved shirt and long pants, socks, chemical resistant shoes and gloves, and when mixing and loading wear a chemical-resistant apron. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

PHYSICAL AND CHEMICAL HAZARDS

When exposed to fire, Sterilex Ultra Disinfectant Cleaner Solution 1 may release oxygen and oxides of carbon and/or nitrogen.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep product in closed original container when not in use. Store in a cool area away from potential sources of heat, open flames, sunlight or other chemicals. Store product below 90°F.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed [or pressure rinsed] these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations. Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

[{NOTE TO REVIEWER: *The following language is for containers having a capacity of 5 gallons or less*} Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.]

[{NOTE TO REVIEWER: *The following language is optional for containers having a capacity of 5 gallons or less when "or pressure rinse" appears in the above Container Disposal statement*} Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[{NOTE TO REVIEWER: *The following language is for containers having a capacity greater than 5 gallons*} Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.]

[{NOTE TO REVIEWER: *The following language is optional for containers having a capacity greater than 5 gallons when "or pressure rinse" appears in the above Container Disposal statement*} Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[[For refillable containers]

Refillable container. Refill this container with this product only. Do not reuse this container for any other purpose. Cleaning or pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the owner. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

{NOTE TO REVIEWER: All use sites and marketing claims listed below are meant to be optional language on the final market label.}

{NOTE TO REVIEWER: "ViveSecure" Alternate Brand Names apply to final printed labels intended for use in, and for claims associated with, the Vivarium and Animal Research markets. "Vive" refers directly to use in vivariums and is not meant to imply enhanced safety.}

[Alternate Brand Names for Sterilex Ultra Disinfectant Cleaner Solution 1:]

[Sterilex Ultra CIP]

[Ultra Disinfectant]

[FortiSolve 100]

[ViveSecure 100]

[Alternate Brand Names for Sterilex Ultra Activator Solution:]

[Ultra Soft Metal Activator]

[Ultra Activator]

[Ultra High Foam Additive]

[FortiSolve 200]

[FortiSolve 200 SM]

[ViveSecure 200]

[ViveSecure 200 SM]

[POWERED BY] [BASED OFF OF] [BASED ON] [INCLUDING] [INCLUDES] [USING] [PATENTED] [PERQUAT® TECHNOLOGY]

FOR DISINFECTION AND NON-FOOD CONTACT SURFACE SANITIZATION FOR PUBLIC HEALTH USES, THIS IS PART ONE OF A TWO-PART PRODUCT AND MUST BE USED WITH STERILEX ULTRA ACTIVATOR SOLUTION

FOR BIOCIDES AND NON-PUBLIC HEALTH USES, THE USE OF STERILEX ULTRA ACTIVATOR SOLUTION MAY NOT BE REQUIRED.

PLEASE CONSULT THE DIRECTIONS FOR USE FOR SPECIFIC INSTRUCTIONS.

[ONE-STEP DISINFECTANT – CLEANER] [DISINFECTANT]

[FOOD CONTACT SURFACE DISINFECTANT]

[NON-FOOD CONTACT SURFACE DISINFECTANT]

[SANITIZER]

[NON-FOOD CONTACT SURFACE SANITIZER]

[FOOD CONTACT SURFACE SANITIZER]

[VIRUCIDE§]

[[Federally Inspected] Food Processing Facilities], [[Federally Inspected] Meat And Poultry Plants], [Pet Food Processing Facilities], [Dairies], [Wineries], [Breweries], [Beverage Manufacturing Facilities], [Dairy Farms], [Swine Farms], [Poultry] [And] [Turkey] Farms, [Farm Premises], [Hatcheries], [Refrigerated Trucks^{††}], [Kennels], [Pet Animal Quarters], [Zoos], [Pet Shops], [Animal Laboratories], [Veterinary Facilities], [Animal Care Facilities], [Vivariums]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is for use on washable hard, nonporous surfaces in:]

[hospitals, medical and dental offices, nursing homes, health care facilities, ultrasonic baths (ultrasonic cleaning units), federally inspected food processing facilities, federally inspected meat and poultry plants, dairies, wineries, breweries, beverage manufacturing facilities, dairy farms, swine farms, poultry and turkey farms, farm premises, hatcheries, brooders, harvesters, harvesting equipment, refrigerated trucks^{††}, kennels, pet animal quarters, zoos, pet shops, animal laboratories, veterinary facilities, animal care facilities, transportation terminals, hotels and motels, factories, business and office buildings, barber shops, salons, boats, ships, campers, trailers, mobile homes, homes, kitchens, bathrooms, household areas, cars, buses, trains, taxis, airplanes, restaurants, bars, cafeterias, institutional kitchens, food preparation and storage areas, convenience stores, food storage areas, public rest rooms, institutions, schools and colleges, athletic facilities, sports facilities, gym rooms, locker rooms, dressing rooms, shower and bath areas, exercise equipment, large, inflatable, non-porous plastic and rubber structures (animals, promotional items, moonwalk, slides, obstacle course play equipment, exercise equipment and non-porous wrestling mats).]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is for use on washable hard, nonporous surfaces including:]

[Medical machines (x-ray, MRIs, CAT scanners), and equipment surfaces (that do not penetrate the body or contact the human bloodstream or mucous membranes): This product is not to be used as a terminal Sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.].]

[Steam sterilizer water reservoirs, steam sterilizer water reservoir tubing;]

[Water reservoir tanks, water reservoir pipes, tanks, and piping systems used in food processing;]

[Trench drains, floor drains and other visible drain surfaces;]

[Floors, walls, countertops, stovetops, sinks, appliances, refrigerators (exteriors), plastic and other nonporous cutting boards and chopping blocks; [coolers¹], [ice chests¹]; [nonresidential refrigerator bins¹] (exteriors) for meat, fruit, vegetable and egg; food processing equipment (k-pac equipment, injectors, slicers, knives, steel mesh gloves, deboners, saws, grinders, cutters, racks, dairy equipment, interlocking belts, spiral freezers^{††}, [HVAC units¹], [cooling coils¹], [individual quick freezing (IQF) freezers^{††}], [cooling tunnels¹], [refrigeration units¹], outside surfaces of kitchen equipment, beer fermentation and holding tanks, brewery pasteurizers, wine fermentation tanks, beverage dispensing equipment, beverage transfer lines, bottling or premix dispensing equipment;]

[[Drinking water coolers¹], [ice making machines^{††}], transfer line tubing, water lines, [automatic] watering systems, farm animal nipple drinkers;]

[Cabinets, highchairs, garbage cans, garbage storage areas, refrigerated storage and display equipment (exteriors), tables, picnic tables (non-wooden and finished/sealed or painted), outdoor furniture, chairs, desks, telephones, doorknobs, shower stalls, tubs and glazed tiles, whirlpool bathtubs, bathtubs, sinks, urinals, exterior toilet bowl surfaces, other bathroom fixtures;]

[Kennel runs, nest pads, cages, waterers and feeders, automatic feeders, hauling equipment, dressing plants, loading equipment, farrowing barns and related areas (nursery blocks, creep areas), chutes, feed racks, mangers, troughs, fountains and waterers, forks, shovels, scrapers, barns, pens, stalls, facilities and fixtures occupied or traversed by animals, hatchery areas (egg receiving and holding, setter room, tray dumping, chick holding, processing and loading), trays, buggies, racks, egg flats, trucks, reapers, poultry buildings, ceilings, sidewalls and floors, drinkers and other poultry house related equipment;]

[Coils and drain pans of air conditioning and refrigeration equipment and heat pumps, conductive flooring;¹]

[Nonporous salon/barber tools and instruments (combs, brushes, scissors, blades, pedicure and manicure instruments, pedicure and manicure tubs);]

[Hard nonporous surfaces, listed on this label, that are made of metal, stainless steel, glazed porcelain, glazed ceramic, sealed stone, hard fiberglass (bathtubs, tubs, shower stalls, and sinks), plastic (such as polystyrene, polypropylene), glazed porcelain tiling, enameled surfaces, finished/sealed and painted woodwork.]

Optional Marketing Claims

[General cleaning]

[Specifically formulated for maximum stability]

[Made in [The USA][The United States]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, kills household germs in ten minutes; effective against *Streptococcus pyogenes* (*Strep*), *Pseudomonas aeruginosa* (*Pseudomonas*), *Staphylococcus aureus* (*Staph*) while effectively reducing non-pathogenic mold and mildew.]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, cleans and deodorizes non-porous surfaces by killing odor-causing microorganisms and [preventing] [inhibiting][reducing] the growth of non-pathogenic mold and mildew.]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, [kills up to 99.9999%] pathogens in ten minutes on non-porous surfaces†;

† Effective against][Such as] [*Streptococcus pyogenes* (*Strep*)] [*E. coli*] [*Listeria monocytogenes*] [and] [*Salmonella enterica*]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, inhibits the growth of non-pathogenic mold and mildew on non-porous surfaces, leaving bathrooms and kitchens clean and fresh smelling.]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, eliminates odors on non-porous surfaces caused by non-pathogenic bacteria, mildew, and non-fresh foods.]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, controls non-pathogenic mold and mildew on large, inflatable, non-porous plastic and rubber structures (such as animals, promotional items, moonwalk, slides, obstacle course play equipment, exercise equipment and wrestling mats).]

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, cleans and deodorizes non-porous surfaces in locker rooms, gym rooms and sports facilities by killing odor-causing microorganisms and [preventing] [inhibiting] [controlling] the growth of non-pathogenic mold and mildew.]

[Germicide and disinfectant][Antibacterial] [Fungicidal] [Bactericide] [Bactericidal]

[Staphylocidal^Y] –]Pseudomonacidal^Y] – [Salmonellacidal^Y] – [Virucidal^S] – [Fungistat/Mildewstat]

[Effective against mycotoxin-producing molds†]

†*Aspergillus niger*

[Reduce cross-contamination between treated surfaces [in high traffic areas]]

[Helps reduce [cross contamination] between treated surfaces of [Avian Influenza A] [animal] [avian] virus[es]^S [in [food plants] [animal health facilities]]

^SAvian Influenza A virus, PED virus, PRRS virus, and Newcastle Disease virus

[Kills up to 99.9999% germs, bacteria, and viruses^S [household germs] within 10 minutes

[Effective, economical and convenient germicide]

[Use this product to sanitize or disinfect hard, nonporous surfaces in chicken houses, brooders, and other poultry equipment to help reduce cross contamination of [Avian Influenza A][animal and avian[§]] virus[es] between treated surfaces [in [food plants] [animal health facilities]]

[§]Avian Influenza A virus, PED virus, PRRS virus, and Newcastle Disease virus

[This product is an effective deodorizer.][This product is an essential institutional item]

[Effective against spoilage microorganisms[‡] that can adversely affect product quality.]

[[‡]For a list of organisms, see directions for use.]

[Reduces harmful pathogens found on food processing plant surfaces.]

[Reduces harmful microorganisms on critical meat, poultry, and fruit/vegetable conveyor surfaces.]

[Controls *Avibacterium paragallinarum*, the cause of Avian coryza.]

[Nipple drinker lubricant]

[Disinfects and lubricates nipple drinkers]

[Deep][Powerful] Cleaning

- [Essential] [Deep] cleaner]
- [Makes cleaning easy]
- [Versatile] [Multi-purpose cleaner]
- [Concentrated [power] [foam] to clean]
- [Cost-effective disinfectant]
- [Reduces dairy odors, colorings and flavorings from equipment and milk storage areas]
- [Removes][Washes away] soils such as [dairy soils][turmeric][red sauces][residual food additives]
- [Reduces [appearance of] stains on equipment surfaces]
- [Disinfects and deodorizes by [killing] [most] [up to 99.9999%] germs* and their odors]
- [Quickly removes soap scum from shower tile, fiberglass, chrome and glass, toilet bowls, sinks, tubs and spas]

Promotes quality assurance, cost savings and environmental impact

- [Low foam formulation improves CIP system efficiency and minimizes cycle time]
- [Contact sales representative for ideal application temperatures]
- [Improved productivity as removal of minerals associated with biofilm helps reduce frequency of acid washes and labor needs]
- [Use of this product helps meet compliance standards established by FSMA]
- [Use of this product helps meet regulatory standards outlined in FSIS Compliance Guideline Appendix B through reduction of non-pathogenic spore forming bacteria]
- [Limit wastewater impact with ability to apply at a lower rate for disinfection]
- [This product does not [need][require] hot water to work]
- [Increases heat transfer efficiency through removal of biofilm insulation on heat transfer surfaces]

Helps Protect Valuable Equipment

- [Compatible with] [Safe on] soft metals, including aluminum]
- [Soft metal disinfection and sanitization applications include spiral freezers^{††} and [chillers[†]]]
- [Optimized for use in [spiral freezers^{††}] [chillers[†]] [individual quick freezing (IQF) freezers^{††}] [cooling tunnels[†]]] and other soft metal surfaces]
- [Helps reduce cross contamination between treated surfaces]
- [Compatible with] [Safe on] aluminum surfaces, (including) (specifically): Al 1100, Al 3003, Al 5052, Al 6061]
- [Compatible with] [Safe on] stainless steel and soft metal surfaces including: [Aluminum], [Copper]]
- [Semi-compatible with [Al 7075] [Brass] [Carbon Steel]]]
- [Controls corrosion causing bacteria]
- [Soft metal safe technology helps reduce corrosion and wear]
- [Not for use on galvanized surfaces]
- [Contact Sterilex for full compatibility profile]
- [When mixed with Ultra Soft Metal Activator, this product is [compatible with] [safe on] the following material surfaces: [Al 1100], [Al 5052], [Al 6061], [Al 3003], [Bronze], [Copper], [SS 304], [SS 316]]]
- [This product is semi-compatible with [Al 7075], [Brass], [Carbon Steel]]]
- [Note: This product may also be [compatible with] [safe on] hard non-porous materials that have not been listed on this label. If you have any questions regarding the compatibility of this product, please contact [your local distributor][Sterilex]]]
- [May be corrosive to metal] [when used with Sterilex Ultra Activator Solution]

Biofilm Claims

- [Removes** biofilm from dental unit water lines]
- [Reduces formation of biofilm in [dental unit water lines][animal drinking lines]]
- [Kills biofilm bacteria]
- [Maintains [dental unit water] line effluent water <500 cfu/ml]
- [Controls [bacteria*] [biofilm] found in harborage niches]
- [For control of biofilm found on hard, non-porous [environmental] [surfaces] [including] [but not limited to] [associated with] [restrooms, floors, shower stalls, sink basins, drain covers, walls, countertops, instrument trays in patient care areas of hospitals, healthcare, clean room, and residential environments]]]
- [[Kills] [Controls] slime-forming bacteria]
- [[Kills][Controls] up to 99.9999% of bacteria* in biofilms on a hard, non-porous surface]
- [[Controls] [Kills] up to 99.9999% of bacteria* in biofilms]
- [Reduces up to 99.9999% of bacteria* growing in biofilms]
- [[Proven] [Formulated] to kill up to 99.9999% of bacteria* in biofilms]
- [[Penetrates and removes] [biofilms] [biofilm matrix] [biofilms found in [harborage niches][hard-to-reach areas], [killing][destroying] up to 99.9999% of the [pathogenic] bacteria* living there]
- [Removes biofilm and kills biofilm bacteria* on [all hard non-porous surfaces][food [and non-food] contact surfaces]
- [[Patented technology that is][Proven] [Developed][Approved][Formulated] to [kill][reduce][penetrate]][remove][collapse][breakdown] biofilms]
- [Eliminate up to 99.9999% biofilm associated recontamination by [removing][destroying] the biofilms where pathogens live]
- [[Controls][Reduces] biofilm [formation][buildup][deposits]]]
- [Biofilm disinfectant]
- [Attack and kill bacteria where they live]
- [[Disinfect] [and] [wash away] biofilm matrix]
- [Sterilex Ultra Disinfectant Cleaner delivers 1, 2 punch by disinfecting biofilm matrix and then washing the biofilm away]
- [Increases shelf-life when used as part of a sanitation program]
- [Biofilm control may reduce [product recalls][product losses][food recall risk]]]

- [[Kills][Controls] biofilm [forming][producing] bacteria such as [*Listeria monocytogenes*], [*Pseudomonas aeruginosa*], [*Salmonella enterica*], [*Staphylococcus aureus*] [*E. coli*] [*Campylobacter jejuni*]]
- [[Kills][Controls] bacteria* inside and outside the biofilm]
- [Removes protective biofilm matrix helping to reduce repopulation on treated surface]
- [Seek out and destroy biofilm with Indicon Gel and Sterilex Ultra Disinfectant Cleaner Solution 1]
- [Only biofilm removal controls [dormant] bacteria [hiding] [deep] within the [biofilm matrix][extracellular polymeric substance]
- [Provides 6 log reduction of biofilm bacteria]
- [Effectively removes biofilm in hard to reach areas such as gaskets, probes, filling equipment, etc.]
- [Helps Improve heat transfer efficiency through removal of biofilm [deposits][fouling]]
- [Removes scale [found within][associated with] biofilm]
- [Control of persistent biofilms helps reduce positives and reduces QA/QC time spent finding source of contamination]
- [Biofilm removal reduces need for physical removal methods]

Associated biofilm disclaimers

*[*Pseudomonas aeruginosa* and *Staphylococcus aureus*]

**laboratory studies show >90% removal

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For disinfection and sanitization for public health uses, this is part one of a two-part product and must be used with Sterilex Ultra Activator Solution.

For biocide and non-public health uses, the use of Sterilex Ultra Activator Solution, may not be required.

Ultra Soft Metal Activator may replace Sterilex Ultra Activator Solution in all listed claims and application instructions.

Please consult the directions for use for specific instructions

PRECLEANING INSTRUCTIONS (OPTIONAL STEP):

Although Sterilex Ultra Disinfectant Cleaner Solution 1 provides one-step cleaning and disinfection, precleaning may be desired for some areas, articles and surfaces in certain cases. Prepare a precleaning solution by adding 1 –12.8 fl. oz. of Sterilex Ultra Disinfectant Cleaner Solution 1 and an equivalent amount of Sterilex Ultra Activator Solution per 1 gallon of water, depending on the level of visible soil to be precleaned. [If high foam is desired, refer to High Foam Application Instructions.] Clean surfaces with precleaning solution by pouring, wiping, brushing, scrubbing, foaming, spraying with a coarse trigger sprayer, sponging, immersing, using a clean in place (CIP) system, pumping it through the system, drawing it through the system or mopping. Rinse all food contact surfaces thoroughly with a potable water rinse.

SANITIZATION DIRECTIONS

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, reduces the following bacteria by 99.9% in 3 minutes according to the Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces:

[*Enterobacter aerogenes* [ATCC 13048]]

[*Staphylococcus aureus* [ATCC 6538]]

SANITIZING DIRECTIONS FOR NON-FOOD CONTACT SURFACES:

Add 2 – 4 fl. oz. [59.1 -118.3 mL] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 -118.3 mL] of Sterilex Ultra Activator Solution to 1 gallon [3.79 L] of water (1:1:64 – 1:1:32) (or equivalent use dilution) to sanitize hard, non-porous surfaces. Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces using a cloth, mop, sponge, sprayer or foaming device. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming, sponging, using a clean in place (CIP) system, pumping it through the system, drawing it through the system, mopping or by

immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches [15-22 cm] from surface. [If high foam is desired, refer to High Foam Application Instructions.] Do not breathe spray. Treated surfaces must remain visibly wet for 3 minutes, then wipe with sponge, mop or cloth, rinse, or allow to air dry. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

SANITIZING DIRECTIONS FOR FOOD CONTACT SURFACES:

Add 2 - 4 fl. oz. [59.1 - 118.3 mL] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 - 118.3 mL] of Sterilex Ultra Activator Solution to 1 gallon [3.79 L] of water (1:1:64 - 1:1:32) (or equivalent use dilution) to sanitize hard, non-porous surfaces. Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces using a cloth, mop, sponge, sprayer or foaming device. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming, sponging, using a clean in place (CIP) system, pumping it through the system, drawing it through the system, mopping or by immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches [15-22 cm] from surface. [If high foam is desired, refer to High Foam Application Instructions.] Do not breathe spray. Allow surfaces to remain visibly wet for at least 3 minutes. Rinse all surfaces thoroughly with a potable water rinse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

[[¶]] ONE STEP DISINFECTION AND CLEANING DIRECTIONS FOR HARD, NON-POROUS SURFACES

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is a one-step, cleaner and hospital-use disinfectant at 12.8 - 16.0 fl. oz. [378.5 - 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 [378.5 - 473.2 ml] fl. oz. of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 - 1:1:8), or equivalent use dilution. [If high foam is desired, refer to High Foam Application Instructions.] Bactericidal according to the current AOAC Use-Dilution Test Method modified in the presence of 400 ppm hard water plus organic soil against:

[*Pseudomonas aeruginosa* (*Pseudomonas*) [ATCC 15442]]

[*Staphylococcus aureus* (*Staph*) [ATCC 6538]]

[*Staphylococcus aureus* - Methicillin Resistant [ATCC 33591] [MRSA]]

[*Salmonella enterica* (*Salmonella*) [ATCC 10708]]

[*Salmonella enterica* subspecies *enterica* serovar Enteritidis (*S. Enteritidis*) [ATCC 13076]]

[*Salmonella enterica* subspecies *enterica* serovar Heidelberg (*S. Heidelberg*) [ATCC 8326]]

[*Escherichia coli* O157:H7 [ATCC 35150]]

[*Listeria monocytogenes* [ATCC 19111]]

[*Campylobacter jejuni* [ATCC 29428]]

[*Cronobacter sakazakii* [ATCC 29004]]

[*Burkholderia cepacia* [ATCC 25416]]

APPLICATION INSTRUCTIONS:

To clean and disinfect in one step, remove gross visible soil from all areas, articles and surfaces to be disinfected using a pre-clean, pre-flush, or pre-scrape and, if necessary, presoak. Mix 12.8 - 16.0 fl. oz. [378.5 - 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 - 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming, spraying with a coarse trigger sprayer, sponging, using a clean in place (CIP) system, pumping it through the system, drawing it through the system, mopping or immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches [15-22 cm] from surface. [If high foam is desired, refer to High Foam Application Instructions.] Do not breathe spray. Allow surfaces to remain visibly wet for at least 10 minutes. Rinse all food contact surfaces thoroughly with a potable water rinse. Treated nonfood contact surfaces do not require a rinse and may be wiped with sponge, mop or cloth, rinse, or allow to air dry following the 10 minute contact time. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration. [For initial start-up, repeat on all surfaces for [3] [3-5] consecutive [nights, treatments]]. [For routine maintenance applications treat all surfaces 1x/week, or as needed to maintain environmental counts.]

[ANIMAL HEALTH] DISINFECTION FOR HARD, NON-POROUS SURFACES

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is a [hospital-use][broad spectrum] disinfectant at 2 - 4 fl. oz. [59.1 - 118.3 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 - 118.3 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:64 - 1:1:32), or equivalent use dilution. [If high foam is desired, refer to High Foam Application Instructions.] Bactericidal according to the current AOAC Use-Dilution Test Method modified in the presence of 400 ppm hard water:

[*Pseudomonas aeruginosa* (*Pseudomonas*) [ATCC 15442]]
[*Staphylococcus aureus* (*Staph*) [ATCC 6538]]
[*Salmonella enterica* subspecies *enterica* serovar Enteritidis (*S. Enteritidis*) [ATCC 13076]]
[*Salmonella enterica* subspecies *enterica* serovar Heidelberg (*S. Heidelberg*) [ATCC 8326]]
[*Salmonella enterica* subspecies *enterica* serovar Braenderup (*S. Braenderup*) [ATCC 700136]]
[*Escherichia coli* O157:H7 [ATCC 35150]]
[*Avibacterium paragallinarum* [ATCC 29545]] [Precogning not required]

APPLICATION INSTRUCTIONS:

To clean and disinfect in one step, remove gross visible soil from all areas, articles and surfaces to be disinfected using a pre-clean, pre-flush, or pre-scrape and, if necessary, presoak. Mix 2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming, spraying with a coarse trigger sprayer, sponging, using a clean in place (CIP) system, pumping it through the system, drawing it through the system, mopping or immersion. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-9 inches [15-22 cm] from surface. [If high foam is desired, refer to High Foam Application Instructions.] Do not breathe spray. Allow surfaces to remain visibly wet for at least 10 minutes. Rinse all food contact surfaces thoroughly with a potable water rinse. Treated nonfood contact surfaces do not require a rinse and may be wiped with sponge, mop or cloth, rinse, or allow to air dry following the 10 minute contact time. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration. [For initial start-up, repeat on all surfaces for [3] [3-5] consecutive [nights, treatments]]. [For routine maintenance applications treat all surfaces 1x/week, or as needed to maintain environmental counts.]

[TOUGH][HEAVY DUTY] [SOIL] [INANIMATE SCUM] [INANIMATE MATERIAL] REMOVAL APPLICATION INSTRUCTIONS

Precogning surfaces per the Precleaning Instructions. Mix 64.0 fl. oz. [1892.7 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 64.0 fl. oz. [1892.7 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:2), or equivalent use dilution. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, spraying, sponging, mopping, or immersion. Allow surfaces to remain visibly wet for at least 10 minutes. Do not breathe spray. Rinse all food contact surfaces thoroughly with a potable water rinse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration. For a more concentrated option, prepare a 1:1 solution by mixing equal parts Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution. Do not use Ultra Soft Metal Activator at the 1:1 or 1:1:2 application rate.

HIGH FOAM APPLICATION INSTRUCTIONS

Prepare a 1:1 solution by mixing equal parts Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming, spraying, sponging, mopping, or immersion. Allow surfaces to remain visibly wet for at least 10 minutes. Do not breathe spray. Rinse all food contact surfaces thoroughly with a potable water rinse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration. For a less concentrated, high foam option, mix 64.0 fl. oz. [1892.7 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 64.0 fl. oz. [1892.7 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:2). [Ultra High Foam Additive may be added to mixed solution to enhance foam]. Do not use Ultra Soft Metal Activator at the 1:1 or 1:1:2 application rate.

Kills Biofilm Bacteria [on] [in] [insert hard, non-porous use site]

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, effectively penetrates biofilm and kills the following biofilm bacteria:

Pseudomonas aeruginosa (*Pseudomonas*) [ATCC# 15442]
Staphylococcus aureus (*Staph*) [ATCC# 6538]

APPLICATION INSTRUCTIONS:

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per the [Tough][Heavy Duty][Soil] [Inanimate Scum] [Inanimate Material] Removal Application Instructions. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

FUNGICIDE: SPOILAGE ORGANISMS AND OTHER FUNGI

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution kills the following spoilage organisms on hard, non-porous inanimate surfaces in one step at the concentrations listed in the table below, according to the current AOAC Use-Dilution Test Method modified in the presence of 400 ppm hard water plus organic soil. Allow surfaces to remain visibly wet for at least 10 minutes.

Spoilage Organism	Concentration
[<i>Zygosaccharomyces parabaillii</i> [(ATCC 56075)]]	[4 - 8 oz. [118.3 – 236.6 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 4- 8 oz. [118.3 – 236.6 ml] of Sterilex Ultra Activator Solution per gallon of water (1:1:32– 1:1:16)]
Other Fungus	Concentration
[<i>Trichophyton mentagrophytes</i> [(ATCC 9533)]]	[12.8 - 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon of water (1:1:10 – 1:1:8)]

[If high foam is desired, refer to High Foam Application Instructions.]

§VIRUSES

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is effective against the following viruses on hard, non-porous inanimate surfaces at the concentrations listed in the table below, in the presence of 400 ppm hard water:

Virus	Concentration	Minimum Contact Time	Precleaning Required?
[Avian Influenza]	2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:64 – 1:1:32)	5 minutes	No
[Feline calicivirus]	[12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)]	10 minutes	No
[Norovirus]	[12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)]	10 minutes	No
[Rotavirus]	[12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)]	10 minutes	No
[Newcastle Disease Virus]	[12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)]	10 minutes	No
[Porcine Respiratory & Reproductive Syndrome (PRRS) Virus]	[12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)]	10 minutes	No

[PED Virus]	12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)	10 minutes	Yes. Refer to Precleaning Instructions prior to application.
[Mouse Parvovirus]	12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)	10 minutes	Yes. Refer to Precleaning Instructions prior to application.
[Murine Minute Virus]	12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8)	10 minutes	Yes. Refer to Precleaning Instructions prior to application.

Application Instructions

Mix Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution per table instructions above. Visibly wet surfaces by pouring, wiping, brushing, scrubbing, foaming spraying with a coarse trigger sprayer, sponging, using a clean in place (CIP) system, pumping it through the system, drawing it through the system or mopping. Allow surfaces to remain visibly wet for listed contact time. [If high foam is desired, refer to High Foam Application Instructions.] Do not breathe spray. Rinse all food contact surfaces thoroughly with a potable water rinse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

CONTROL OF NON-PUBLIC HEALTH, VEGETATIVE, SPORE-FORMING BACTERIA

To reduce the number of food spoilage non-public health microorganisms *Bacillus coagulans* [ATCC 7050], *Alicyclobacillus acidoterrestris* [ATCC 49025] and *Geobacillus stearothermophilus* [ATCC 7953], apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

FUNGISTAT/MILDEWSTAT

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, controls the growth of mold and mildew organisms (*Aspergillus niger*, [ATCC# 6275]) and the odors they cause on hard, non-porous surfaces in one step at 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8), or equivalent use dilution, according to the current AOAC Use-Dilution Test Method modified in the presence of 400 ppm hard water plus organic soil. [If high foam is desired, refer to High Foam Application Instructions.] Allow surfaces to remain visibly wet for at least 10 minutes.

[(For food processing or other facilities that have installed entryway sanitizing systems)]

ENTRYWAY SANITIZING SYSTEMS:

Set the entryway system to apply 2 - 16 fl. oz. [59.1 – 473.2 mL] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 16 fl. oz. [59.1 – 473.2 mL] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:64 – 1:1:8), or equivalent use dilution. The entire doorway path must be covered with the [spray] [foam] [footbath]. Ensure a continuous wet blanket of sanitizer solution is delivered to the [floor] [bath]. [If high foam is desired, refer to High Foam Application Instructions.] Allow surfaces to remain visibly wet for at least 3 minutes.

SANITIZATION OF HARD, NON-POROUS [INANIMATE] [ENVIRONMENTAL] SURFACES ON [PERSONAL PROTECTIVE EQUIPMENT] [PPE]:

[Add] [Mix] 2 - 4 fl. oz. [59.1 – 118.3 mL] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 – 118.3 mL] of Sterilex Ultra Activator Solution [per] [to] 1 gallon [3.79 L] of water (1:1:64 – 1:1:32), or equivalent use

dilution. Gently mix for uniform solution. Apply solution to hard, non-porous [inanimate] [environmental] surfaces of the [Personal Protective Equipment] [PPE] with a brush, cloth, mop, sponge, auto scrubber, spray, foam or by immersion. Visibly wet all surfaces to be sanitized. Treated surfaces must remain visibly wet for three (3) minutes. Remove excess solution from equipment prior to storage. Change cloth, sponge or towels frequently to avoid redeposition of soil. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

SHOE FOAM DIRECTIONS:

Apply a foam layer that is approximately 0.5 to 2 inches [1.2 cm to 5 cm] thick at all entrances using a foam generating machine or aerator and a solution containing 2 - 16 fl. oz. [59.1 – 473.2 mL] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 16 fl. oz. [59.1 – 473.2 mL] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:64 – 1:1:8), or equivalent use dilution. [If high foam is desired, refer to High Foam Application Instructions.] Scrape waterproof shoes and boots. Walk through foam. Boots must remain visibly wet for at least 3 minutes. Foam entranceways must be washed and the foam replaced at least daily or when it foam/area appears dirty. Apply foam to entrances of buildings, hatcheries, production and packaging rooms to reduce cross contamination between treated surfaces in the packaging and storage areas of food plants.

CLEANING AND DISINFECTION OF [POTABLE WATER], [FOUNTAIN DRINK] OR [BEVERAGE PREPARATION], [STORAGE], [TRANSFER] AND [DISPENSING] EQUIPMENT:

[Confirm compatibility of Storage, Transfer and Dispensing equipment before use of this product.]

Surfaces must be visibly clean prior to disinfection. Prepare enough solution to complete fill storage, transfer and dispensing equipment by adding 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution to 1 gallon [3.79 L] tap water (1:1:10 – 1:1:8), or equivalent use dilution, in an appropriate container and stir. [If high foam is desired, refer to High Foam Application Instructions.] Completely fill storage, transfer and dispensing equipment with the prepared solution. Allow the mixture to remain in contact with the surfaces of the storage, transfer and dispensing equipment for at least 10 minutes. Drain storage, transfer and dispensing equipment. Rinse storage, transfer and dispensing equipment with potable water.

DISINFECTION AND SANITIZATION OF DRAINS, TRENCH DRAINS, FLOOR DRAINS AND OTHER HARD, NON-POROUS DRAIN SURFACES

Surfaces must be visibly clean prior to disinfection. Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

FOGGING

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, controls odors in hatcheries, farms, animal health facilities, food plants, and other use sites when [fogged] [misted] [applied] at a rate of 2 – 16 fl. oz. [59.1 – 473.2 mL] Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 – 16 fl. oz. [59.1 – 473.2 mL] Sterilex Ultra Activator Solution per gallon [3.79 L] of water.

People, animals, food products, and packaging material must be removed from the room/barn/building or carefully protected prior to fogging [or misting] applications. All surfaces must be cleaned and disinfected in accordance with label directions prior to fogging [or misting]. The area must be vacated of all personnel during [fogging][misting] and for a minimum of 2 hours after fogging [or misting]. All food contact surfaces must be thoroughly rinsed with potable water prior to reuse.

Fog [or mist] desired areas using 1 gallon [3.79 L] per 4000 cubic feet [113.26 cubic meters] of room area with a solution containing 2 – 16 fl. oz. [59.1 – 473.2 mL] Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 – 16 fl. oz. [59.1 – 473.2 mL] Sterilex Ultra Activator Solution to 1 gallon [3.78 L] of water (1:1:64 – 1:1:8), or equivalent use dilution. following fogger manufacturer's directions for use. Electrostatic sprayers, backpack sprayers, or other fine

sprayers (average droplet size <340 micron), may be used for misting applications. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

NOTE: The fog [or mist] generated is irritating to the eyes, skin and mucous membranes. No one is allowed to enter the room or building within two hours of the actual fogging [or misting] unless wearing the appropriate personal protective equipment (PPE). If the room or building must be entered, then the individuals must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants. All surfaces must be cleaned and disinfected in accordance with label directions prior to fogging [or misting].

[Do not use thermal foggers with this product.]

DEODORIZATION DIRECTIONS

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION AND SANITIZATION OF [LIFE SCIENCE FACILITIES], [LABORATORIES], [RESEARCH FACILITIES], [ANIMAL CAGES], [BIOMEDICAL RESEARCH FACILITIES], [CLEANROOMS], [MANUFACTURING EQUIPMENT], [INCUBATORS], [VIVARIUMS], [TERRARIUM, ANIMAL RACKS], AND OTHER HARD, NONPOROUS SURFACES

Remove all animals and feed from premises, room and enclosures. Remove all litter and droppings from floors, walls and surfaces of cages, racks, and other facilities and fixtures occupied or traversed by animals. Empty all feeding and watering appliances. Visibly clean all surfaces with soap or detergent and rinse with water. Optionally, preclean surfaces per the Precleaning Instructions. Saturate all surfaces with a solution of Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution, as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. [If high foam is desired, refer to High Foam Application Instructions.] Immerse all equipment used in handling and restraining animals or equipment used for removing litter and droppings. Ventilate closed spaces. Allow to air dry. Do not introduce animals until treatment has been absorbed, set, or dried. Rinse food contact surfaces with potable water before reuse. Prepare fresh solution of Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution dilution daily

DISINFECTION AND SANITIZATION OF FOOD PROCESSING EQUIPMENT AND HARD SURFACES IN [FOOD PROCESSING FACILITIES], [BREWERIES], [WINERIES] AND [OTHER BEVERAGE MANUFACTURING FACILITIES], [RESTAURANTS], [BARS], [CAFETERIAS], [INSTITUTIONAL KITCHENS], [CONVENIENCE STORES], [FOOD PREPARATION AREAS], [FOOD STORAGE AREAS] AND [FOOD SERVICE AREAS]:

NOTE:

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. Follow all applicable local health and sanitation codes for sanitizing food processing equipment post-disinfection.

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Rinse food contact surfaces with potable water before reuse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION AND SANITIZATION OF HEMP PRODUCTION FACILITIES, STORAGE FACILITIES, GREENHOUSES, PROCESSING EQUIPMENT, IRRIGATION LINES, GROW ROOMS, FARMS, PHARMACEUTICAL-PROCESSING FACILITIES, AND OTHER HARD NONPOROUS SURFACES USED IN THE PRODUCTION/PROCESSING OF HEMP

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Rinse food

contact surfaces with potable water before reuse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION AND SANITIZATION OF [VETERINARY FACILITIES], [ANIMAL CARE FACILITIES], [ANIMAL LABORATORIES], [ZOOS], [PET SHOPS], [KENNELS AND ANIMAL QUARTERS], [FARM AND DAIRY FARM PREMISES], [SWINE PREMISES], [FARM AND DAIRY FARM PREMISE EQUIPMENT], [HARVESTORS], [SWINE PREMISE EQUIPMENT], [ANIMAL PREMISES][SICK ANIMAL BARNs].

Any visibly soiled surfaces should be cleaned prior to disinfection. Remove all animals and feed from premises/room/barn/building, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Visibly clean all surfaces with soap or detergent and rinse with water. Optionally, preclean surfaces per the Precleaning Instructions. Saturate all surfaces with a solution of Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution, as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. [If high foam is desired, refer to High Foam Application Instructions.] Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats and other closed spaces. Allow to air dry. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Rinse food contact surfaces with potable water before reuse. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION OF POULTRY PREMISES, POULTRY EQUIPMENT, BROODERS AND HATCHERIES

Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Visibly clean all surfaces with soap or detergent and rinse with water. Optionally, preclean surfaces per the Precleaning Instructions. Saturate surfaces with the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution, as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. [If high foam is desired, refer to High Foam Application Instructions.] Ventilate buildings, coops, and other closed spaces. Allow to air dry. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Rinse food contact surfaces with potable water before reuse.

Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

SANITIZING [AND] [OR] [BIOFILM REMOVAL* IN] NON-POROUS ANIMAL DRINKING LINES:

This product can be used to [sanitize] [and] [or] [remove biofilms* found in] tanks, pumps or medicators used to deliver the product, piping and tubing associated with water lines in animal, livestock and poultry housing facilities. The water lines can be used for delivering drinking water, air chiller water and cooling spray lines. Prior to use in the water lines, remove all animals, livestock or birds from [premises][room][barn][building]. Do not reintroduce animals, livestock or birds until treatment is complete and water lines have been thoroughly rinsed with potable water.

Ensure all animals are removed from the [premises][room][barn][building]. Apply sanitizing solution to the lines by adding 2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 2 - 4 fl. oz. [59.1 – 118.3 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:64 – 1:1:32), or equivalent use dilution, to the drinking water system. Fill the system by pumping it through the system or drawing it through the system. Make sure there is enough solution to fill the lines. Trigger drinkers to ensure solution flows over the nipple. If a drinker is not working, continue to trigger it until water flows freely. Allow surfaces to remain visibly wet for at least 3 minutes. Flush system thoroughly with a potable water rinse.

[If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

[*[Biofilm removal claim is not approved in California.] Biofilm removal claim for non-public health uses only.]

[CLEANING] [AND] [DISINFECTION] [AND] [BIOFILM REMOVAL* IN] [OF] NON-POROUS ANIMAL DRINKING LINES:

This product can be used to [clean] [and] [disinfect] [and] [remove biofilms* found in] tanks, pumps or medicators used to deliver the product, piping and tubing associated with water lines in animal, livestock and poultry housing facilities. The water lines can be used for delivering drinking water, air chiller water and cooling spray lines. Prior to use in the water lines, remove all animals, livestock or birds from [premises][room][barn][building]. Do not reintroduce animals, livestock or birds until treatment is complete and water lines have been thoroughly rinsed with potable water.

Ensure all animals are removed from the [premises][room][barn][building]. Apply solution to the lines by adding 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8), or equivalent use dilution, to the drinking water system. Fill the system by pumping it through the system or drawing it through the system. Make sure there is enough solution to fill the lines. Trigger drinkers to ensure solution flows over the nipple. If a drinker is not working, continue to trigger it until water flows freely. Allow surfaces to remain visibly wet for at least 10 minutes. Flush system thoroughly with a potable water rinse.

[If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

[*Biofilm removal claim is not approved in California.]Biofilm removal claim for non-public health uses only]

DISINFECTION AND SANITIZATION IN [VEHICLES], [INSTITUTIONS], [INDUSTRIAL FACILITIES], [SCHOOLS], [ATHLETIC FACILITIES], [REST ROOMS], [HOTELS], [MOTELS], [OFFICE BUILDINGS AND OTHER PUBLIC AREAS], AND [HOUSEHOLDS]:

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. For any visibly soiled surfaces that should be cleaned prior to disinfection, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION OF SALON (BARBER) INSTRUMENTS AND TOOLS:

APPLICATION INSTRUCTIONS:

Add 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution to 1 gallon [3.79 L] tap water (1:1:10 – 1:1:8), or equivalent use dilution, in an appropriate plastic container and stir. [If high foam is desired, refer to High Foam Application Instructions.] Completely immerse combs, brushes, rollers, manicure and other hard, nonporous salon instruments and tools for 10 minutes. Allow surfaces to remain visibly wet for at least 10 minutes. Rinse all visible soil with potable water and dry before use. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Commercial AGRICULTURAL OR INDUSTRIAL water system injection for the oxidation of odors when they form: In well systems, Sterilex Ultra Disinfectant Cleaner Solution 1 is injected with a peristaltic pump through a ¼ inch plastic tube (or other comparable injection system) inserted into the well all the way to the well intake screens. The most distant faucet is tested with a supplied test strip until a residual of 20-25 ppm hydrogen peroxide is detected indicating that the system has been thoroughly treated. The injection pump should then be adjusted to deliver 25 ppm on a consistent basis to the water system. When treating the watering system for sick animals, the most distant faucet is tested with a test strip until a residual of 100-1,000 ppm hydrogen peroxide is detected indicating that the system has been thoroughly treated. When this level is reached the injection is immediately reduced until the test strip indicates the 25 ppm is reached and the system is maintained at the 25 ppm level. This process can be repeated monthly to ensure a clean water system and avoid screen clogging.

WATER AND SMOKE DAMAGE REMEDIATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI IN HOSPITAL, INSTITUTIONAL AND INDUSTRIAL AREAS:

Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, is for use in water and smoke damage remediation to control odor causing bacteria and fungi on porous and semi-porous materials such as subflooring, drywall, lumber (trim and frame), tackles strip and paneling.

REMEDICATION PROCESS FOR SEWER BACKUP AND RIVER FLOODING:

Mix 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8), or equivalent use dilution, allowing for the dilution by absorbed water within the affected materials. [If high foam is desired, refer to High Foam Application Instructions.] Remove visible, heavy soil and non-salvageable materials. Saturate all affected materials directly with a sprayer using a coarse spray tip before and after cleaning and extraction. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

REMEDICATION PROCESS FOR POROUS MATERIALS SUCH AS SUBFLOORING, DRYWALL, LUMBER (TRIM AND FRAME), TACKLESS STRIP AND PANELING:

If water damage is from a clean source, extract excess water from affected materials. Mix 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8), or equivalent use dilution, allowing for the dilution by absorbed water within the affected materials. [If high foam is desired, refer to High Foam Application Instructions.] Remove visible, heavy soil and non-salvageable materials. Saturate all affected materials directly with a sprayer using a coarse spray tip. Roll, brush or agitate use solution into affected materials and allow affected materials to remain saturated for ten minutes. Follow with a thorough extraction of the affected materials to remove excess moisture. Dry affected materials rapidly and thoroughly. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

REMEDICATION PROCESS FOR SMOKE DAMAGE AGAINST ODOR CAUSING BACTERIA AND FUNGI IN HOSPITAL, INSTITUTIONAL AND INDUSTRIAL AREAS SUCH AS MOTELS, HOTELS, NURSING HOMES, AND SCHOOLS:

Follow applicable instructions outlined in the preceding sections for water damage remediation. [If high foam is desired, refer to High Foam Application Instructions.] Saturate affected materials with the use solution with enough product to remain visibly wet for 10 minutes. Use proper ventilation during treatment of affected materials.

DISINFECTION AND SANITIZATION IN HOSPITALS, MEDICAL OFFICES, DENTAL OFFICES, DENTAL OPERATORIES, NURSING HOMES AND OTHER HUMAN HEALTH CARE FACILITIES:

NOTE: This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

INSTRUMENT PRE-SOAK:

For use as an instrument pre-soak, prepare a solution by adding 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 - 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] tap water (1:1:10 – 1:1:8), or equivalent use dilution, in an appropriate plastic container and stir. Immerse instruments completely and allow to soak for 10 minutes. Rinse all surfaces visibly with a potable water rinse. Instruments must undergo high-level disinfection or sterilization following pre-soak treatment.

Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION IN ULTRASONIC BATHS:

NOTE:

Use Sterilex Ultra Disinfectant Cleaner Solution 1 to disinfect hard nonporous non-critical objects compatible with Ultrasonic Cleaning Units.

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Optionally, preclean surfaces per the Precleaning Instructions. [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

FOR PRECLEANING OF WHIRLPOOL BATHTUBS, JACUZZIS, SPAS, POOLS, AND FOOTBATHS circulate a solution of 0.5 fl. oz. [14.8 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 and 0.5 fl. oz. [14.8 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] tap water, or equivalent use dilution, through tub, spa, pool, or bath and associated piping and filters at a temperature > 90°F [32.2°C]. [If high foam is desired, refer to High Foam Application Instructions.] Drain tub, spa, pool, or bath and rinse with water.

CLEANING AND DISINFECTION OF WHIRLPOOL BATHTUBS, JACUZZIS, SPAS, POOLS, AND FOOTBATHS After using unit, drain and fill with a use solution of 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 12.8 – 16.0 fl. oz. [378.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:10 – 1:1:8), or equivalent use dilution, to just cover the intake valve. Start the pump to circulate the solution. Wash down the unit sides, seat of the chair lift, and any related equipment with a clean swab, brush or sponge. Treated surfaces must remain visibly wet for a minimum of 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse disinfected surfaces with fresh water. Repeat for visibly soiled units. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

DISINFECTION OF [DENTAL UNIT WATER][ANIMAL DRINKING][ANIMAL WATER] LINES:

IMPORTANT:

Always use Sterilex Ultra Disinfectant Cleaner Solution 1 with Sterilex Ultra Activator Solution.

This product is to be used in conjunction with regular testing of [dental unit water] [animal drinking water] .

Consult the manufacturer of the [dental unit] [water line system] before use of this product about compatibility of Sterilex Ultra Disinfectant Cleaner Solution 1 with the [dental unit] [water line system].

INITIAL START-UP TREATMENT: Treat the [Dental Unit Water (DUWL)] [Animal Drinking] [Animal Water] Lines [for three consecutive nights].

ROUTINE TREATMENT: Treat Dental Unit Water (DUWL) [Animal Drinking] [Animal Water] Lines [one night/week] after Initial Start-up treatment.

Note: If two or more weekly treatments are skipped, repeat initial start-up treatment.

HOW TO APPLY STERILEX ULTRA DISINFECTANT CLEANER SOLUTION 1 WITH STERILEX ULTRA ACTIVATOR SOLUTION:

- 1) Add 1 - 2 fl. oz. [29.6 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 1 - 2 fl. oz. [29.6ml] of Sterilex Ultra Activator Solution and 10 fl. oz. [295.7ml] of water (1:1:10 – 1:1:5), or equivalent use dilution, in a measuring cup or other empty container and stir. Pour appropriate amount of the combined mixture into an empty external [dental unit] water bottle. Do not use PET bottles (See note below). [If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.
- 2) Run Sterilex Ultra Solution through the system until the mixture appears at the end of the air/water syringes and handpiece lines. Always remove the handpiece. It may be advisable to remove coupler depending on the type of coupler. Contact manufacturer for specific recommendations.
- 3) Allow the Sterilex Ultra mixture to sit in the unit overnight (Place the ends of water lines into a sink or cuspidor in case any pink mixture drips overnight).

- 4) At the beginning of the next workday, discard any remaining Sterilex Ultra mixture. Rinse the external water bottle with water.
- 5) Fill the external water bottle with water. Flush each line (air/water syringe, handpiece lines) until the bottle is empty.

Note: Do Not Use Thin Walled Polyethylene Terephthalate (PET) Bottles. We Recommend Either A High Density Polyethylene Bottle With A Minimum Thickness Of 0.08 Inches [0.20 cm] Or A High/Low Density Polyethylene Blend With A Minimum Thickness Of 0.14 Inches [0.36 cm].

[++][DISINFECTION AND SANITIZATION OF SPIRAL FREEZERS, COOLING UNITS, COOLING TUNNELS, EVAPORATIVE CONDENSERS, REFRIGERATED TRUCKS, INDIVIDUAL QUICK FREEZING (IQF) FREEZERS, AND OTHER COIL SURFACES

Apply Sterilex Ultra Disinfectant Cleaner Solution 1 and Ultra Soft Metal Activator as a disinfectant, per One Step Disinfection and Cleaning Directions, or as a sanitizer, per Sanitization Directions. Visibly soiled surfaces should be cleaned prior to disinfection.. Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration. Surfaces must be defrosted and at room temperature prior to application.]

[1][Allow surfaces to come to room temperature.]

Optional, Non-Label Marketing Claims

Emerging Viral Pathogen Claims

[Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution,][This product] qualifies for emerging viral pathogen claims per the EPA’s ‘Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA-Registered Disinfectant Labels’ when used in accordance with the appropriate use directions indicated below. [Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution,][This product] meets the criteria to make claims against certain emerging viral pathogens from the following viral categories:

- Enveloped viruses
- Large non-enveloped viruses
- Small Enveloped viruses

Emerging Viral Pathogen Claim – Terms of Registration

Emerging Viral Pathogen Claim	Supporting Virus(es)
Enveloped Virus Families	Rotavirus or Feline calicivirus
Large, Non-Enveloped Virus Families	Feline calicivirus
Small, Non-Enveloped Virus Families	Mouse Parvovirus and/or Murine Minute virus

- **Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, has demonstrated effectiveness against viruses similar to [insert name of emerging virus(es)] on hard, non-porous surfaces. Therefore, Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, can be used against [insert name of emerging virus(es)] when used in accordance with the directions for use against [name of supporting virus(es)] on hard, non-porous surfaces. Refer to the [insert CDC or OIE] website at [insert pathogen-specific website address] for additional information.**
- **[Insert name of illness/outbreak] is caused by [insert name of emerging virus(es)]. Sterilex Ultra Disinfectant Cleaner Solution 1, when mixed with Sterilex Ultra Activator Solution, kills similar viruses and therefore can be used against [insert name of emerging virus(es)] when used in accordance with the directions for use against [name of supporting virus(es)] on hard, non-porous surfaces. Refer to [insert CDC or OIE] website at [insert pathogen-specific website address] for additional information.**

{NOTE TO REVIEWER: The “Biocide and Non-Public Health Uses” section title will always precede the application instructions below on final printed labels.}

BIOCIDES AND NON-PUBLIC HEALTH USES

For the biocide and non-public health uses below, the use of Sterilex Ultra Activator Solution is not required. Sterilex Ultra Activator Solution may be used as an adjuvant for Sterilex Ultra Disinfectant Cleaner Solution 1 in biocide and non-public health uses to adjust pH, inhibit corrosion and enhance cleaning.

Penetrates, Removes and Destroys up to 99.9999% Biofilm and other Organic Contaminants

For use as a [Bactericide], [Slimeicide], [Mildewstat] [Mildewcide][Odor] and [Algicide] [in Recirculating Cooling Water Systems,] [in Evaporative Cool Cells,] [in Brewery Pasteurizing Systems,] [in Air Washers,] [in Water-Jet Looms,] [Tanks and Piping used to store and convey Industrial Process Water or Aqueous Mixtures not related to food processing,] [Deionizer Water Supply Systems not used in food processing, for human drinking water or in dental lines,] [in Drains,] [in Chair Side Evacuator Traps and Pump Filter Traps].

Controls Mold and Mildew

RECIRCULATING COOLING WATER SYSTEMS, INCLUDING COOLING TOWERS, EVAPORATIVE CONDENSERS, EVAPORATIVE COOL CELLS, DAIRY SWEET WATER SYSTEMS AND BREWERY PASTEURIZERS. Effectively removes and controls [biofilm] [slime] [bacteria] [fungi] [mildew] [odor] [algae] [and other] organic contaminants in dairy sweet water systems and brewery pasteurizers. **[Biofilm claim not approved in California]**

DOSAGE RATES. Initial Dose for badly fouled systems: Add 0.5 to 4 gal. (500-4000 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water in the system. If cleaning is desired or if the pH of the system is below 8, optionally add simultaneously Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1. Repeat until control is evident.

Routine Dose, when microbial control is evident: Subsequent slug additions of 5 oz. to 64 fl. oz. (40-500 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water (and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) are employed every 2 to 5 days or as needed. The frequency of addition depends on the relative amount of bleed-off, the quality of the makeup water and rate of inflow of airborne or other contaminants. Make slug additions in the sump of water cooling towers.

Continuous Dosage: For continuous, or semi-continuous, low level dosage of Sterilex Ultra Disinfectant Cleaner Solution 1, add 1 oz. to 20 oz. (8-165 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water (and if cleaning is desired or the pH of the system is below 8, Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1).

PULP AND PAPER MILLS. For use only in the production of pulp and paper that will not come in contact with food. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in pulp and paper mill fresh and seawater influent, water systems, wastewater treatment systems, non-potable water systems, and other process water. [If high foam is desired, refer to High Foam Application Instructions.] **[Biofilm and biocidal claims not approved in California]**

DOSAGE RATES. Initial Dose for badly fouled systems: Add 3 to 16 gal. (3100-16150 ppm) of **Sterilex Ultra Disinfectant Cleaner Solution 1** per 1000 gallons of water in the system. If cleaning is desired or if the pH of the system is below 8, optionally add simultaneously Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1. If necessary, apply a solution containing 0.4 to 2 oz. Sterilex Ultra Disinfectant Cleaner Solution 1 per gallon of water (and if cleaning is desired or if the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of [Sterilex Ultra Disinfectant Cleaner Solution 1) onto interior surfaces of the system that are not continuously submerged.

Routine Dose, when microbial control is evident: Subsequent slug additions of 8 oz. to 50 gal. (65-3100 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water (and if cleaning is desired or if the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) are employed every 1 to 5 days as needed. The frequency of addition depends on the quality of the makeup water and rate of inflow of airborne or other contaminants.

Continuous Dosage: For continuous low level dosage of Sterilex Ultra Disinfectant Cleaner Solution 1, add 1 oz. to 20 oz. (8-165 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water (and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the

volume of Sterilex Ultra Disinfectant Cleaner Solution 1).

AIR WASHERS. For use only in industrial air washing systems that maintain effective mist eliminating components. Effectively removes and controls [biofilm] [slime] [mildew] [algae] [odor] [and other] organic contaminants in Industrial Water Scrubbing Systems. [If high foam is desired, refer to High Foam Application Instructions.] [**Biofilm and biocidal claims not approved in California**]

DOSAGE RATES. Initial Dose for badly fouled systems: Add 3 to 16 gal. (3100-16150 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water in the system. If cleaning is desired or if the pH of the system is below 8, optionally add simultaneously Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1. If necessary, apply a solution containing 0.4 to 2 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon of water onto interior surfaces of the system that are not continuously submerged. Repeat until control is evident.

Routine Dose, when microbial control is evident: Subsequent slug additions of 28 oz. to 3 gals. (225-3100 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1 are employed every 1 to 5 days or as needed. The frequency of addition depends on the relative amount of bleed-off, the quality of the makeup water and rate of inflow of airborne or other contaminants. Make slug additions in the sump of Industrial Water Scrubbing Systems.

DYE MACHINES. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in Textile Dyeing Machines. [If high foam is desired, refer to High Foam Application Instructions.] [**Biofilm and biocidal claims not approved in California**]

DOSAGE RATES. Add 16 to 40 gal. (16150-38500 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water in a dye storage tank. If cleaning is desired or the pH of the mixture is below 8, optionally add simultaneously Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1. Circulate the mixture from the dye storage tank through the dye machine and all associated piping and other dye storage tanks. If necessary, apply a solution containing 2 to 5 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon of water onto surfaces of the system that are not continuously submerged.

WATER-JET LOOMS. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in Water-Jet Looms. [If high foam is desired, refer to High Foam Application Instructions.] [**Biofilm and biocidal claims not approved in California**]

DOSAGE RATES. Initial Dose for badly fouled systems: Prepare a mixture of 2 to 5 fl. oz. [59.1 – 147.9 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 per gallon [3.79 L] of water. If cleaning is desired or if the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1. Pump the mixture through the Water-Jet Loom and all associated piping. If necessary, apply a solution containing 2 to 5 fl. oz. [59.1 – 147.9 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon [3.79 L] of water onto surfaces of the system that are not continuously submerged.

Subsequent Dose, when microbial control is evident: Subsequent slug additions of 5 to 50 fl. oz. (40-406 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per 1000 gallons of daily water use are employed every 1 to 5 days or as needed. The frequency of addition depends on the quality of the makeup water and rate of inflow of airborne or other contaminants.

TANKS, SUMPS, AND OTHER HOLDING VESSELS, PIPING, WATERLINES AND INDUSTRIAL EQUIPMENT USED TO STORE AND CONVEY INDUSTRIAL OR COMMERCIAL WATER OR AQUEOUS MIXTURES. Do Not Use in equipment that will contact food. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in Tanks, Piping, Waterlines, and Industrial Equipment used to store and convey Industrial Process Water or Aqueous Mixtures. [If high foam is desired, refer to High Foam Application Instructions.] [**Biofilm and biocidal claims not approved in California**]

DOSAGE RATES. Dose for badly fouled systems: Fill equipment as completely as possible with a solution of 8 to 80 gal. (8215-70000 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per 1000 gallons of system capacity (or equivalent use dilution). Circulate and then drain system. Rinse system thoroughly. If necessary, apply a solution containing 1 to 10 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon onto surfaces of the equipment that are not submerged.

Subsequent Dose, when microbial control is evident: Subsequent slug additions of 5 oz. to 8 gals.. (40-8215 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per 1000 gallons of system capacity (or equivalent use dilution) are employed every 1 to 5 days or as needed. The frequency of addition depends on the quality of the makeup water and rate of inflow of airborne or other contaminants.

BIOFILM REMOVAL IN ANIMAL DRINKING LINES:

This product can be used to clean and remove biofilms found in tanks, pumps or medicators used to deliver the product, piping and tubing associated with water lines in animal, livestock and poultry housing facilities. The water lines can be used for delivering drinking water, air chiller water and cooling spray lines. Prior to use in the water lines, remove all animals, livestock or birds from [premises][room][barn][building]. Do not reintroduce animals, livestock or birds until treatment is complete and water lines have been thoroughly rinsed with potable water.

Ensure all animals are removed from the [premises][room][barn][building]. Apply solution to the lines by adding 1.0 - 16.0 fl. oz. [29.5 – 473.2 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 1.0 - 16.0 fl. oz. [29.5 – 473.2 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] of water (1:1:128 – 1:1:8), or equivalent use dilution, to the drinking water system. Fill the system by pumping it through the system or drawing it through the system. Make sure there is enough solution to fill the lines. Trigger drinkers to ensure solution flows over the nipple. If a drinker is not working, continue to trigger it until water flows freely. Allow surfaces to remain visibly wet for at least 3 minutes. Flush system thoroughly with a potable water rinse.

[If high foam is desired, refer to High Foam Application Instructions.] Mix fresh regularly. If the Sterilex Ultra Disinfectant Cleaner Solution 1 and Sterilex Ultra Activator Solution mixture must be used after 8 hours, use testing kit to ensure appropriate active ingredient concentration.

Biofilm removal claims [at rates below 1:1:10 are][is] not approved in California

WATER SUPPLY SYSTEMS (WATERLINES) NOT USED IN FOOD PROCESSING, FOR HUMAN DRINKING WATER OR DENTAL LINES. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other]organic contaminants in Tanks, Filters and Piping used to store and convey water.
[Biofilm and biocidal claims not approved in California]

DOSAGE RATES. Add 3 to 100 gal. (3100-86361 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of system capacity (or equivalent use dilution). Circulate for at least one hour and then drain system. Rinse system thoroughly. If necessary, apply a solution containing 0.4 to 2 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the system is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon onto surfaces of the equipment are not submerged. [If high foam is desired, refer to High Foam Application Instructions.]

Controls Slime Forming Bacteria and Biofouling

Penetrates and Removes Biofilm and Other Organic Contaminants. Use as a Bactericide, Slimicide, Mildewstat, and Algicide in:

DRAINS. Effectively removes and controls plugging and odors caused by accumulation of [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in drains. Prevents stoppage and odors. **[Biofilm and biocidal**

claims not approved in California. See DISINFECTION AND SANITIZATION OF DRAINS, TRENCH DRAINS, FLOOR DRAINS AND OTHER HARD, NON-POROUS DRAIN SURFACES instructions for drain use instructions in California]

DOSAGE RATES. Initial Dose for badly fouled systems. This product can be poured, foamed, wiped, brushed, applied using a clean in place CIP system, pumping it through the system or mopping. Apply from 6 fl. oz. to 15 fl. oz. [177.4 – 443.6 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 per gallon [3.79 L] of water to drain. Repeat until control is evident. Optionally, pour from 1 fl. oz. [29.6 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 around or into one-inch [2.54 cm] diameter or smaller drains up to 90 fl. oz. [2661.6 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 around or into three-inch [7.62 cm] diameter drains. If cleaning is desired, add simultaneously Sterilex Ultra Activator Solution at the same volume as Sterilex Ultra Disinfectant Cleaner Solution 1 (1 – 90 fl. oz.) [29.6 – 2661.62 ml]. For foam cleaning, add 6-15 oz. [177.4 – 443.6 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 and an equal amount of Sterilex Ultra Activator Solution per gallon [3.79 L] of water to a foaming device and foam the mixture into the drain.

Subsequent Dose: When microbial control is evident, subsequent additions of 1-6 fl. oz. [29.6 – 177.4 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 per gallon of water are employed every 7 days or as needed. This product can be poured, foamed, wiped, brushed, applied using a clean in place CIP system, pumping it through the system or mopping. If cleaning is desired, add simultaneously Sterilex Ultra Activator Solution at the same volume as Sterilex Ultra Disinfectant Cleaner Solution 1 (1 – 6 fl. oz.) [29.6 – 177.4 ml] per gallon [3.79 L] of water. Optionally, subsequent additions of 1 – 6 fl. oz. [29.6 – 177.4 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 into one-inch [2.54 cm] diameter or smaller drains up to 6 fl. oz. [177.4 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 into three-inch [7.62 cm] diameter drains, are employed every 7 days or as needed. If cleaning is desired, add simultaneously Sterilex Ultra Activator Solution at the same volume as Sterilex Ultra Disinfectant Cleaner Solution 1 (1 – 6 fl. oz.) [29.6 – 177.4 ml]. For foam cleaning, add 1 – 6 oz. [29.6 – 177.4 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 and an equal amount of Sterilex Ultra Activator Solution per gallon [3.79 L] of water to a foaming device and foam the mixture into the drain.

CHAIR SIDE EVACUATOR TRAPS AND PUMP FILTER TRAPS. Effectively removes and controls plugging and odors caused by accumulation of [biofilm] [slime] [mildew] [odor] [algae] [and other organic contaminants] in chair side evacuator traps and pump filter traps. Prevents stoppage and odor. [**Biofilm and biocidal claim not approved in California]**

DOSAGE RATES. Initial Dose for badly fouled systems. Mix 10 fl. oz. [295.7 ml] of Sterilex Ultra Disinfectant Cleaner Solution 1 and 10 fl. oz. [295.7 ml] of Sterilex Ultra Activator Solution into one gallon [3.79 L] of hot water at the end of the working day. Draw all of the solution into the trap and allow to stand overnight. Repeat until control is evident.

Subsequent Dose: When microbial control is evident, subsequent slug additions of 1.5 - 2 fl. oz. [44.4 – 59.1 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 and 1.5 - 2 fl. oz [44.4 – 59.1 ml] of Sterilex Ultra Activator Solution mixed into one gallon [3.79 L] of hot water are employed every 7 days or as needed.

FOR PRECLEANING OF WHIRLPOOL BATHTUBS circulate a solution of 0.5 fl. oz. [14.8 ml] Sterilex Ultra Disinfectant Cleaner Solution 1 and 0.5 fl. oz. [14.8 ml] of Sterilex Ultra Activator Solution per gallon [3.79 L] tap water through tub and associated piping and filters at a temperature > 90°F [32.2°C].

[DECORATIVE POOLS, FOUNTAINS AND WATER DISPLAYS][WHIRLPOOL BATHTUB, JACUZZI, SPA, POOL, AND FOOTBATH WATER SYSTEMS]. Effectively removes and controls [biofilm] [slime] [mildew] [odor] [algae] [and other] organic contaminants in [decorative pools, fountains and water displays] [whirlpool bathtub, Jacuzzi, spa, pool, and footbath water systems]. [If high foam is desired, refer to High Foam Application Instructions.] **[Biofilm claim not approved in California]**

DOSAGE RATES. Initial Dose for badly fouled systems. After using unit, drain and fill with a use solution of 4 to 80 gal. (4000-70000 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per 1000 gallons of system capacity [to just cover the intake valve]. [Start the pump to circulate the solution]. Circulate [solution] and then drain system. Rinse system thoroughly [with clean water]. If necessary, apply a solution containing 1 to 10 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon onto surfaces of the equipment that are not submerged.

Subsequent Dose, when microbial control is evident After using unit, drain and fill with a use solution of 5 fl. oz. to 4 gal. (40 - 4000 ppm) of Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per 1000 gallons of system capacity to just cover the intake valve. Start the pump to circulate the solution. Circulate and then drain system. Rinse system thoroughly. If necessary, apply a solution containing 1 to 10 fl. oz. Sterilex Ultra Disinfectant Cleaner Solution 1 (and if cleaning is desired or the pH of the mixture is below 8, optionally add Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1) per gallon onto surfaces of the [equipment][unit] that are not submerged. [Continuous Dosage (Decorative Pools, Fountains and Water Display use only). For continuous, or semi-continuous, low level dosage of Sterilex Ultra Disinfectant Cleaner Solution 1, add 1 oz. to 20 oz. (8-165 ppm) Sterilex Ultra Disinfectant Cleaner Solution 1 per 1000 gallons of water (and if cleaning is desired or the pH of the system is below 8, Sterilex Ultra Activator Solution at the rate of 1 to 3 times the volume of Sterilex Ultra Disinfectant Cleaner Solution 1).]

[Procedure for Initial Dose

- 1.) Measure the temperature of the Spa water and raise spa water > 90degrees
- 2.) Remove Filter Cartridges or isolate sand/DE filters for duration of the treatment.
- 3.) Add Pulsar® Biofilm Removal System (Solution 1) at a rate of 1 gallon per 250 gallons of spa water to reach a desired residual of 4000 ppm
- 4.) Let circulate for 2 minutes
- 5.) Add Pulsar® Biofilm Removal System (Solution 2) at a rate of 1 gallon per 250 gallons of spa water to reach a desired residual of 4000 ppm
- 6.) Run jets – for 5 minutes and turn off
- 7.) Circulate water for the balance of treatment 6 to 10 Hours/overnight.
- 8.) Begin to drain treatment while scrubbing all submerged surfaces with brush to remove any remaining biofilm. Drain to waste.
- 9.) Fill spa with fresh water, circulate through jets to remove any dislodged organic material/soil and drain to waste.
- 10.) Fill spa above water jets and circulate. If foam is present, add de-foamer and circulate until foam is no longer present. Rinse and drain to waste.
- 11.) Fill spa with water, rebalance chemistry and add 75g (PER 100 gallons of spa water) of calcium chloride to increase hardness by approx. 200ppm.]

[Procedure for Monthly Maintenance Dose

- 1.) Measure the temperature of the Spa water and raise spa water > 90degrees
- 2.) Remove Filter Cartridges or isolate sand/DE filters for duration of the treatment.
- 3.) Add Pulsar® Biofilm Removal System (Solution 1) at a rate of 0.5 gallons per 250 gallons of spa water to reach a desired residual of 2000 ppm
- 4.) Let circulate for 2 minutes
- 5.) Add Pulsar® Biofilm Removal System (Solution 2) at a rate of 0.5 gallons per 250 gallons of spa water to reach a desired residual of 2000 ppm
- 6.) Run jets – for 5 minutes and turn off
- 7.) Circulate water for the balance of treatment 6 to 10 Hours/overnight.
- 8.) Begin to drain treatment while scrubbing all submerged surfaces with brush to remove any remaining biofilm. Drain to waste.
- 9.) Fill spa with fresh water, circulate through jets to remove any dislodged organic material/soil and drain to waste.
- 10.) Fill spa above water jets and circulate. If foam is present, add de-foamer and circulate until foam is no longer present. Rinse and drain to waste.
- 11.) Fill spa with water, rebalance chemistry and add 75g (PER 100 gallons of spa water) of calcium chloride to increase hardness by approx. 200ppm.]



LISTED 153111 (D1) [Sterilex Ultra Disinfectant Cleaner Solution 1]



LISTED 153115 (D1) [Sterilex Ultra CIP]