



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

64014-9

Date of Issuance:

JUN 25 2007

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Harpoon

Name and Address of Registrant (include ZIP Code):

Dr. Roger S. Webb
Florida Silvics, Inc.
950 S.E. 215th Ave.
Morriston, FL 32668

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Decision, EPA has reregistered the product listed above. This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Generic data requirements that have been identified in the RED as confirmatory data for the active ingredient contained in the subject product will continue to be tracked and assessed in the Agency's continual reassessment of pesticides.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing of your label, which ever occurs first, must bear the new revised label.

A stamped copy of the label is enclosed for your records.

Continued on Page 2

Signature of Approving Official:

Date:

for

Meredith Laws, Chief
Insecticide-Rodenticide Branch
Registration Division (7505P)

Page 2
Notice of Reregistration (continued)
EPA Reg. No. 64014-9

The Agency has completed the cumulative assessment for the organophosphate pesticides (OPs) and finalized the Interim Reregistration Eligibility Decisions (IREDs) for the OP pesticides. The assessment and supporting documents may be found on the Agency's web site at www.epa.gov/pesticides/cumulative and in the docket (EPA-HQ-OPP-2006-0618).

If you have any questions or comments regarding this letter, please contact Stephanie Nguyen at (703) 605-0702 or via e-mail at Nguyen.Stephanie@epa.gov.

Meredith F. Laws, Chief
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

3 8 7

**RESTRICTED USE PESTICIDE
Due to Reproductive Effects**

For sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, equipment repair, and equipment cleaning. Certified applicators must ensure that all persons involved in these activities under their direct supervision are informed of the precautionary statements.

HARPOON®

**FOREST, ORNAMENTAL, NON-CROP-BEARING AND CHRISTMAS TREE INSECTICIDE
Containing META SYSTOX®-R Insecticide for Effective Systemic Protection**

This product is an organophosphate Insecticide

ACTIVE INGREDIENT	By Wt.
Oxydemeton methyl (S-(2-(Ethysulfinyl) ethyl) 0,0 dimethyl phosphorothioate).....	50%
OTHER INGREDIENTS	50%
	<u>100%</u>

Contains aromatic petroleum distillate
Meta Systox-R is a Reg. TM of the Gowan Co.

EPA REG. 64014-9

EPA EST. NO. 64014-FL-001

STOP: READ THE ENTIRE LABEL BEFORE USE

**FOR USE BY COMMERCIAL ARBORISTS
(APPLICATORS)**

KEEP OUT OF REACH OF CHILDREN

DANGER



POISON

ACCEPTED

JUN 25 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
As amended, for the pesticide
Registered under
EPA Reg. No. 64014-9

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. •Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> •Call poison control center or doctor immediately for treatment advice. •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Also, contact 1-800-457-2022 for emergency treatment information.	
NOTE TO PHYSICIAN	
<p>Oxydemeton methyl is a cholinesterase inhibitor. Signs of poisoning include a sense of tightness in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea. If signs of cholinesterase inhibition appear, atropine is antidotal. Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema that may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically. 2-PAM also may be used in conjunction with atropine but should not be used alone. The solvent, methyl isobutyl ketone (MIBK), contains petroleum distillate and vomiting may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

SEE REAR PANEL FOR PRECAUTIONARY STATEMENTS

NET CONTENTS: Each microinjection unit contains 3 mL (0.1 oz.) of Meta Systox-R 50% Specialty Concentrate. There are 100 microinjection units per case.

Manufactured and Sold by
 FLORIDA SILVICS, INC. (dba Tree Tech® Microinjection Systems).
 950 SE 215th Avenue, Morriston, FL 32668
 352-528-5335

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. May be fatal if absorbed through the skin. May be fatal if inhaled. Harmful if swallowed. Do not get in eyes, skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

REPEATED EXPOSURE TO CHOLINESTERASE INHIBITOR SUCH AS THOSE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP WORK, and get help right away.

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber or polyvinyl chloride. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and handlers must wear:

- Coveralls over short sleeved shirt and short pants,
- Chemical-resistant gloves made of barrier laminate or butyl rubber,
- Socks and chemical-resistant footwear,
- Goggles, face shield or safety glasses,

- A chemical-resistant apron for loading, and cleaning equipment.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not contaminate water by cleaning equipment or disposal of wastes.

AVISO

PRECAUCION AL USUARIO: Si Usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. Workers must follow the PPE instructions as described in this label when installing or removing HARPOON microinjection units.

The requirements in this box only apply to the uses of this product that are covered by the Workers Protection Standard (WPS).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Since HARPOON is only applied directly into individual tree stems, there is no restricted worker re-entry interval associated with the use of this product. **THE RESTRICTED-ENTRY INTERVAL (REI) FOR THIS PRODUCT IS "0" HOURS.**

IMPORTANT: Read these entire **DIRECTIONS FOR USE** and **CONDITIONS OF SALE** before using HARPOON insecticide.

GENERAL DIRECTIONS

Applying HARPOON using the Tree Tech microinjection system:

HARPOON label and injection instructions must be read and understood prior to use or installation of Tree Tech Microinjection

Spruce	Needle scale	Apply when crawlers appear.
	Adelgids Mites	Apply in May-June. After mite eggs hatch.

Systems' microinjection units. Failure to follow these instructions may lead to injury to the installer or other persons as well as mechanical or phytotoxic damage to treated trees. The following instructions must be heeded to ensure safe and effective use of the injectors containing HARPOON insecticide.

1. Protective eye wear (goggles, safety glasses or face shield) and chemical-resistant gloves such as barrier laminate or butyl rubber must be worn while handling or installing the microinjection units. See sections dealing with protective clothing.
2. When properly installed, the microinjection unit generates internal pressure resulting in the flow of HARPOON solution through the dispenser tube. The microinjection unit must never be activated unless installed correctly and securely in the tree to be treated.
3. Microinjection units containing HARPOON may require up to several minutes or more to empty depending on the health of the treated tree and local weather conditions. Never assume that microinjection units have depressurized completely because they appear nearly empty or empty. Microinjection units may be rotated in place to determine if any liquid remains in the unit. When removing microinjection units, individuals must wear proper eye protection and chemical-resistant gloves such as barrier laminate or butyl rubber. The individual should then cover the microinjection unit with one hand near the point of insertion into the stem while grasping the barrel end of the microinjection unit with the other hand. The microinjection unit should be turned slightly as it is slowly withdrawn from the tree. Applicators shall not leave Tree Tech Microinjection Systems microinjection units unattended while in use. If microinjection units do not drain within a few hours, carefully remove each unit and place in the accompanying heavy-duty plastic bag for disposal in accordance with state and local regulations. Remove microinjection units promptly after treatment.
4. Discard used microinjection units according to **INJECTOR (CONTAINER) DISPOSAL** in **STORAGE AND DISPOSAL** statement.
5. Disinfect drill bit between the injection of each tree with a mixture of 1 part household bleach to 4 parts of water.

USE RESTRICTIONS

Do not inject trees that are less than two inches in diameter.

Do not apply this product through any type of irrigation system.

Do not inject trees within two weeks of any other spray or soil chemical treatment. Do not treat trees that are suffering from stress such as lack of moisture or herbicide damage.

Do not use on trees that will produce food within the year following treatment (such as nuts, berries, maple syrup, etc.).

SITES, PESTS AND APPLICATION TIMING

Plant	Insect/Pest	Remarks
Ash	Lecanium scale	Treat any time from early spring through October. Repeat after new invasion.
Cedars	Bark beetles	Treat any time from early spring through October. Repeat after new invasion.
Conifers, general	Borers Bagworm	Treat as soon as insects appear Treat as soon as insects appear
Cottonwoods	Aphids	Apply when wingless forms appear.
Douglas-fir	Cone moths Engraver beetles	Apply to heavy, current cone crop trees during or shortly after eggs are laid. Treat any time from early spring through October. Repeat after new invasion.
Elms	Elm leaf beetle Lecanium scale Obscure scale Tea scale Cankerworm	Apply after eggs are present or during early larval stage. Treat any time from early spring through October. Repeat after new invasion. Treat as soon as caterpillars appear.
Hemlock	Hemlock woolly adelgid	Apply in early spring (April/May) after the crawler stage appears. Treat again in mid-summer if the infestation is still present.
Juniper	Bark beetles	Treat any time from early spring through October. Repeat after new invasion.
Pines (Except pinon pines)	Bark beetles Engraver beetles (6 spined) Engraver beetles (Monterey) Flathead borer Mountain pine beetle Red turpentine beetle Black turpentine beetle Jeffrey pine needle miner Mites Nantucket pine tip moth Pine needle aphid Pine needle scale	Treat any time from early spring through October. Repeat after new invasion.
Redwood	Bark beetles Mites Needle scale	When active infestation sites are present. Apply after mite eggs hatch. Apply when crawlers appear.
Spruce	Adelgids Mites	Apply in May-June. After mite eggs hatch.

Installing Microinjection Units

1. Determine the number of microinjection units to be installed based upon a) the target insect pest(s), b) stem circumference at breast height, and c) the registered dosage rate as administered by proper spacing of microinjection units around the stem circumference. Unless otherwise noted, microinjection units should be installed in the stem and root flares every 6 inches around the trunk near the ground line, i.e., 6.0-to-8.0 inches (15-to-20 cm) from the soil surface.
2. Using a portable electric drill (600-800 rpm capacity is preferred) with a sharp, clean 1/64-in. (0.4-cm) bit, the installer should drill a hole at the correct stem circumference spacing to a depth of 1/4-to-1/2 in. (0.60-to-1.3 cm) into the wood (xylem) under the bark. A slight downwardly angle is recommended for more complete drainage of the microinjection unit.
3. After reaching the proper depth range, the drill bit should be withdrawn carefully to avoid dislodging bark fragments around the exterior opening of the hole. The rear barrel portion of the microinjection unit should be compressed partially (about 1/2 in. or 1/3 cm.) and the microinjection unit should be inserted into the hole. Placing the plastic installation cap over the rear barrel end, strike the cap with a plastic hammer to seat the microinjection unit firmly in the hole. If the microinjection unit is not properly positioned in the hole, strike the cap again until correctly seated. By striking the microinjection unit, the back end of the feeder tip is forced back into the funnel-shaped section dislodging a septum that allows the HARPOON solution to flow from the microinjection unit into the tree. When the microinjection unit is positioned correctly in the tree and the internal septum is dislodged to permit the flow of HARPOON, remove the cap and, if necessary, push the rear barrel portion of the unit further downwardly until it is flush with the edge of the locking mechanism. This engages the locking mechanism that pressurizes the microinjection unit and assists in the evacuation of HARPOON from the microinjection unit and movement into the vascular system of the tree.
4. Each hole should be drilled and a microinjection unit installed without undue delay. After the microinjection unit is properly seated, it should be activated. This sequence minimizes the flow of tree sap or resin into the hole prior to injection.
5. Disinfect the drill bit between the injection of each tree with a mixture of 1 part household bleach to 4 parts water.

RECOMMENDED APPLICATIONS

Install one (1) Tree Tech microinjection unit containing HARPOON insecticide for every 6 inches of tree trunk circumference.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of microinjection units.
Open dumping is prohibited.

PESTICIDE STORAGE: Store microinjection units at room temperature (45 -to- 74 degrees F) away from heat and open flame. Do not freeze. Store microinjection units in a locked, cool, dry storage area out of the reach of children. In spill or leak incidents, keep unauthorized people away. Wash down with large amounts of water.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

INJECTOR (CONTAINER DISPOSAL):

IF EMPTY: Do not reuse this container.

IF PARTLY EMPTY: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Used microinjection units may be placed in the heavy-duty plastic bag that accompanies each case of microinjection units. The bag must be properly sealed, placed into the original shipping carton and returned freight prepaid for disposal to Tree Tech Microinjection Systems, 950 SE 215th Ave., Morriston, FL 32668.

WARRANTY AND DISCLAIMER STATEMENT

To the extent consistent with applicable law, Tree Tech Microinjection Systems warrants that this product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions. Buyer and user assume all risk from use of this product when such use is contrary to label instructions. Read and follow label directions carefully.

HARPOON is a registered trademark of Florida Silvics, Inc.
Label revised 09/2006