

ORTHO BOLERO 8 EC
EPA Reg. No. 239-2450
Submitted: June 11, 1987


COPY A

ENDANGERED SPECIES RESTRICTION


The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of its habitat have been identified by county:

Pocketbook Pearly Mussel
(Potamilus capox)

Arkansas (Cross, Crittenden, Lee,
Mississippi, Poinsette,
St. Francis)



62499-20



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if swallowed. Avoid contact with skin and clothing. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, wash eyes immediately with plenty of water. See a doctor if irritation persists. If swallowed, give a small amount of water to drink, make person vomit, and call doctor. Note to Physicians: Emergency Information - call 1-800-233-3737.

ENVIRONMENTAL HAZARDS

This product is toxic to shrimp. Do not apply directly to lakes, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

INSTRUCTIONS FOR USE

Do not violate Federal law to use this product in any manner inconsistent with its labeling.

Read ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

PRECAUTIONS

Store pesticide in original container.
Do not put concentrate or dilute into food or drink containers.
Store in cool, dry place.
Protect from excessive heat.
Help with any spill, leak, fire or exposure involving this product, call day or night (415) 233-3737.

HERBICIDE DISPOSAL

Residue resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Recycle (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or dispose of by other procedures allowed by state and local authorities.



ORTHO BOLERO® 8 EC

(HERBICIDE)

Active Ingredient	By Wt.
*Thiobencarb	84%
Inert Ingredients	
*S-[(4-chlorophenyl)methyl] diethylcarbamothioate	16%
Contains 8 pounds S-[(4-chlorophenyl)methyl] diethylcarbamothioate per gallon.	

BOLERO®—Reg. TM of Kumiai Chemical Industry Co., Ltd. for Thiobencarb Herbicide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

BEST AVAILABLE COPY

Chevron Chemical Company © 1984

Ortho Agricultural Chemicals Division

San Francisco CA 94120-7144

Product 5281

Made in U.S.A.

Form 10129-G

EPA Est. 239-MO-1

EPA Reg. No. 239-2450

NET CONTENTS 5
GALLONS

WEED CONTROL IN DOWN SEEDED RICE
MISSISSIPPI AND ARKANSAS.

Application Directions—For Dry Seeded Rice in Mississippi and Arkansas.

Late Preemergence Application (Preemergence) — For control of watergrass, Sprangletop (Lepidochloa spp.), Apply prior to rice emergence, generally 10 to 14 days after planting of the rice. Apply 4 pts. per acre after soils have been sealed by rainfall or well prepared seedbeds free of large clumps at time of application. Do not apply BOLERO 8 EC to cracked soil.

Early Postemergence Application—For control of grass. Apply 4 pts. of BOLERO 8 EC per acre. It has developed no further than the rice has fully emerged at time of application.

Do not apply BOLERO 8 EC to stressed rice. Do not apply BOLERO 8 EC to second rice crop. Water Management—After application, necessary to prevent crusting and dry spots. Do not flood as soon as the rice plants are fully emerged.

Application Equipment

Aircraft: Apply BOLERO 8 EC in no more than 10 to 20 gals. per acre.

Ground Sprayers: Apply in 10 to 20 gals. per acre.

Precautions: Do not apply more than 10 gals. per acre per year. Do not plant subsequent crops within 6 months of last application. Do not farm out of crayfish will be practiced. In Texas and other areas where the agricultural practice, do not apply BOLERO 8 EC to rice crop. Water drained directly from rice fields to irrigate other crops.

Application to stressed rice can result in growth inhibition, delayed maturity and other factors include but are not limited to temperatures below 65° F., or above 95° F., deficiency, high salt content, high pH, above field capacity while rice seed is in the field, poor field drainage or deep water.

Do not apply under conditions involving forage or other plantings that might be rendered unfit for sale, use or consumption.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the label and is reasonably fit for use. Chevron neither makes, nor authorizes anyone to make, any other warranty of fitness, MERCHANTABILITY, or representation, including this material.

2. Critical and unforeseeable factors prevent it from eliminating all risks in the use of chemicals. Such risks include, but are not limited to, plants and crops to which the material is applied, and damage caused by such risks occur even though the user follows the use as stated herein and even follows the label. Buyer and user acknowledge liability (except those caused by Chevron) for such risks.