


Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0080. Approval expires 05-31-98

	United States Environmental Protection Agency Washington, DC 20460	Registration Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 250993
----------------------------------------------------------------------------------	---------------------------------------------------------------------------------	------------------------------------------------------------------------	----------------------------------------

Application for Pesticide - Section I

1. Company/Product Number 64321-1	2. EPA Product Manager LaRocca	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) BioKill brand insecticide	PM# a3	
5. Name and Address of Applicant (Include ZIP Code) Bio-Environmental Products Corporation 768 Walker Road, Ste. 301 Great Falls, VA 22066 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
SEP 29 1997

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
NOTIFICATION: 1) Compliance with PR Notice 96-6 re pet pesticide product label statements. 2) Additional brand names. 3) Expand marketing copy. 4) Add pests (hornet/wasps/yellow jackets) via notification per PR Notice 95-2.

This notification is consistent with the provisions of PR Notices 95-2 and 96-6 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 95-2 and 96-6 and 40 CFR 152.46, this product may be in violation of FIFRA and BioEnvironmental Products Corporation may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Rebecca M. Horton	Title Consultant	Telephone No. (Include Area Code) 540-342-4148
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped) _____
2. Signature Rebecca M. Horton	3. Title Consultant	
4. Typed Name Rebecca M. Horton	5. Date 8/28/97	

2911

[FRONT PANEL]

BIO KILL® brand INSECTICIDE

[Additional Brand Names: Multi-Purpose BioKill brand insecticide; Multi-Purpose BugBusters]

For commercial use
 Makes up to 16 gallons
 For residential use
 Water-based Permethrin insecticide
 Odorless in application.
 For use INDOORS on houseplants and in indoor landscaping
 A READY-TO-USE and DILUTABLE Permethrin-based emulsion insecticide
 For use OUTDOORS (residential site only) in gardens, on ornamental trees, shrubs and flowers, on lawns.
 For use outdoors, in homes and non-food areas of food/feed processing establishments, hotels, motels, commercial buildings, schools, warehouses, theaters, office buildings and kennels
 Will not stain water-safe fabrics and surfaces
 For use indoors and outdoors on a wide variety of crawling and flying insects
 Use on DOGS and CATS to control fleas and ticks
 Can be used as a DIP for fleas on DOGS
 Use on Poultry to control lice and mites
 Provides effective broad-spectrum control of building and garden insect pests
 Effective for up to four weeks
 Multipurpose, ready-to-use, water based bug killer
 Biodegradable, non-flammable, do-it-yourself product
 Easy to use - just spray affected areas
 Kills bugs on contact, indoors and outdoors - broad spectrum insecticide includes: cockroaches, mosquitoes, ants, spiders, flies, moths, fleas, lice, bedbugs, crickets, grasshoppers and many, many other listed species
 No odor - Non-staining - No sticky residue
 Long lasting - residual effect up to 4 weeks
 Sold in over 30 countries
 Ready-to-use, water-based bug killer
 Kills *crawling* insects: cockroaches, ants, fireants spiders, fleas, ticks, silverfish, lice, beetles, bedbugs, crickets and more!
 Kills *flying* insects: flies, moths, mosquitoes, gnats, black flies, hornets, wasps, and more!

ACTIVE INGREDIENT:

*Permethrin	0.25%
-------------------	-------

INERT INGREDIENTS:	99.75%
	100.00%


*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate.
 Cis/trans ratio: max. 28% (+/-) cis and min. 72% (+/-) trans.

[NOTE TO EPA: Front panel of label may include graphics representing product use; e.g.: insects controlled]

EPA Reg. No. 64321-1
 MASTER LABEL
 Approved: Oct. 21, 1996
 Notification: Aug. 28, 1997

NOTIFICATION
 SEP 29 1997

EPA Est. _____
 Page 1 of 40 Pages



3 of 11

Keep Out of Reach of Children

CAUTION

See back/side/other panel(s) (& accompanying booklet) for additional precautionary statements.

NET CONTENTS _____

[BACK/SIDE/OTHER PANEL(S) &/OR BOOKLET, WHEN NECESSARY]

READ ENTIRE LABEL BEFORE EACH USE.

COMMERCIAL APPLICATIONS:

Hotels, Motels, Commercial buildings, Schools, Warehouses, Theaters, Office Buildings, Kennels, Drains, Apartments and Condominium buildings, and non-food contact areas of food-feed processing establishments.

Inside and outside landscaping

Controls bugs inside garbage bins, garbage dumpsters and in garbage chutes

Controls bugs in hallways, common areas and inside apartments, office units and public areas

HOUSEHOLD APPLICATIONS:

House plants and indoor landscape

Inside - along baseboards, in carpets, mattresses, around cupboards, etc.

Outside - in gardens, on ornamental flowers, lawns, vegetables and fruit trees.

DOMESTIC ANIMALS:

Dogs and cats to control fleas & ticks

Poultry to control lice, fleas & ticks

Livestock - protects beef and dairy cattle, goats, sheep and horses from horn flies, face flies, mosquitoes and gnats.

[NOTE TO EPA: When marketed for use on dogs or cats, the following precautionary statement will be added to label:


PRECAUTIONARY STATEMENTS

Hazards to Domestic Animals

CAUTION: Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.]

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater. This pesticide is toxic to honey bees and other beneficial pollinators exposed to an application. Do not apply when bees are actively visiting blooming plants (vegetables, flowers, fruit/ornamental trees). Do not allow this product to come in direct contact with beehives at anytime.



4 9 11

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

INDOOR APPLICATIONS:

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. The product may be used in non-food/feed areas of food/feed establishments such as garbage rooms, lavatories, floor drains (to sewer entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

In the home, cover all food handling surfaces, cover or remove all food and cooking utensils or wash thoroughly after treatment.

Not For Use In Federally Inspected Meat and Poultry Plants.

Do not apply to classrooms while in use.

Surface Spraying: Provides control of crawling and flying household insect pests for up to 4 weeks. To control *cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, crickets, cigarette beetles, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, drugstore beetles, mealworms, grain mites and cadelles:* Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

To control *carpet beetles:* Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and under baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control *bedbugs:* Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Allow all sprayed articles to dry thoroughly before use.

To control *fleas* (adult and larvae) and *ticks* (adult and nymphs): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, beds, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. Make a test application to upholstery or drapery fabrics in an inconspicuous place before use. Allow to dry thoroughly before allowing pets to contact treated areas.

To control *clothes moths:* Remove and brush thoroughly any infested articles from storage and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage areas directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly by keeping nozzle at least 3 feet from fabric to avoid possible staining. Allow all sprayed articles to dry thoroughly before use.

To kill *lice* and *louse eggs:* Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

Industrial Warehouse Fogging: For area control of crawling and flying insect pests for up to 4 weeks.

To control *spiders, millipedes, centipedes, sowbugs, pillbugs, ants, firebrats, cigarette beetles, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, mealworms, Indian meal moths, coried grain beetles, tobacco beetles, grain mites and cadelles*: Close all doors and windows and fog at the rate of 1 ounce per thousand cubic feet, moving toward the exit. Fill room with fog. Direct to all parts of the room especially windows and other light sources which attract these insects. Leave area and keep closed for a period of 2 hours before reentry. Open and ventilate before reoccupying. Fogged area must be cleared of all personnel and pets while fogging is taking place. This product is to be used solely for warehouse space used to store non-food/feed materials.

OUTDOOR APPLICATIONS:

For control of Garden Insects. For use on roses, dahlias, asters and other ornamentals to kill *Japanese beetles* as well as *whiteflies, aphids, lace bugs, ants, exposed thrips, spiders, red mites and leafminers*. May be used on plants in small gardens as a multi-purpose spray and for spot treatments of incipient insect infestations to prevent spreading into larger garden areas.

When spraying plants, do not hold nozzle closer than 18 inches. Use a sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperature exceeds 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Wasps, Hornets, Yellow Jackets: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, including nest openings in ground, bushes, under attic rafters and other cracks and crevices which may harbor pests. Saturate nest openings and contact as many insects as possible. Repeat as necessary.

ON DOGS & CATS:

USE ONLY ON DOGS OR CATS. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

On Dogs. To control *fleas, ticks and lice*: Start spraying at the tail, moving the sprayer rapidly and making sure that the animal's body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat every three weeks. **As a dip for dogs:** Using one part product and three parts water, make up enough material to thoroughly soak animal. Wet animal with water before treatment. Sponge, dip or pour diluted product on the animal until fur and skin are thoroughly wet, making sure that the animal's entire body is treated, including the legs and underbody. Do not treat animal's face. Do not use on puppies less than 12 weeks old.

On Cats. To control *fleas*: Start spray at the tail, moving the sprayer rapidly and making sure that the animal's body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or on face. Avoid contact with the genitalia. Repeat every 16 days. **DO NOT DIP CATS.** Do not use on kittens less than 12 weeks old.

ON LIVESTOCK:

To protect beef and dairy cattle, goats, sheep and horses from *horn flies, face flies, mosquitoes and*

gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control *stable flies, horse flies and deer flies*: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give control protection.

To control *blood-sucking lice*: Apply to infested area of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control *poultry lice*: Spray roosts, walls and nests or cages thoroughly. Follow by spraying over the bird with a fine mist.

To control *bedbugs and mites in poultry houses*: Spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

DIRECTIONS FOR USE AS A DILUTABLE PRODUCT FOR USE ON INDOOR AND OUTDOOR PLANTS:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or lawns and grounds.

For use on ornamental plants, mix BIO KILL 0.25% with water in the following manner:

<u>Finished Spray</u>	<u>BIO KILL</u>	<u>Water</u>
0.015%	8 ounces (1 cup)	1 gallon
	2 ounces (4 Tbs)	1 quart

Note: Do not use more BIO KILL than recommended, as this may cause some plant injury. Use this dilution unless otherwise specified.

CONTACT AND RESIDUAL SPRAY - Apply a finished spray of BIO KILL. Use sufficient spray to obtain full coverage of all foliage. Adjust spray volume used to plant size.

INDOOR APPLICATION: To protect plants in indoor landscaping, spray directly on insects to kill *leafrollers, armyworms, cutworms, cabbageworms* on chrysanthemums; *fungus gnats* on velvet plants. Spray as directed above.

Spray as directed above to control *aphids, armyworms, corn earworms, fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale, spider mites, spring cankerworms, exposed thrips and whiteflies* on the following: Ageratum, Ardisia, Aster, Azalea, Baby's breath, Begonia, Birdsnest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Crown of thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English ivy, Fuchsia, Gladiolus, Grape ivy, Hypoestes Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-back plant, Poinsettia, Portulaca, Pothos, Prayer plant, Purple passion, Rabbits foot fern, Rose, Schefflera, Snapdragon, Snake plant, Statice, Velvet plant, Verbena, Zinnia.

If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars and, therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as

7 7 11

described above, wait 24 hours, and check for possible injury.

Repeat applications may be necessary to control citrus blackflies on non-bearing citrus; adult Japanese beetles on roses.

OUTDOORS: Use 8 ounces (1 cup) of BIO KILL 0.25% and add to 1 gallon of water to make 0.015% finished spray.

CONTACT & RESIDUAL SPRAY - Apply a finished spray of BIO KILL. Use sufficient spray to obtain full coverage of all foliage. Adjust spray volume used to plant size.

Spray in the late afternoon or evening when temperature ranges from 50°F to 78°F, and when there is little or no wind. Spray at the first indication of insect pests. On those plants that will flower during the season, the spray schedule should be completed before blossom formation. Repeat as necessary; use intervals of 4 to 8 days. BIO KILL application can be made up to the day of collection.

Spray as directed above to control *aphids, cicadas, corn earworms, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, bagworms, spider mites, exposed thrips and whiteflies* on the following plants: Ageratum, Arizona cypress, Aster, Azalea, Begonia, Coleus, Common ninebark, Common snowberry, Elm, Exacum, Euonymous, English ivy, Gladiolus, Gold bells, Hypoestes, Ivy, Lilac, Marigold, Mock orange, Nannyberry, Orchid, Palm, Pansy, Pea shrub, Petunia, Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Taxus, Wax plants, Weeping fig, Zinnia.

For ornamental trees grown around the home only: Use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: *pine moths, pine needleminers, pine beetles and needle scales* on pine; *budworms* on conifers; *oakworms* on oak; *fall webworms* on cherry, elm and honeysuckle; *aphids* on tulip poplar; *citrus black flies* on non-bearing citrus; *tussock moths* on Douglas and grand firs; *leafminers* on birch; *elm leaf beetle and elm spanworms* on elms.

To kill *tent caterpillars*: Application should be made when caterpillars are young and tents are first noticed. For best results, apply in late afternoon or evening, when caterpillars have returned to their tents. Apply BIO KILL FULL STRENGTH. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

FOR USE ON LAWNS: For control of *ants (including fire ants), armyworms, chinch bugs, crickets, cutworms, Japanese beetle grubs, grasshoppers, mole crickets and sod webworms* consult the following chart:

<u>Pest</u>	<u>Dilution</u>
Armyworms	4.5 oz. per gal., 10 gal./1000 sq. ft.
Cutworms	1 oz. per gal. 10 gal./1000 sq. ft.
Chinch bugs	"
Japanese beetle grubs	"
Mole crickets	"
Sod webworms	"
Crickets	6.4 oz. per gal., 3 gal./1000 sq. ft.
Grasshoppers	"

EPA Reg. No. 64321-1
MASTER LABEL
Approved: Oct. 21, 1996
Notification: Aug. 28, 1997

EPA Est. _____

Ants (including fire ants) "

Lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed above. For heavy infestations, repeat application after 2 weeks.

Also for control of *cutworms on dichondra and Irish moss*. Use as directed above for lawns.

DIRECTIONS FOR USE OF BIO KILL ON VEGETABLES

For Residential or Home Vegetable Gardens Only

Prepare BIO KILL spray mix as directed in the following chart. Use 0.25 gallons (1 qt.) of finished spray per 110 sq. ft. or 1 liter per 10.7 square meters of surface area.

For Backyard Vegetable Gardens Only

Prepare spray mix as directed in the following chart.

Vegetable	Target Pest(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Sweet corn	Corn earworm, European Corn borer, Fall armyworm, Southern armyworm	Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.	13 oz./gal.
Tomatoes (Florida only). Only for application to tomatoes grown in Florida for varieties yielding tomatoes less than one inch in diameter.	Beet and Southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle, Cabbage looper	Apply for uniform coverage when needed but not more than 6 applications per season. Can be applied up to date of harvest.	13 oz./gal.

Use for Fruit Trees and Nut Crops Ground Around The Home

Apply finished spray to trees until foliage is dripping wet.

Trees (fruit or nut)	Target Pests(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Almonds	Naval orangeworm, Peach twig borer	Apply when insects appear. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest	16 oz./gal.

9 7 11

Trees (fruit or nut)	Target Pests(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Apples	Plum curculio, Redbanded leafroller, Rosy apple aphid, Spotted tentiform leafminer, Tarnished plant bug, White apple leafhopper, Obliquebanded leafroller, Green fruitworm	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.	16 oz./gal.
Peaches	Lesser peachtree borer, Peach twig borer, Plum curculio, Tarnished plant bug, Green fruitworm, Oriental fruit moth	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.	16 oz./gal.
Pears (dormant through delayed dormant)	Pear psylla	Apply during the dormant through delayed dormant growth period only. Add, if desired, 8 oz. of spray oil to each gallon of finished spray solution. Do not make more than 2 applications per season.	16 oz./gal.
Pears (summer)	Pear psylla, Green fruitworm, Codling moth	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.	16 oz./gal.

Vegetable	Target Pest(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Asparagus (preharvest)	Cutworm, Asparagus beetle	Apply for uniform coverage as needed. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.	7 oz./gal.
Asparagus (postharvest)	Larval stages of Asparagus beetle, tarnished plant bug, lygus bugs, and adult Japanese beetle	Treat ferns or bush growth after spear harvest when insects are present.	7 oz./gal.
Celery	Vegetable leafminer, Cabbage looper, Black and Granulate cutworm, Fall and Beet Armyworm	Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.	13 oz./gal.

EPA Reg. No. 64321-1
 MASTER LABEL
 Approved: Oct. 21, 1996
 Notification: Aug. 28, 1997

EPA Est. _____

10 7 19

Vegetable	Target Pest(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Cole Crops: Broccoli, Brussels sprouts, Cabbage, Cauliflower	Beet armyworm, Diamond back moth, Imported cabbageworm, Cabbage-aphid, Cabbage looper	Apply every 5 days or as needed. Can be applied up to one day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels sprouts and broccoli, and 10 applications on cabbage	7 oz./gal.
Eggplant	Colorado potato beetle, Vegetable leafminer	Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.	13 oz./gal.
Horseradish (spring preplant dip)	Imported crucifer weevil	Soak sets for 30 minutes in 0.1 % solution and air dry before planting.	16 oz./qt.
Horseradish (foliar application)	Imported crucifer weevil	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.	13 oz./gal.
Head lettuce only (Do Not use on leaf lettuce)	Cabbage looper, Alfalfa looper, Beet armyworm, Corn earworm and Tobacco budworm	Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.	13 oz./gal.
Potatoes	Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato flea beetle, Aster leafhopper, Potato aphid, Potato tuberworm, Cabbage looper, Cutworms, European corn borer, Potato psyllid, Tarnished plant bug	Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest.	13 oz./gal.
Peppers	Vegetable leafminer, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer	Apply as needed but not more than 8 sprays per season. Can be applied up to 3 days prior to harvest.	13 oz./gal.

Vegetable	Target Pest(s)	Remarks	Recommended Dilution Use Rate Finished Spray
Spinach	Beet armyworm, Corn earworm, Cabbage looper, Cutworms, European corn borer, Green cloverworm, Leafhoppers	Apply for uniform coverage when needed, but not more than 7 applications per season. Can be applied up to 7 days prior to harvest	7 oz./gal.

Storage & Disposal

Do not contaminate water, food or feed by storing or disposal.

Pesticide Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal:

CONTAINERS ONE GALLON AND SMALLER: Do not reuse empty container. Wrap container and put in trash collection.

CONTAINERS LARGER THAN ONE GALLON:

Metal Containers: Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Plastic Containers: Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Buyer assumes all risks of use, storage of handling of this product in strict accordance with directions given herewith.

Distributed by
Bio-Environmental Products
768 Walker Road, Ste. 301
Great Falls, VA 22066