

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF

PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Rebecca M Horton, Agent for Jesmong Holding, Ag. Registration by Design, Inc. 118 ½ East Main Street. Suite 1 Salem, VA 24153—3805

JUN 12 2008

Subject: Notification(s) for Label Revisions under PRN 98-10 and PRN 2007-4

Storage & Disposal and Other Changes (Warranty Statement)

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 98-10 and 2007-4 dated May 14, 2008 for:

EPA Registration 64321-1 "BIOKILL brand INSECTICIDE"

The Registration Division (RD) has conducted a review of the request(s) for applicability under PRN 98-10 and PRN 2007-4 and finds that the label changes requested fall within the scope of PRN-98-10 and PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on non-refillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

rease reed instructions or	n reverse before complet	ting form.		Form Approv	ed. OMB No. 20	070-0060	. Approval exp	ires 2-28-9
≎EPA	اں Environmental	rited States		Registrat	Registration Amendment		Number	
		Application	for Pesticid	le - Section	n i			
1. Company/Product Numb 64321-1				roduct Manager		1 - 1	posed Classifica	stion Restricted
4. Company/Product (Name BIO KILL brand Ins			PM# 13				110110	Nasa seco
5. Name and Address of A Jesmond Holding Ag Chollerstrassee 35, Zug, Switzerland	g	de)	(b)(i), m to: EPA R		imilar or identic		- -	
			Section - I					
Amendment - Explain Resubmission in rec	sponse to Agency letter	dated		Final printed lal Agency letter d "Me Too" Appl Other - Explain	lication.	to		
NOTIFICATION of: (1) Label This notification is consistent 156.146 and 156.156. No oth is a violation of 18 U.S.C. Se PR Notices 2007-4 and 98-10	Explanation: Use additional page(s) if necessary. (For section I and Section II.) NOTIFICATION of: (1) Label Change per PR Notice 2007-4; (2) revised Warranty/Disclaimer per October 2006 Agency guidelines. This notification is consistent with the provisions of PR Notices 2007-4 and 98-10 and the requirements of EPA regulations at 40 CFR 152.46, 156.10, 156.140, 156.144, 156.146 and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. Jesmond Holding Ag understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA, and further understands that if the amended label is not consistent with the requirements of PR Notices 2007-4 and 98-10, and 40 CFR 152.46, 156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and Jesmond Holding Ag may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.							
			Section - II	1				
1. Material This Product W							 -	
Child-Resistant Packaging Yes No	Unit Packaging Yes No If "Yes"	No. per	Water Soluble Po	eckeging No. per	2. Type of C	Metal Plastic Glass Paper		·
* Certification must be submitted	Unit Packaging wgt.	1	Package wgt	container		Other (S	ipecify)	
3. Location of Net Content		4. Size(s) Retail	l Container	5.	Location of Labo	el Directio	ns,	
6. Manner in Which Label	Label Container 1 Qt. on label 6. Manner in Which Label is Affixed to Product Paper glued Stendiled							
			Section - I\	V				
1. Contact Point /Comple	te items directly below f	for identification (of individual to be	e contacted, if n	ecessery, to pro	cess this	application.)	
Name Rebecca M. Horton			itte onsultant/Agen	nt for Jesmono		Telephone 540-375-8	8 No. (Include Ai 8826	rea Code)
i certify that the sta i acknowledge that both under applicable	stements I have made on any knowlingliy false or le law.	Certification this form and all misleading states	ll attachments the	areto are true, a vishable by fine	courate and com or imprisonment	oplete.	6. Date Applica Received (Stampe	
2. Signature	W Hortz	1	. Title Consultant/Age	ent for Jesmor	nd Holding Ag	٠		
4. Typed Name Rebecca M. Horton		5.	Date .6/1.	3/08 lev	ised			



118½ East Main Street Suite 1 Salem, VA 24153-3805 phone 540.375.8826 fax 540.375.8827 CERTIFIED MAIL #7007 0710 0002 1185 5637 May 14, 2008

Document Processing Desk (NOTIF)
Office of Pesticide Programs/Reg. Div. (7505P)
U.S. Environmental Protection Agency
Ariel Rios Bldg.
1200 Pennsylvania Ave., NW
Washington, DC 20460

Re:

NOTIFICATION FOR COMPLIANCE WITH PR NOTICE 2007-4

EPA Reg. No. 64321-1, BIO KILL brand INSECTICIDE

Attn:

Linda Arrington

Notifications & Minor Formulations Team Leader

On behalf of Jesmong Holding, Ag. of Zug, Switzerland, enclosed is notification of administrative label revisions, including revision to the Storage & Disposal text for compliance with PR Notice 2007-4 and revised Warranty/Disclaimer per October 2006 Agency guidelines.

The following documents are enclosed:

1) EPA Form #8570-1, Application for Pesticide - Notification (2 copies)

2) Copy of the proposed label with revisions highlighted in red.

If further action or information is required on Jesmond Holding's behalf, please advise.

Best regards,

Rebecca M. Horton Consultant/Agent for

JESMOND HOLDING AG

e-mail: hortonb@ntelos.net

RH/alb Enclosures



Please read	instructions of	on reverse	before	completing	for
					_

 Registration
Amendment

Please read instructions on	reverse <u>before comp</u>	leting form.		Form A	Dp <u>rove</u>	od. O <u>MB No. 207</u>	70-00 <u>60</u>	. Approval expires 2-28-9
≎EPA	Environmenta	United States	•		V	Registration Amendme Other	on	OPP Identifier Number
		Application	n for Pestic	ide - Sec	ction	1		
1. Company/Product Numbe 64321-1	er .	,, ,	2. EPA LaRo	A Product Mer DCCA	nagar			posed Classification None Restricted
4. Company/Product (Name) BIO KILL brand INSEC			PM# 13					10000
5. Name and Address of App	•	ode) .						FIFRA Section 3(c)(3)
Jesmond Holding Ag Chollerstrassee 35, C Zug, Switzerland	CH 6301		to:			nilar or identical	in con	nposition and labeling
Check if this	s is a new address			uct Name				
`			Section -	11				
Amendment - Explain Resubmission in resp Notification - Explain	ponse to Agency letter	r dated		Final printed Agency lett "Me Too" A Other - Exp	ter dat Applica	ation.		
This notification is consistent with changes have been made to the hamiltully make any false statement 156.156, this product may be in vi	labeling or the Confidential at to EPA, and further under	 Statement of Formularstands that if the amount 	ula for this product. Jo mended labet is not co	Jesmond Holding consistent with the rement action a	g Ag und ne requir	derstands that it is a versents of 40 CFR 15	violation c 56.10, 15	of 18 U.S.C. Sec. 1001 to 6.140, 156.144, 156.146 and
1. Material This Product Will	i Be Packaged in:							
Child-Resistant Packaging Yes No	Unit Packaging Yes V No		Weter Soluble P Yes No	ackaging		V PIE	tainer etal astic ass	
* Certification must be submitted	If "Yes" Unit Packaging wgt.		If "Yes" Package wgt	No. per container	r	· —	per ther (Spe	acify)
3. Location of Nat Contents I	Information Container	4. Size(s) Retail	l Conteiner 1 qt.	.]	5. Loc	cation of Label Di	irections	,
5. Manner in Which Label is A		Lithograp Paper glu Stenciled	ph ued d	Other	r			
		·	Section - I	V				
1. Contact Point (Complete i	items directly below I	for identification c	of individual to b	e contected,	if nece	ssary, to proces:	s this ap	pplication } o
Name Rebecca M. Horton		1	itle Consultant/Ager	nt for Jesma	ond H		phone N)-375-882	No. (Include Ayea Code) 26 ຶ່ວລັ້ນ
I certify that the statem I acknowledge that any both under applicable le	y knowlinglly false or a		li attachments the			mprisonment or	()	Date Application Received () (Stamped)
Liberta M.	. Horton		Title Consultant/Age	ent for Jes	mono		66630	0 0 0000 0000
. Typed Name Rebecca M. Horton		5.1	Date 5/14	1/08				

IFRONT PANEL

BIO KILL® brand INSECTICIDE

[Additional Brand Name: Multipurpose BioKill brand insecticide; Multi-Purpose BugBusters; BioKill brand insecticide Multi-purposel

Optional Statements

For commercial use

Makes up to 16 gallons

For residential use

Water-based Permethrin insecticide

Odorless in application.

For use INDOORS on houseplants and in indoor landscaping

A READY-TO-USE and DILUTABLE Permethrin-based emulsion insecticide

Dilutable for use on plants

For use OUTDOORS (residential site only) in gardens, on ornamental trees, shrubs and flowers, on lawns.

For use outdoors, in homes and non-food areas of food/feed processing establishments, hotels, motels, commercial buildings, schools, warehouses, theaters, office buildings and kennels

Will not stain water-safe fabrics and surfaces

For use indoors and outdoors on a wide variety of crawling and flying insects

Use on DOGS and CATS to control fleas and ticks

Can be used as a DIP for fleas on DOGS

Use on Poultry to control lice and mites

Provides effective broad-spectrum control of building and garden insect pests.

Effective for up to four weeks

Multipurpose, ready-to-use, water based bug killer

Non-flammable, do-it-yourself product

Easy to use - just spray affected areas

Use indoors and outdoors - broad spectrum insecticide includes: cockroaches, mosquitoes, ants, spiders, flies, moths, fleas, lice, bedbugs, crickets, grasshoppers and many, many other listed species No odor - Non-staining - No sticky residue

Odorless and Multi-purpose Insecticide

Long lasting - residual effect up to 4 weeks

Sold in over 30 countries

Ready-to-use, water-based bug killer

Kills crawling insects, including cockroaches, ants, fire ants, fleas, spiders, ticks, lice, silverfish, earwigs, beetles, bedbugs, crickets and more!

Kills flying insects, including flies, moths, mosquitoes, gnats, hornets, wasps, black flies and more! Rx for bugs.

For Bedding & Furniture

NOT FOR USE ON HUMANS .

The Only Lice Control Spray Available in a Non-Aerosol Bottle

Observe all precautionary statements and follow directions for use carefully.

Household insects: cockroaches, fleas, bed bugs, many, many more

Household plants and indoor landscaping

Pet spray for dogs/cats to eliminate fleas, ticks and lice

Lawns & gardens: many listed insects

NOTIFICATION

JUN 1 2 2008

EPA Est.

ACTIVE INGREDIENT:

Permethrin (CAS #52645-53-1) 0.25%

OTHER INGREDIENTS: 99.75%

100.00%

[NOTE TO EPA: Front panel of label may include graphics representing product use; e.g., insects controlled]

Keep Out of Reach of Children CAUTION

See back/side/other panel(s) (& accompanying booklet) for additional precautionary statements

NET	CONTE	NTS	

[BACK/SIDE/OTHER PANEL(S) &/OR BOOKLET, WHEN NECESSARY]

READ ENTIRE LABEL BEFORE EACH USE.

SHAKE BOTTLE WELL BEFORE USING

COMMERCIAL APPLICATIONS:

For use in non-food areas indoors and outside premises of Hotels, Motels, Commercial buildings, Schools, Warehouses, Theaters, Office Buildings, Kennels, Drains, Apartments and Condominium buildings, and non-food contact areas of food-feed processing establishments.

Inside and outside landscaping

Controls bugs inside garbage bins, garbage dumpsters and in garbage chutes

Controls bugs in hallways, common areas and inside apartments, office units and public areas

HOUSEHOLD APPLICATIONS:

House plants and indoor landscape

Inside - along baseboards, in carpets, mattresses, around cupboards, etc.

Outside - in gardens, on ornamental flowers, lawns, vegetables and fruit trees.

DOMESTIC ANIMALS:

Dogs and cats to control fleas & ticks

Poultry to control lice, fleas & ticks

Livestock - protects beef and dairy cattle, goats, sheep and horses from horn flies, face flies, mosquitoes and gnats

[NOTE TO EPA: When marketed for use on dogs or cats, the following precautionary statement will be added to label:

PRECAUTIONARY STATEMENTS

Hazards to Domestic Animals

CAUTION: Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult

EPA Reg. No. 64321-1 May 14, 2008 notification EPA Est.
Page 2 of 10 Pages

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater. This pesticide is toxic to honey bees and other beneficial pollinators exposed to an application. Do not apply when bees are actively visiting blooming plants (vegetables, flowers, fruit/ornamental trees). Do not allow this product to come in direct contact with beehives at anytime.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE IN NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF HOTELS, MOTELS, HOMES, SCHOOLS, PRESCHOOLS, NURSERIES, DAY CAMPS, GREENHOUSES, LATHHOUSES, SEED HOUSES, FOOD HANDLING ESTABLISHMENTS, PACKAGING AND PROCESSING PLANTS, BREWERIES; COMMERCIAL, INDUSTRIAL, OFFICE, INSTITUTIONAL AND STORAGE BUILDINGS; DAIRIES, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, STORES, GROCERY STORES/SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOWER AND FEED MILLS, BAKERIES, WAREHOUSES, PORTABLE TOILETS, FLOOR DRAINS, AND IN HOSPITALS. IN HOSPITALS, PATIENTS SHOULD BE REMOVED PRIOR TO TREATMENT. ALSO FOR USE IN CAMPERS, MOBILE HOMES, BUSES, AUTOMOBILES, TRUCKS, BOXCARS, MARINAS, BOATS AND SHIPS. ALSO FOR USE IN DOG KENNELS AND HORSE STABLES. IN HORSE STABLES, NOT FOR APPLICATION IF HORSES WILL BE USED FOR FOOD.

INDOOR APPLICATIONS:



Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where

foods may be prepared or held. The product may be used in non-food/feed areas of food/feed establishments such as garbage rooms, lavatories, floor drains (to sewer entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling). In the home, cover all food handling surfaces, cover or remove all food and cooking utensils or wash thoroughly after treatment. Not For Use In Federally Inspected Meat and Poultry Plants. Do not apply to classrooms while in use.

Surface Spraying: Provides control of crawling and flying household insect pests for up to 4 weeks. To control ants, boxelder bugs, brown dog ticks, cadelles, centipedes, cheese mites, cluster flies, clover mites, cockroaches, crickets, earwigs, fire ants, flying moths, fruit flies, gnats, grasshoppers, house flies, Lone star ticks, mealworms, millipedes, mosquitoes, palmetto bugs, rice weevils, scorpions, silverfish (firebrats), sowbugs (pillbugs), spider beetles, spiders and waterbugs: Direct the spray into hiding places, cracks and crevices, behind shelves and drawers, inoist areas, openings around pipes and sinks, under refrigerators. Spray directly into and around floor drains. For silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

To control carpet beetles: Hold container 14-18 inches from carpet. Apply spray evenly, covering the entire surface until slightly moist. Avoid wetting or soaking. Spray underside of carpet when possible. Also spray floor and along baseboards, cracks and crevices wherever these insects are seen

EPA Est. ___

or suspected.

To control *bedbugs*: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Allow all sprayed articles to dry thoroughly before use.

To control *fleas* (adult and larvae) and *ticks* (adult and nymphs): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, beds, entire floor and floor covering (e.g., rugs and carpeting) with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. Make a test application to upholstery or drapery fabrics in an inconspicuous place before use. Allow to dry thoroughly before allowing pets to contact treated areas.

To control *clothes moths*: Remove and brush thoroughly any infested articles from storage and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage areas directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly by keeping nozzle at least 3 feet from fabric to avoid possible staining. Allow all sprayed articles to dry thoroughly before use.

To kill *lice* and *louse eggs*: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

Grain Insects: To control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths and carpet beetles remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the area by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed. Any utensils that are contacted with spray should be washed before reusing. Repeat as necessary.



Industrial Warehouse Fogging: For area control of crawling and flying insect pests for up to 4 weeks.

To control spiders, millipedes, centipedes, sowbugs, pillbugs, ants, firebrats, cigarette beetles, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, mealworms, Indian meal moths, coried grain beetles, tobacco beetles, grain mites and cadelles: Close all doors and windows and fog at the rate of 1 ounce per thousand cubic feet, moving toward the exit. Fill room with fog. Direct to all parts of the room especially windows and other light sources which attract these insects. Leave area and keep closed for a period of 2 hours before reentry. Open and ventilate before reoccupying. Fogged area must be cleared of all personnel and pets while fogging is taking place. This product is to be used solely for warehouse space used to store non-food/feed materials.



OUTDOOR APPLICATIONS:

Garden Insects: For use on roses, dahlias, asters and other ornamentals to kilk Japanese beetles as well as whiteflies, aphids, lace bugs, ants, exposed thrips, spiders, red mites and leafminers. May be used on plants in small gardens as a multi-purpose spray and for spot treatments of incipient insect infestations to prevent spreading into larger garden areas.

EPA Est.
Page 4 of 10 Pages

When spraying plants, do not hold nozzle closer than 18 inches. Use a sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperature exceeds 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Nuisance Insect Pests: 1 gallon BIO KILL brand insecticide treats 800 linear feet with a 6-inch band. [16 fl. oz. BIO KILL brand insecticide treats 100 linear feet with a 6-inch band.]

Outside Surfaces of Buildings: To control ants, clover mites, crickets, earwigs, millipedes, sowbugs (pillbugs). To help prevent infestations, spray the building foundation to a height of 2-3 feet where insects are active and may find entrance. Apply as a residual spray to porches, screens, window frames, eaves, patios, garages, refuse dumps and other outside building areas where these pests congregate or are seen. Repeat treatment as needed to maintain effectiveness. To aid in reducing annoyance from cluster flies, flying moths, gnats, house flies and mosquitoes, spray outside surfaces of screens, doors, window frames or wherever these insects may enter. Treat surfaces around light fixtures on porches, in garages and other places where these insects gather. Repeat as necessary.

Other Outdoor Areas: To kill *crickets*, *earwigs* and *fleas* thoroughly spray infested areas of weeds or brushy non-crop areas. For *ants* thoroughly wet hills and runways. Repeat application as reinfestations occur.



Wasps, Hornets, Yellow Jackets: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, including nest openings in ground, bushes, under attic rafters and other cracks and crevices which may harbor pests. Saturate nest openings and contact as many insects as possible. Repeat as necessary.

Fire Ants: Apply 1 gallon of BIO KILL as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter (1.22 m). For best results apply in cool weather, 65°-80° F (18.5-27°C), or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound.

Ticks (including ticks which may carry lyme disease): Apply as a residual treatment to control *Deer tick (Ixodes dammiei)*, *Lone Star tick (Amblyomma americanium)*, *Brown dog tick (Rhipicephalus sanguineus)*, *American dog tick (Dermacentor variabilis)* and *Western black-legged tick (Ixodes pacificus)*. Apply until surface is wet (approximately 1 gal/750-1000 ft.) [(approximately 16 fl. oz./94-125 ft.)] Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter. Treat a band of vegetation and leaf litter 6-10 feet wide around the outside of buildings to help control ticks occurring there. To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.



ON DOGS & CATS:

USE ONLY ON DOGS OR CATS. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

On Dogs. To control *fleas*, ticks and lice: Start spraying at the tail, moving the sprayer rapidly and making sure that the animal's body is covered, including the legs and underbody. While spraying fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat every three weeks. As a dip for dogs: Using one part product and three parts water, make up enough material to thoroughly soak animal.

EPA Est.
Page 5 of 10 Pages

Wet animal with water before treatment. Sponge, dip or pour diluted product on the animal until fur and skin are thoroughly wet, making sure that the animal's entire body is treated, including the legs and underbody. Do not treat animal's face. Do not use on puppies less than 12 weeks old.

On Cats. To control *fleas*: Start spray at the tail, moving the sprayer rapidly and making sure that the animal's body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or on face. Avoid contact with the genitalia. Repeat every 16 days. DO NOT DIP CATS. Do not use on kittens less than 12 weeks old.



ON LIVESTOCK:

To protect beef and dairy cattle, goats, sheep and horses from *horn flies, face flies, mosquitoes and gnats*: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control stable flies, horse flies and deer flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give control protection.

To control **blood-sucking lice**: Apply to infested area of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To aid in control of above listed insects and reduce annoyance to livestock, thoroughly spray barn walls, stalls and surfaces where these pests may alight or congregate.

To control *poultry lice*: Spray roosts, walls and nests or cages thoroughly. Follow by spraying over the bird with a fine mist.

To control *bedbugs and mites in poultry houses*: Spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

To control *flies, mosquitoes* and *gnats* in barns, milk rooms, dairy plants, slaughter and poultry houses: Apply as a fog or fine mist, directing spray toward the ceiling and upper corners of the room until the area is filled with mist, using 1 oz. of Bio Kill® brand insecticide in 2 quarts of water to cover 1,000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for 10 to 15 minutes. Vacate the area and ventilate before re-occupying. Repeat application as necessary.

DIRECTIONS FOR USE AS A DILUTABLE PRODUCT FOR USE ON INDOOR AND OUTDOOR PLANTS:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or lawns and grounds.

Note: Do not use more BIO KILL than recommended, as this may cause some plant injury. Use this dilution unless otherwise specified. Use with care on ferns, palms and fuchsia. As with all plants libred above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars and, therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

CONTACT & RESIDUAL SPRAY - Apply as a thorough coverage spray, wetting leaves and branches to run-off. Use sufficient spray to penetrate dense foliage. Adjust spray volume used to plant size.

EPA Reg. No. 64321-1

May 14, 2008 notification

EPA Est.

Page 6 of 10 Pages



Indoor Application: Mix 8 oz (1 cup) of BIO KILL 0.25% with 1 gallon of water [2 oz. (4 Tbs) of BIO KILL with 1 quart of water]. If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in

an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

To protect plants in indoor landscaping, spray directly on insects to kill *leafrollers*, *armyworms*, *cutworms* and *cabbageworms* on Chrysanthemums; *fungus gnats* on Velvet plants. Spray as directed above to control *aphids*, *armyworms*, *corn earworms*, *fall cankerworms*, *inchworms*, *Japanese beetles*, *leafminers*, *leafrollers*, *mealybugs*, *scale*, *spider mites*, *spring cankerworms*, *exposed thrips* and *whiteflies* on the following: Ageratum, Ardisia, Aster, Azalea, Baby's breath, Begonia, Birdsnest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Crown of thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English ivy, Fuchsia, Gladiolus, Grape ivy, Hypoestes Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-back plant, Poinsettia, Portulaca, Pothos, Prayer plant, Purple passion, Rabbits foot fern, Rose, Schefflera, Snapdragon, Snake plant, Statice, Velvet plant, Verbena, Zinnia.



Outdoor Application: Mix 8 oz (1 cup) of BIO KILL 0.25% with 1 gallon of water 12 oz. (4 Tbs) of BIO KILL with 1 quart of water]. Spray in the late afternoon or evening when temperature ranges from 50°F to 78°F, and when there is little or no wind. Spray at the first sign of insect pests. On those plants that will flower during the season, the spray schedule should be completed before blossom formation. Repeat as necessary; use intervals of 4 to 8 days. BIO KILL application can be made up to the day of collection. Repeat applications may be necessary to control citrus blackflies on non-bearing citrus; adult Japanese beetles on roses.

Roses & Flowers: Spray as directed above to control aphids, cicadas, corn earworms, fall cankerworms, fall webworms, inchworms, leafminers, leafrollers, mealybugs, bagworms, spider mites, exposed thrips and whiteflies on the following: Ageratum, Aster, Azalea, Begonia, Coleus, Common ninebark, Common snowberry, Exacum, Gladiolus, Gold bells, Hypoestes, Ivy, Lilac, Marigold, Mock orange, Nannyberry, Orchid, Pansy, Pea shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Wax plants, Zinnia.

Trees & Shrubs: Use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control aphids, bagworms, cicadas, exposed thrips, fall cankerworms, inchworms, leafminers, leaf rollers, mealybugs, spider mites and whiteflies on the following: Arizona cypress, Birch, Cherry, non-bearing Citrus, Conifers, Elm, Euonymous, English ivy, Fir, Honeysuckle, Oak, Palm, Philodendron, Pine, Taxus, Tulip Poplar and Weeping fig. Use to control pine moths, pine needleminers, pine beetles and needle scales on Pine; budworms on Conifers; oakworms; on Oak; fall webworms on Cherry, Elm and Honeysuckle; citrus black flies on non-bearing citrus; tussock moths on Douglas and grand firs; leafminers on Birch; elm leaf beetle and elm spanworms on Elms. To kill tent caterpillars: Application should be made when caterpillars are young and tents are first noticed. For best results, apply in late afternoon or evening, when caterpillars have returned to their tents. Apply BIO KILL FULL STRENGTH. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die

EPA Est.

Page 7 of 10 Pages

LAWNS: For armyworms mix 4½ oz. in 1 gal. of water and use 10 gals. of diluted BIO KILL solution to cover 1,000 sq. ft. For brown dog ticks, chinch bugs, cutworms, fleas, Japanese beetle grubs, mole crickets and sod webworms mix 1 oz. in 1 gal. of water. Use 10 gals. of diluted BIO KILL solution to cover 1,000 sq. ft. To control ants (including fire ants), crickets and grasshoppers mix 6.4 oz. in 1 gal. of water. Use 3 gals. of diluted BIO KILL solution to cover 1,000 sq. ft.

Thoroughly wet down grass a few hours before applying. Lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed above. For heavy infestations, repeat application after 2 weeks. Also for control of *cutworms on dichondra and Irish moss*. Use as directed above for lawns.

FRUIT & NUT TREES: For use on fruit and nut trees grown around the home. Mix 1 pt. (16 fl. oz.) BIO KILL in 1 gallon of water. Apply when insects appear. Spray until foliage is dripping wet.

Almonds: Naval orangeworm, Peach twig borer. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.

Apples: Plum curculio, Redbanded leafroller, Rosy apple aphid, Spotted tentiform leafminer, Tarnished plant bug, White apple leafhopper, Obliquebanded leafroller, Green fruitworm.

Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.

Peaches: Lesser peachtree borer, Peach twig borer, Plum curculio, Tarnished plant bug, Green fruitworm, Oriental fruit moth. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.

Pears (dormant through delayed dormant): Pear psylla. Apply during the dormant through delayed dormant growth period only. Add, if desired, 8 oz. of spray oil to each gallon of finished spray solution. Do not make more than 2 applications per season.

Pears (summer): *Pear psylla, Green fruitworm, Codling moth.* Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.

VEGETABLES: For Residential or Home Vegetable Gardens Only

Prepare BIO KILL spray mix as directed. Use 0.25 gallons (1 qt.) of finished spray per 110 sq. ft. or 1 liter per 10.7 square meters of surface area.

Asparagus (preharvest): Cutworm, Asparagus beetle - Mix 7 oz. in 1 gal. of water. Apply for uniform coverage as needed. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest. (postharvest): Larval stages of Asparagus beetle, tarnished plant bug, lygus bugs, and adult Japanese beetle. Mix 7 oz. in 1 gal. of water. Treat ferns or bush growth after spear harvest when insects are present.

Celery: Vegetable leafminer, Cabbage looper, Black and Granulate cutworm, Armyworms (fall, beet & southern). Mix 13 oz. in 1 gal. of water. Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.

Cole Crops (Broccoli, Brussels sprouts, Cabbage, Cauliflower): Beet armyworm, Diamond back moth, Imported cabbageworm, Cabbage-aphid, Cabbage looper. Mix 7 oz. in 1 gale of water. Apply every 5 days or as needed. Can be applied up to one day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels sprouts and broccoli, and 10 applications on cabbage.

Eggplant: Colorado potato beetle, Vegetable leafminer. Mix 13 oz. in 1 gal. of water. Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to

harvest.

Horseradish: Imported crucifer weevil. Spring preplant dip - Mix 1 pt. (16 oz.) in 1 qt. of water. Soak sets for 30 minutes and air dry before planting. Foliar application - Mix 13 oz. in 1 gal. of water. Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.

Lettuce (Head lettuce only - do not use on leaf lettuce): Cabbage looper, Alfalfa looper, Beet armyworm, Corn earworm and Tobacco budworm. Mix 13 oz. in 1 gal. of water. Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.

Peppers: Vegetable leafminer, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer. Mix 13 oz. in 1 gal. of water. Apply as needed but not more than 8 sprays per season. Can be applied up to 3 days prior to harvest.

Potatoes: Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato flea beetle, Aster leafhopper, Potato aphid. Potato tuberworm, Cabbage looper, Cutworms, European corn borer, Potato psyllid, Tarnished plant bug. Mix 13 oz. in 1 gal. of water. Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest.

Spinach: Beet armyworm, Corn earworm, Cabbage looper, Cutworms, European corn borer, Green cloverworm, Leafhoppers. Mix 7 oz. in 1 gal. of water. Apply for uniform coverage when needed, but not more than 7 applications per season. Can be applied up to 7 days prior to harvest.

Sweet corn: Corn earworm, European Corn borer, Fall armyworm, Southern armyworm. Mix 13 oz. in 1 gal. of water. Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.

Tomatoes (Florida only. Only for application to tomatoes grown in Florida for varieties yielding tomatoes less than one inch in diameter.): Beet and Southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle, Cabbage looper.

Mix 13 oz. in 1 gal. of water. Apply for uniform coverage when needed but not more than 6 applications per season. Can be applied up to date of harvest.

Storage & Disposal

Do not contaminate water, food or feed by storing or disposal.

Pesticide Storage and Spill Procedures: Keep from freezing. Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

[CONTAINERS ONE GALLON AND SMALLER]:

Pesticide Disposal: If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain

Container Disposal: If empty - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

[CONTAINERS LARGER THAN ONE GALLON]

Pesticide Disposal: Pesticide or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container.

Metal Containers: Triple rinse or equivalent, then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

EPA Est.

Page 9 of 10 Pages

Plastic Containers: Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage of handling of this product not in strict accordance with directions given herewith.

Jesmond Holding Ag, Zug, Switzerland

[Lot no. to be added at production.]

EPA Est.

Page 10 of 10 Pages