


PH 14 64405-2

1043



**REDZONE
BEETLE BAIT**

MOLE CRICKET, ROACH AND BEETLE BAIT

**Weather Resistant Bait for the Control of
Crickets, Mole Crickets and Cockroaches**

**Kills Darkling Beetles and Hide Beetles
and Their Larva Stages**

For Both Interior and Exterior Use

ACTIVE INGREDIENT:

Ortho Boric Acid

5%

INERT INGREDIENTS

95%

TOTAL

100%

Keep Out of Reach of Children

C A U T I O N

See [side] panel for additional precautionary statements.

EPA Reg. No. 64405-2

EPA Est. 64405-TN-1

NET CONTENTS: __ POUNDS

**NISUS CORPORATION
Cherokee Place
101 Concord Street, North
Knoxville, TN 37919**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a ready-to-use, weather resistant bait for the control of crickets, mole crickets and cockroaches including:

CRICKETS: Field and Mole Crickets

COCKROACHES: Asian, American, Brown Banded, Smokey Brown, German and Oriental

BEETLES: Darkling and Hide (and their larva stages)

APPLICATION INSTRUCTIONS

FOR THE CONTROL OF MOLE CRICKETS ON LAWNS, ORNAMENTAL TURF, PLAYING FIELDS, PARKS, FAIRWAYS AND GREENS OF GOLF COURSES: Use a mechanical spreader to apply bait evenly to affected area at the rate of 2 pounds per 1,000 square feet or 90 pounds per acre. Areas to be treated should be well irrigated prior to application and product applied late in the afternoon. For maximum effectiveness, do not irrigate treated area immediately after application.

FOR THE CONTROL OF ASIAN COCKROACHES ON LAWNS, ORNAMENTAL TURF, PLAYING FIELDS, PARKS AND AREAS ADJACENT TO HOUSEHOLDS OR PUBLIC FACILITIES: Use a mechanical spreader to apply bait evenly to lawns and playing fields at a rate of 2 pounds per 1,000 square feet or 90 pounds per acre. Scatter bait evenly in flower beds, leaf litter, in trash cans or refuse areas.

For perimeter treatments, spread bait at a rate of 6 ounces per 100 square feet in a band at least 2 feet wide around foundations, patios, driveways, sidewalks, entrances to homes and buildings or other areas where roaches are present. Apply around landscape lighting, floodlights, streetlamps or other exterior fixtures.

FOR THE CONTROL OF COCKROACHES AND CRICKETS IN AND AROUND HOMES, HOTELS, APARTMENT BUILDINGS, STORES AND RESTAURANTS (NON-FOOD AREAS) WAREHOUSES AND SEWERS: Apply at the rate of 4 pounds per 1,000 square feet (6 ounces per 100 square feet) of surface area. Spread evenly in crawl spaces, attics, drop ceilings, cellars with dirt or gravel floors. In warehouses, garages and basements, concentrate application along walls and baseboards. Apply in inaccessible areas such as cracks and crevices where insects may hide. Bait may also be placed in removable trays. Locate trays in areas accessible to insects but away from children and pets. Always record the location and number of bait stations. In sewers, apply along ledges and around manhole entrances. Treat exterior perimeter areas of buildings in a band at least 2 feet wide and scatter bait in flower beds, leaf litter, wood piles, trash cans and refuse areas.

TO KILL DARKLING BEETLES AND HIDE BEETLES (ADULT AND LARVAE): For Dirt-Floored Grow Out Houses: Apply dry with a fertilizer or seed spreader directly to soil at the rate of two pounds per 100 square feet when house is empty and after litter has been removed. Reapply annually at the rate of one pound per 100 square feet. Prior to the warm season apply at the two pound rate to the surrounding two feet outside the house under eaves. For interior

sidewalls mix at the rate of one pound per three gallons of water. Using a pressure sprayer, saturate cement sidewalls, top plates, posts and framing. Spray mix into cracks and crevices around installation. 150 gallons of mix will cover a 10,000 square foot house. Reapply spray annually or after each time a house has been disinfected. For best results, allow house to remain empty for at least two to four days before replacing litter. Wear a dust mask or a respirator and goggles when applying dust. Do not apply directly to the litter.

TO KILL DARKLING BEETLES AND HIDE BEETLES (ADULT AND LARVAE): For Layer Houses and Cement-Floored Grow Out Houses: Mix at the rate of one pound per three gallons of water. Using a pressure sprayer, saturate sidewalls, top plates, posts and framing. Spray mix into cracks and crevices around insulation. Apply at the rate of one pound product (three gallons of mix) for each 100 square feet of treated surface. 200 to 300 pounds will treat a 30,000 square foot house. Product must be applied after each clean-out and after standard sanitation procedures have been completed. Do not apply directly to manure. **NOTE:** Prior to the warm season apply as a dust to the two feet outside the house under the eaves at the rate of two pounds per 100 square feet.

Storage & Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place. Do not store where children or animals may gain access. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Wash hands thoroughly after handling. May cause eye irritation. Avoid eye contact. Wear a dust mask or respirator.

Statement of Practical Treatment

If Swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious or convulsing person. **If Inhaled:** Remove person to fresh air. Apply artificial respiration if indicated. Contact a physician if warranted. **If On Skin or In Eyes:** Flush with plenty of water. Contact a physician if irritation persists.