

## LONG LASTING

Resistant to rain and watering DEADLINE BULLETS are not easily dissipated by water.

## DIRECTIONS: AGRICULTURAL USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Apply as a soil surface treatment. May be broadcast by air or ground equipment unless specified otherwise in the label. Use a band treatment between the rows after formation of edible parts. Do not contaminate edible portions of the plant. May be applied to the following crops to protect from slugs and snails.

**FIELD CROPS:** Cereal grains, corn, legumes, mint: Apply 10 to 40 lbs. per acre.

**SEED CROPS:** Sugar beets, flowers, grasses, vegetable and legume crops grown for seed: Apply 10 to 40 lbs. per acre.

**VEGETABLES:** Asparagus, beans, beets, black-eyed peas, broccoli, brussel sprouts, cabbage, cantaloupe, carrots, cauliflower, celery, collards, corn, cowpeas, cucumbers, eggplants, endive, garlic, ginseng, horseradish, kale, kohlrabi, leeks, legumes, lettuce, melons, mustard greens, okra, onions, parsnips, peas, peppers, pimentos, potatoes, pumpkins, radishes, rhubarb, rutabagas, salsify, shallots, spinach, squash, sweet potatoes, swiss chard, tomatoes, turnips, watermelons: Apply 20 to 40 lbs. per acre.

**Artichokes:** By air or ground application before tight bud stage, apply 25 to 70 lbs. per acre. After tight bud stage, apply by ground application only.

**TREE CROPS:** Citrus: Apply 20 to 40 lbs. per acre to give even distribution.

Avacado, cherries, peaches, apricots, apples, pears: Apply 20 to 40 lbs per acre.

**SMALL FRUIT:** Blackberries, blueberries, boysenberries, caneberries, loganberries, raspberries, strawberries, grapes: Apply 10 to 40 lbs per acre

**TURF & ORNAMENTALS:** Ornamental greenhouses, ornamental planting, commercial ornamental nurseries, ground covers, turfgrass areas and grasses grown for sod, dichondra, and around home and building foundations: Normally apply 1/2 to 2 lbs per 1,000 sq ft. Repeat treatment as necessary to maintain control.

For best results, apply in evening. Especially beneficial, if applied following rain or watering. Apply every 3 to 4 weeks during growing season, or as needed.

**DO NOT APPLY DIRECTLY TO EDIBLE PORTION OF THE PLANT.**

## STORAGE AND DISPOSAL

**Storage:** Do not contaminate water, food, or feed by storage or disposal. Use product only from original container & keep in locked storage area away from children & pets. Do not store near food, feed, fertilizers, or other pesticides or seeds.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state & local authorities, by burning. If burned, stay out of smoke.



# One last meal for slugs

## ACTIVE INGREDIENT:

Metaldehyde

(2, 4, 6, 8 - Tetramethyl-1, 3, 5, 7 - Tetraox

INERT INGREDIENTS: . . . . .

TOT/

EPA REG. NO. 648

EPA EST. NO. 485

## KEEP OUT OF REACH OF WARNING STATEMENT OF PRACTICAL

**If Swallowed:** Drink 1 or 2 glasses of water and induce vomiting with finger. Do not induce vomiting or give anything to person.

**If In Eyes:** Flush eyes with plenty of water for at least 15 minutes. Attention immediately.

**If On Skin:** Wash thoroughly with soap and water. See side panel for additional precautions.

## PRECAUTIONARY STATEMENT HAZARDS TO HUMANS & DOMESTIC ANIMALS

**WARNING:** Harmful if swallowed. Avoid eye irritation. Avoid contact with skin, eyes. Wash thoroughly with soap and water after handling. Fatal to dogs or other pets if eaten. Keep pets out of treated areas.

**ENVIRONMENTAL HAZARDS:** Do not apply to wetlands (swamps, bogs, marshes and potholes) or surface water when disposing of equipment washwater.

**Deadline™**  
Reg. trademark of  
Pace International LP

**MANUFACTURED BY**  
Pace International LP  
Kirkland, WA 98033  
Made in U.S.A.

# BEST AVAILABLE COPY

NET CONTENTS



- Resistant to rain & watering
- Goes further; Lasts longer

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

If ground is dry, thoroughly water area to be treated. For best results, apply in the evening. Scatter the pellets around flowers, ornamental plants and lawn areas. Do not place pellets in piles. Apply one pound of pellets to each 1,250 sq. ft. Apply only to the soil surface around plants. Do not apply directly to foliage or other plant parts. See below for directions for use around fruits, berries, and vegetables. Always wear protective rubber gloves.

Bait areas likely to be infested at the beginning of the spring planting season before damage occurs. Treat every 3 to 4 weeks during the growing season, or as needed. During periods of high rainfall or frequent waterings, it may be necessary to treat more often.

DEADLINE® BULLETS is specially formulated to attract and control snails and slugs. The highly compressed pellets are easy to use, clean to handle and economical. One pound will bait 1,250 sq. ft.

The presence of snails and slugs can be detected by their shiny mucous-like trail. They are night feeders and prefer damp, cool places in the ornamental garden, flowers, under boards or flower pots or piles of leaves and around fences, rocks, hedges and bushes.

**LESS VISIBLE TO CHILDREN AND PETS**

Because each pellet is compressed and dense, it falls through plant foliage down to the soil surface where it becomes less visible to children or pets, but more available to snails and slugs as they move about in search of food.

**FRUITS, BERRIES AND VEGETABLES:** To control snails and slugs around the following fruits, berries and vegetables, apply to the soil surface around these plants. Do not contaminate edible parts. Thoroughly wash all parts used for food before eating.

**Tree Fruits:** Citrus, avocado, cherries, peaches, apricots, apples, pears.

**Berries:** Strawberries, blackberries, boysenberries, caneberrries, loganberries, raspberries.

**Vegetables:** Artichoke, asparagus, beans, beets, blackeyed peas, broccoli, brussel sprouts, cabbage, cantaloupe, carrots, cauliflower, celery, collards, corn, cowpeas, cucumbers, eggplants, endive, garlic, ginseng, horseradish, kale, kohlrabi, leeks, lettuce, melons, mustard greens, okra, onions, parsnips, peas, peppers, pimientos, potatoes, pumpkins, radishes, rhubarb, rutabagas, salsify, shallots, spinach, squash, sweet potatoes, swiss chard, tomatoes, turnips, watermelons.

**FLOWERS AND ORNAMENTALS:** Apply to soil surface around following plants: azaleas, begonias, camellias, chrysanthemums, daffodils, dahlias, daisies, geraniums, honeysuckle, hydrangeas, ice plants, impatiens, iris, mangolds, pansies, petunias, rhododendrons, roses, snapdragons, sweet peas, tulips.

**NOTE:** When setting out transplants, sprinkle a few pellets around each plant after watering. Those new soft, succulent plants are very tempting to foraging snails and slugs for the first few weeks after planting.

See right side panel for storage and disposal instructions.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

(Iraoxycyclo-octane) . . . . . 4%  
 . . . . . 96%  
 TOTAL . . . . . 100%

54864-2  
 232-WA-1

OF CHILDREN  
 TO  
 CHEMICAL TREATMENT

Prevent vomiting by touching back of  
 throat by mouth to an unconscious

Wait 15 minutes and get medical

Precautionary statements.

PRECAUTIONS  
 DOMESTIC ANIMALS

Avoid breathing dust. May  
 irritate eyes, or clothing. Wash  
 thoroughly. This pesticide may be  
 washed out of treated areas.

Do not apply directly to water or  
 swimming holes. Do not contaminate  
 fish or shellfish.

DATE:

LOT:

**ACCEPTED**  
 MAY 05 1992  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 54864-2

**STORAGE AND DISPOSAL**  
**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Use product only from original container and keep in locked storage area away from children and pets. Do not store near food, food, fertilizers, other pesticides, or fuels. Store in a cool dry place, away from sunlight, in original container.  
**PESTICIDE DISPOSAL:** Product - If a substantially filled container may be re-used, it should be thoroughly cleaned and disposed of in accordance with local, state, and federal regulations.  
 Container - Do not reuse container. Recycle or dispose of in accordance with local, state, and federal regulations.

**BEST AVAILABLE COPY**

**IMPORTANT:** Keep pellets away from children, dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

0492