



READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS ON THE LABEL AND OBSERVE ALL PRECAUTIONS.

Hinder Deer and Rabbit Repellent protects plants by forming an odor barrier that is inoffensive to humans. For use on vegetable, fruit, flower, vine and nursery plants and shrubs.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
DANGER

Causes irreversible eye damage. Harmful if inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist. Wear safety glasses when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product may be hazardous to aquatic invertebrates. Do not spray directly to water, areas where surface water is present, or to wetland areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

See a website at www.fedweb.com for use this product to a summer insecticide with its labeling.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warning must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER" Area to be treated with HINDER on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid recommendations.

USE RESTRICTIONS

This product may be used to limit browsing by white tail deer (*Odocoileus virginianus*) and black tail or mule

(CONTINUED ON RIGHT SIDE PANEL)

LEFFINGWELL® HINDER® DEER AND RABBIT REPELLENT

COMPOSITION

Active Ingredient (% by weight)	13.9%
Ammonium Salts of Higher Fatty Acids	86.1%
Inert Ingredients	100.0%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

PELIGRO

PUEDE CAUSAR DAÑO PERMANENTE A LOS OJOS.

Si usted no lee inglés, no use este producto hasta que alguien se lo explique arrojándole la etiqueta.

STATEMENT OF PRACTICAL TREATMENT/FIRST AID

IF IN EYES: Hold eyelids open and flush with a steady stream of water. Get medical attention. **IF SWALLOWED:** Drink promptly a large quantity of milk, soy milk, gelatin solution, or if those are not available, drink large quantities of water. Avoid alcohol. Get medical attention. Do not induce vomiting. **IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention. **NOTE TO PHYSICIAN:** Possible structural damage may compromise the use of gastric lavage.

In case of an emergency involving this product, call, anytime, 1-800-424-9300

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS IF THESE TERMS ARE NOT ACCEPTABLE. RETURN AT ONCE UNOPENED.

EPA REG. NO. 64884-24 EPA EST. NO. 64884-1M-1

NET CONTENTS: 2½ LIQUID GALLONS (9.45 Liters) • NET WEIGHT: 20.5 LBS (9.25 Kg)

PACE INTERNATIONAL LP • 600 SEVENTH AVENUE SOUTH

KIRKLAND, WASHINGTON 98033 • USA • (252) 627-5711

PACE
INTERNATIONAL

Accept. Stamp on Pg. 2 of 2
Best Available Copy

PM 04

64864-24

11/19/97

P3182

area (L. B. ...), all bare soil, trees and perennial flowers, nursery stock, ...

This product may also be used to ...

When conditions use of this product ...

Do not apply this product through any type of irrigation system.

WINDER IS NOT CHEMICALLY COMPATIBLE with most strongly ...

Check relevant sections of application directions for information on phytoxicity and other specific limitations on treatments.

APPLICATION DIRECTIONS

SHAKE CONTAINER WELL BEFORE MIXING OR APPLYING TO CROPS. WINDER may be applied by conventional spraying equipment ranging from small hand-held sprayers to commercial sprayers.

Under reasonably dry conditions, Winder should be reapplied every 10-14 days for as long as plant production is needed and the treatments appear to be effective.

Winder may reach Winder off to back vegetation. Treatment should be repeated after heavy rains. Do not apply Winder when rain is expected within the next 24 hours.

For best results, apply Winder when the plants are in the early stages of growth. Winder may be applied to plants that are in the early stages of growth. Winder may be applied to plants that are in the early stages of growth.

Consult specific directions (below) for the target species and type of treatment (removed to your control situation).

DEER

Apply and pour Winder: To make spray mixture, mix 3-5 gallons of Winder per 100 gallons of water. Using ground

(CONTINUED ON BACK PANEL)

ing treatment. Treat areas regularly on all lower surfaces up to 6-8 feet above the ground surface. Apply correct spray with air vegetation to be protected has been

Do not apply Winder to plants that are in the early stages of growth. Winder may be applied to plants that are in the early stages of growth.

When conditions use of this product ...

Do not apply this product through any type of irrigation system.

WINDER IS NOT CHEMICALLY COMPATIBLE with most strongly ...

Check relevant sections of application directions for information on phytoxicity and other specific limitations on treatments.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Under reasonably dry conditions, Winder should be reapplied every 10-14 days for as long as plant production is needed and the treatments appear to be effective.

Winder may reach Winder off to back vegetation. Treatment should be repeated after heavy rains. Do not apply Winder when rain is expected within the next 24 hours.

For best results, apply Winder when the plants are in the early stages of growth. Winder may be applied to plants that are in the early stages of growth.

Consult specific directions (below) for the target species and type of treatment (removed to your control situation).

DEER

Apply and pour Winder: To make spray mixture, mix 3-5 gallons of Winder per 100 gallons of water. Using ground

(CONTINUED ON BACK PANEL)

10/19/87, 11:31 AM

NOV 19 1987
let 864-24