

64864-66

12/14/2011

1/9

**U.S. ENVIRONMENTAL PROTECTION AGENCY**

Office of Chemical Safety and Pollution Prevention  
Office of Pesticide Programs  
Registration Division (7504P)  
1200 Pennsylvania Ave., N.W.  
Washington, DC 20460

**NOTICE OF PESTICIDE:**

☒ Registration  
☐ Reregistration  
(under FIFRA, as amended)

EPA Reg. Number:

**64864-66**

Date of Issuance:

**DEC 14 2011**

Term of Issuance:

Conditional

Name of Pesticide Product:

**Shield-Brite TBZ  
500D**

Name and Address of Registrant (include ZIP Code):

**Pace International, LLC**  
**5661 Branch Road**  
**Wapato, WA 98951**

**MAILED TO:**

**Andrew Kieniksman**  
**Supervisor, Regulatory Affairs**

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or can cel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Page 1 of 2

Signature of Approving Official:

Tony Kish, Product Manager (22)  
Fungicide Branch/Registration Division/OPP/OCSPP (7504P)

Date:

**DEC 14 2011**

2. Make the following change to the label:

- Change the product registration number to "EPA Reg. No. 64864-66".

Submit one copy of the revised final printed label with the change listed above before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted" is enclosed for your records.

3/9

**Shield-Brite TBZ 500D**

Thiabendazole Fungicide for Post-harvest Drench and Dip Treatments

**ACTIVE INGREDIENT:**

Thiabendazole: 2-(4-thiazolyl)benzimidazole\* .....42.3%

**OTHER INGREDIENTS:** ..... 57.7%

Total..... 100.0%

\*3.9 lbs. active ingredient per gallon.

EPA Reg. No. 64864-xx

EPA Est. 64864-WA-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.	

**PRECAUTIONARY STATEMENTS****Hazards to Humans and Domestic Animals**

CAUTION. Harmful if swallowed. Causes eye and skin irritation. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after handling.


**Personal Protective Equipment (PPE)**

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes plus socks

In addition to the above, handlers making application to mushroom houses using hand held sprayers must wear chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of waste.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

NOT FOR USE IN FORMULATING AGRICULTURAL PESTICIDES.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Manual seed treatments are prohibited.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

**General Information**

Shield-Brite TBZ 500D is a systemic fungicide which aids in the control of many important plant diseases. If treatment is not effective following use of Shield-Brite TBZ 500D as recommended, a tolerant strain of fungus may be present. Consideration should be given to prompt use of other suitable non-benzimidazole fungicides. Shield-Brite TBZ 500D is not for use in formulating agricultural products.



Crop Use Directions

**GENERAL - SHAKE OR STIR WELL BEFORE USE.**

Clean equipment before using Shield-Brite TBZ 500D. When Shield-Brite TBZ 500D is dispersed in water, the resulting suspension must be constantly agitated. Avoid foaming.

Use the higher recommended rates under conditions of heavier disease pressure for citrus sprays.

Compatibility information for Shield-Brite TBZ 500D with other chemicals is limited. Consult your supplier or agricultural experiment station for specific recommendations.

Do not apply this product through any type of irrigation system.

**CARROTS -**

Gray Mold (*Botrytis cinerea*) and Sclerotinia Rot (*Sclerotinia sclerotiorum*): Dip carrots, before storage, for 5-10 seconds in a suspension containing 43 fl. oz. of Shield-Brite TBZ 500D per 100 gals. of water. Replenish the suspension when the volume is too low or when it becomes dirty.

The runoff and wastes from the dipping operation should not be discarded in a drainage which could contaminate public water systems.

**CITRUS (One Application Only) -**

Blue mold (*Penicillium digitatum*), Green mold (*Penicillium italicum*), and stem end rot (*Diplodia natalensis*): Dip, flood, or spray harvested fruit with a suspension of product in water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion. Rate will vary according to application method; please refer to the following for specific rates for citrus blue mold control.

**Citrus Bin Dips, Floods and Drenches**

Use 27.5 fl. oz. of product in 100 gallons of water to provide 1,000 ppm TBZ. Citrus must be drenched for three (3) minutes. Use one (1) gallon of diluted mixture per 1,500 - 2,000 lbs. citrus or 100 gallons of diluted mixture for about 30 bins. Avoid excessive foaming, use antifoam if needed.

**Citrus Sprays**

Use 54.5 - 95.5 fl. oz. product in 100 gallons of water, wax/oil emulsions, or an aqueous dilution of wax/oil emulsion to provide 2,000 - 3,500 ppm of TBZ. Keep the mixture under constant agitation. Apply at a rate of one gallon per 8,000-10,000 lbs. citrus fruit.

For sporulation control, apply 136.5 fl. oz. of product in 100 gal of water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion to provide 5000 ppm of thiabendazole. Apply at a rate of one gallon of diluted mixture per 8,000 - 10,000 lbs citrus fruit.

**MUSHROOMS (*Agaricus spp.*) -**

Dry Bubble (*Verticillium fungicola*), Wet Bubble (*Mycogone perniciososa*), Cobweb (*Dactylius dendroides*) and Green Mold (*Trichoderma spp.*):

**Addition to mushroom supplements -**

To a finished batch of mushroom supplement, add up to 0.1 % active ingredient (by weight) of Shield-Brite TBZ 500D.

**Applying to mushrooms -**

Apply Shield-Brite TBZ 500D as a drench during normal watering operations. Apply at spawning, casting, and between breaks. Up to 8.4 fl. oz./1,000 sq. ft can be applied at casting while a maximum of 4.2 fl. oz./1,000 sq. ft. can be applied at the other timings. Do not apply more than 21 fl. oz./1,000 sq. ft. of Shield-Brite TBZ 500D per crop unless a supplement containing Shield-Brite TBZ 500D has been used. If the supplement contains thiabendazole, refer to table below for maximum amount that can be applied to a crop. Do not apply within 12 hours prior to harvest.

6/9

Lbs supplement* Per 8,000 sq. ft.	Maximum amount of Shield-Brite TBZ 500D per 1,000 sq. ft. per crop (fl. oz.)
800	18
1200	16
1400	15
1800	13.5
2200	11.5
2400	9.5

\*Assumes at maximum rate of 0.1 % thiabendazole

NOTE: if you are unsure if the mushroom supplement that you are using contains thiabendazole, contact the manufacturer of the mushroom supplement.

#### ORNAMENTAL BULBS AND CORMS –

Fusarium Basal Rot and Penicillium Blue Mold: Clean and treat bulbs or corm within 24-48 hours of digging. Prepare suspension of 30 fl. oz. of Shield-Brite TBZ 500D per 100 gals, of water. Submerge stock completely in dipping suspension. Suspension temperature should be 55-75°F (13-24°C). Discard suspension (1) when it becomes dirty, (2) after using five times, or (3) after 24 hours, whichever occurs first.

Fusarium (Bulbs and Corms): Dip bulbs 15-30 minutes, corms 15 minutes.

Blue Mold (Bulbs): Dip 10-15 minutes.

After treatment, dry bulbs or corms in a shaded, well-ventilated area. Curing or retarding may precede or follow treatment.

#### POTATO –

Fusarium Tuber Rot: Mist unwashed tubers on a conveyor line, with tumbling action, entering storage with 0.44 fl. oz. of Shield-Brite TBZ 500D to each 2,000 Lbs. of tubers in sufficient water for complete coverage. If an additional treatment is necessary before shipping, mist the tubers at the same rate or dip the tubers for 20 seconds in a solution containing 0.44 fl. oz. of Shield-Brite TBZ 500D per gallon of water. Do not treat seed potatoes after cutting.

Shield-Brite TBZ 500D for potatoes should be used as part of an Integrated Management Program, which includes prevention of bruising during harvest and maintaining proper storage temperature and relative humidity. Neither cultural practices nor chemical measures will prevent Fusarium infection if potatoes are damaged or improperly stored.

#### POME FRUIT (Apples and Pears) –

Blue Mold Rot, Bull's Eye Rot, and Gray Mold (Nest Rot, Cluster Rot), Stem End and Neck Rot: Dip, flood, or spray harvested fruit with a suspension of 16.8 fl. oz. of Shield-Brite TBZ 500D in 100 gals, of water. Do not treat for over 3 minutes. Treat apples only before and after storage for maximum decay control. Treat pears once.

#### SWEET POTATO "Seed Roots" –

Black Rot, Scurf, and Foot Rot: Dip the seed roots in a suspension containing 8.4 fl. oz. of Shield-Brite TBZ 500D per 7.5 gals, of water (112.4 fl. oz. per 100 gals.). Treat the seed roots for 1-2 minutes and plant immediately. Discard the suspension when the volume is too low or when it becomes dirty. Do not use the treated roots for food or feed.

## SEED TREATMENT

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Albaugh, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

Seed treatment with Shield-Brite TBZ 500D fungicide will not provide postemergence protection of plants grown in or near infected fields.

### Mixing Procedures

Apply Shield-Brite TBZ 500D as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of Shield-Brite TBZ 500D into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

Federal law requires that bags containing treated seed shall be labeled with the following information:

"This seed has been treated with thiabendazole fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed."

Mixing with other pesticides is limited to adequate physical compatibility and satisfactory phytotox test results.

### SOYBEAN SEED TREATMENT

Shield-Brite TBZ 500D is recommended for protection against pod and stem blight (*Phomopsis spp.*) on soybean seed to improve germination and early season seedling health. For protection against other seed and soil-borne diseases, it is recommended that a co-fungicide such as Apron® XL LS + Maxim® 4FS, or an Apron MAXX® brand fungicide, be used at EPA registered labeled rates and tank mixed in combination with Shield-Brite TBZ 500D. Consult the label for appropriate use rates and follow all use instructions for the products. Utilized as a seed treatment for protection against seed borne and early season seedling pod and stem blight on soybean: Apply Shield-Brite TBZ 500D at the rate of 0.08 to 0.16 fl. oz. per 100 lbs. of seed.

### SEED TREATMENT FOR:

**PEAS** (Dry), **CHICKPEA** (garbanzo beans) (*Cicer arietinum*), **LENTIL** (*Lens esculenta*), **PEAS** (*Pisum spp.*), **PIGEON PEAS** (*Cajanus cajan*)

Shield-Brite TBZ 500D fungicide is used to effectively suppress seed borne *Ascochyta* blight cause by the pathogen *Ascochyta rabiei*. Shield-Brite TBZ 500D seed treatment must also be used to minimize the threat of highly infectious and limit spread of residue-borne and wind-borne infection of chickpeas and lentil crops by *Ascochyta* blight.



**Chickpea Seed Treatment:**

For slurry solutions in water, mix 2.1 fl. oz. of Shield-Brite TBZ 500D fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. It is essential that thorough coverage of seed be obtained.

**Lentil Seed Treatment:**

For slurry solutions in water, mix 1.1 fl. oz. of Shield-Brite TBZ 500D fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation, it is essential that thorough coverage of seed be obtained.

**Peas Seed Treatment:**

For a ready mix or slurry treaters, mix 1.05 fl. oz. of Shield-Brite TBZ 500D fungicide in adequate water to provide complete coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. It is essential that thorough coverage of seed be obtained.

**For Export Use on Peas Seed:**

Shield-Brite TBZ 500D may be mixed with other registered seed treatment products in accordance with the importing Nation's Phytosanitary requirements.

Treated seed must be labeled "for Export Only"- Not to be Sold or Offered for Sale in the United States.

For best results against Ascochyta blight, plant chickpeas, peas or lentil seed treated with Shield-Brite TBZ 500D fungicide as late in the spring as possible.

**RESTRICTIONS AND LIMITATIONS FOR SEED TREATMENTS**

- Maximum one application per season.
- The product is not to be tank mixed with any pesticide unless physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.
- Do not feed vines or hay grown from treated seed to livestock.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** Store in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

*(for 5-gallon containers or smaller)* Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

*(for containers larger than 5 gallons)* Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

*(Continued for all containers)* Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**



**NOTICE TO BUYER**

**IMPORTANT:** Read the information below before using this product. If the terms are not acceptable, you should return the unopened product container immediately for a complete refund.

**LIMITED WARRANTY, TERMS OF SALE, AND LIMITATION OF LIABILITY**

Upon purchase of this product, purchaser and user agree to the following terms:

**Warranty:** To the extent consistent with applicable law, Pace International, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is remarkably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. The Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose; no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

**Terms of Sale:** The Company's directions for use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. All such risks are assumed by the user.

**Limitations of Liability:** To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury, or other claim. Under no circumstances shall the Company be liable for any special, indirect, incidental, or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale, and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Apron®, Apron MAXX®, Maxim®, are trademarks of Syngenta Group Company

MADE IN USA

NET CONTENTS: \_\_\_\_\_

Distributed By:  
Pace International LLC  
5661 Branch Road  
Wapato, WA 98951  
800.936.6750  
www.paceint.com

