U.S. ENVIRONMENTAL PROTECTION Office of Chemical Safety and Pollution Prev Office of Pesticide Programs Registration Division (7504P) 1200 Pennsylvania Ave., N.W. Washington, DC 20460	AGENCY	EPA Reg. Number: 64962-1	Date of Issuance: MAY 0 3 20
1200 Pennsylvania Ave., N.W. Washington, DC 20460		Ferm of Issuance: Unconditional	
NOTICE OF PESTICIDE: <u>X</u> Registration <u>X</u> Reregistration (under FIFRA, as amended)		Name of Pesticide Pr EarthTec	oduct:
Name and Address of Registrant (include ZIP Code): Earth Science Laboratories Inc. 113 SE 22 nd St, Suite 1 Bentonville, AR 72712	Mailed to: c/o William V Pepper Hamil Hamilton Squ 600 Fourteent Washington, 1	ton LLP are	4
Note: Changes in labeling differing in substance from that a must be submitted to and accepted by the Registration Divis correspondence on this product always refer to the above E	sion prior to use	of the label in con	
On the basis of information furnished by the regist hereby registered/reregistered under the Federal Act. Registration is in no way to be construed as a this product by the Agency. In order to protect he Administrator, on his motion, may at any time sus pesticide in accordance with the Act. The acceptan registration of a product under this Act is not to be right to exclusive use of the name or to its use if it EPA received a label amendment request submitte grants this request under the authority of section Fungicide and Rodenticide Act, as amended. With set forth in the Reregistation Eligibility Decision for Therefore, EPA reregisters the product listed abov authority of section 4(g)(2)(c) of the Federal Insec as amended. Reregistration under this section door reassessment of pesticides. EPA may require subr the registration of your product.	Insecticide, F an endorseme alth and the e pend or cance nce of any nan e construed as has been cove ed by email or 3(c)(5) of the this accepted or coppers hav ye. This action cticide, Fungio es not elimina	Sungicide and I nt or recomment, the environment, the el the registrat me in connections giving the reguered by others n April 26, 201 Federal Insection labeling, all re- ve been satisfient is taken under cide, and Roden the the need for	Rodenticide endation of he ion of a on with the gistrant a 2. EPA ticide, equirements ed. r the nticide Act, r continual
Signature of Approving Official:	2	Date:	0 3 2012

1/15

64962-1

Notice of Pesticide Reregistration EarthTec EPA Reg. No. 64962-1 Page 2 of 2

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

The basic and alternate Confidential Statements of Formula (CSF) dated 3/21/2012 submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act are acceptable. These CSFs supersede all CSFs previously accepted for this product.

If you have any questions or comments regarding this letter, please contact Dominic Schuler at (703) 347-0260 or via e-mail at schuler.dominic@epa.gov.

Enclosure:

Label stamped "Accepted" Acute Toxicity Review DP388015 dated 4/30/11 Product Chemistry Review DP394514 dated 10/13/11

MASTER LABEL FOR EARTHTEC EPA REG. NO. 64962-1

ACCEPTED

MAY 0 3 2012

EARTHTEC

Algicide/Bactericide*

For Impounded Waters, Lakes, Ponds, Lagoons, Wastewater Lagoons, Reservoirs, Livestock Watering Systems, Potable Water Supplies⁺, Sedimentation Basins, Ornamental Water Features or Fountains

Algicide/Bactericide*

For Irrigation Conveyance Systems, Irrigation Reservoirs, Irrigation Canals, Ditches, Chemigation Systems, Rice Fields and to Control Tadpole Shrimp in Rice Fields

Algicide/Bactericide^{*} For Aquacultural Ponds

Algicide/ Bactericide^{*} For Residential Pools, Spas and Hot Tubs

Algicide/Bactericide^{*} For Feedlot Run-Off Lagoons, Animal Waste or Confinement Pits and Organic Sludge Pits

Bactericide^{*} - Nonpublic Health Bacteria Potable Water Supplies⁺ - Water Destined to Be Used as Drinking Water

ACTIVE INGREDIENT

1

Copper Sulfate Pentahydrate [*] (CAS No. 7758-99-8)	19.8%
OTHER INGREDIENTS.	80.2%
Total	100.0%
*Metallic Copper	5%

THIS PRODUCT WEIGHS 9.9 LB. PER GALLON.

1.188 kg/L

NET CONTENTS: EIGHT (8) OUNCES THIRTY-TWO (32) OUNCES SIXTY-FOUR (64) OUNCES ONE (1) U.S. GALLON TWO AND ONE-HALF (2.5) U.S. GALLONS (Commercial Use Only) THIRTY (30) U.S. GALLONS FIFTY-FIVE (55) U.S. GALLONS TWO HUNDRED SEVENTY-FIVE (275) U.S. GALLONS

DANGER PELIGRO

KEEP OUT OF REACH OF CHILDREN

If you do not understand this label, find someone to explain it to you in detail. (Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.)

FIRST AID

IF IN EYES:	Hold eye open and rinse slowly and gently with water for 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for advice.
	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR	
CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15 to 20 minutes. Call a poison control center or doctor for treatment.
NOTE TO	
PHYSICIAN:	Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact INFOTRAC 1-800-535-5053 for emergency medical treatment.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

MFG. BY:

Earth Science Laboratories, Inc. 113 SE 22nd Street, Suite 1 Bentonville, AR 72712 Phone: 800-257-9283

EPA REGISTRATION NO.: EPA ESTABLISHMENT NO.: BATCH NO.: 64962-1 64962-NE-001

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield or safety glasses), long sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material (Chemical Resistance Category A). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Waters treated with this product may be hazardous to aquatic organisms. Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead algae and weeds. This oxygen loss can cause fish and invertebrate suffocation. To minimize this hazard, do not treat more than ½ of the water body to avoid depletion of oxygen due to decaying vegetation. Wait at least 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow fish to move into untreated areas. Consult with the state or local agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required.

Certain water conditions including low pH (≤ 6.5), low dissolved organic carbon (DOC) levels (3.0 mg/L or lower) and "soft" waters (i.e. alkalinity less than 50 mg/L) increases the potential acute toxicity to non-target aquatic organisms. The application rates on this label are appropriate for water with alkalinity greater than 50 mg/L. Do not use these application rates for water with less than 50 ppm alkalinity (e.g., soft or acid waters) because trout and other species of fish may be killed under such conditions.

Consult your local state fish and game agency before applying this product to public waters. Permits may be required before treating such waters.

For applications in waters destined for use as drinking water, those waters must receive additional and separate potable water treatment. Do not apply more than 1.0 ppm as metallic copper in these waters.

PERSONAL PROTECTIVE EQUIPMENT

USER SAFETY REQUIREMENTS

Mixers, loaders, applicators and other handlers must wear the following:

- Long-sleeved shirt
- Long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material (Chemical Resistance Category A)
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.
- Wash the outside of gloves before removing.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your state and tribe, consult the agency responsible for pesticide regulation. For use in residential pools, spas and hot tubs, do not apply this product in a way that will contact adults, children or pets.

AGRICULTURAL USE REQUIREMENTS APPLICABLE TO USE ON RICE FIELDS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, and nurseries, and mixers, loaders, applicators, and other handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted re-entry interval, and notification to workers. Do not enter or allow worker entry into treated areas during the restricted entry interval of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as soil or water. Wear coveralls, protective eyewear, chemical resistant gloves (i.e. gloves made of any waterproof material) and shoes plus socks.

USE INFORMATION

EarthTec is a highly soluble formulation. EarthTec is used to control algae and to suppress nonpublic health bacteria and bacteria that cause taste and odor problems in impounded waters, lakes, ponds, lagoons, wastewater lagoons, reservoirs, livestock watering systems, potable water supplies⁺, sedimentation basins, and ornamental water features or fountains.

EarthTec is used to control algae and to suppress nonpublic health bacteria and bacteria that cause taste and odor problems in irrigation conveyance systems, irrigation reservoirs, irrigation canals, ditches, chemigation systems, rice fields and to control tadpole shrimp in rice fields.

EarthTec is used to control algae and to suppress nonpublic health bacteria and bacteria that cause odor problems in aquacultural ponds.

EarthTec is used to control algae and to suppress nonpublic health bacteria and bacteria that cause odor problems in residential pools, spas and hot tubs.

EarthTec is used to suppress nonpublic health bacteria and bacteria that cause odors (such as odors from hydrogen sulfide and ammonia gas) in feedlot run-off lagoons, animal waste or confinement pits and organic sludge pits.

USE IN CONTROL OF ALGAE, NONPUBLIC HEALTH BACTERIA, AND BACTERIA THAT CAUSE ODOR PROBLEMS

For algae control, apply in the late spring or early summer when algae first appear. The dosages are variable and depend upon algae species, water hardness, water temperature, amount of algae present, as well as whether water is clear, turbid, flowing or static. Preferably, the water should be clear with temperature above 60 degrees F (15.6 degrees C). Higher dosages are required at lower water temperatures, higher algae concentrations and for hard waters. See Specific

Directions for Use. Application should be done by pouring a measured amount of EarthTec directly from a container into the lakes, ponds, reservoirs or irrigation canals. EarthTec is very soluble and will quickly disperse. Several application points speed up dispersal. Static water requires less chemical for algae control than does flowing water. Use higher dosages for chara, nitella and filamentous algae (pond scum) and lower dosages for planktonic algae. If there is uncertainty about the dosage, begin with a lower dosage and increase until control is achieved or until the maximum allowable level has been reached. See Specific Directions for Use.

Treatment of algae can result in oxygen loss from the decomposition of dead algae. This loss can cause fish suffocation. If the algae cover more than $\frac{1}{3}$ of the total water area, treat in sections. Treat $\frac{1}{2}$ of the water area in a single operation and wait for 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow fish to move into untreated areas. In regions where ponds freeze in winter, treatment should be done 6 to 8 weeks before expected freeze to prevent masses of decaying algae under an ice cover. Before treating bodies of water, consult proper state authorities such as the fisheries commission or conservation department to obtain any necessary permits. For use in controlling algae and cyanobacteria at all aquatic application sites do not exceed a copper concentration in water of 1.0 ppm of metallic copper concentration.

For example, if you wish to achieve 1.0 ppm of metallic copper, 1 gallon of EarthTec added to 60,000 gallons of water is equal to 1.0 ppm metallic copper. In order to attain 1.0 ppm of metallic copper in the treated water, the amount of EarthTec added to a water body is equal to the gallons of water being treated divided by 60,000 multiplied by 1 (e.g., see Gallons of EarthTec and Water table below). Use volumetric measurement devices that are calibrated in accordance with manufacturer specifications.

Gallons of EarthTec and Water						
Gallons EarthTec	Gallons Water	Metallic Copper ppm				
0.1 (0.4 quarts or .8 pints)	6,000	1.0				
¹ / ₄ (1 quart)	15,000	1.0				
1	60,000	1.0				
1 2/3	100,000	1.0				

Use formula for calculating water volume and flow rates. Calculate the volume of water (multiply the average depth by surface area). To calculate the gallons of water multiply the volume in cubic feet times 7.5. One cubic foot per second of flow equals 27,000 gallons/hour. One acre foot equals 326,000 gallons. See below for additional directions on methods of application to flowing water.

SPECIFIC DIRECTIONS FOR USE

To Control Algae, Nonpublic Health Bacteria, and Bacteria That Cause Odor Problems in Irrigation Reservoirs, Impounded Waters, Lakes, Ponds, Lagoons, Reservoirs, Livestock Watering Systems, Potable Water Supplies⁺, Sedimentation Basins, and Ornamental Water **Features or Fountains:** For fish-bearing lakes, ponds, drinking water reservoirs, irrigation canals and other applications, apply at the rate of 1 quart of EarthTec per 250,000 gallons of water, or 1 gallon of EarthTec per 1,000,000 gallons of water for preventive treatment of algae and nonpublic health bacteria. This will yield a concentration of 0.06 ppm metallic copper.

If algae are present, treat at the rate of 3 quarts of EarthTec per 250,000 gallons of water, or 3 gallons of EarthTec per 1,000,000 gallons of water. This will yield a concentration of 0.18 ppm metallic copper.

For applications without fish or for wastewater lagoons apply at the rate of up to 1 quart of EarthTec per 15,000 gallons of water, or 1 gallon of EarthTec per 60,000 gallons of water. This will yield a rate of 1.0 ppm metallic copper.

Do not exceed 1 gallon of EarthTec per 60,000 gallons of water (1.0 ppm metallic copper) under any circumstances for water destined for use as drinking water. EarthTec may be poured into the water manually after calculating the volume of water to be treated and measuring the quantity EarthTec necessary to attain a concentration of 0.06 ppm or by using an automated dispenser calibrated to release the required amount. For best results disperse EarthTec evenly throughout the body of water on a sunny day when algae are near the surface. Do not apply copper sulfate to water with less than 50 ppm alkalinity.

To Control Algae and Tadpole Shrimp in Rice Fields: Apply any time the tadpole shrimp appears from planting time until the seedings are well rooted and have emerged through the water or at the first sign of algae growth on the surface of the field. Applications are most effective when made prior to algae leaving the soil surface and rising to the water surface and prior to appearance of the tadpole shrimp. Factors such as water depth, temperature, pH and the amount of algae can affect the amount of EarthTec needed to control algae and tadpole shrimp. If the depth of water is 8 inches, apply 9 gallons of EarthTec per acre. If the depth of water is 4 inches, apply 4½ gallons of EarthTec per acre. EarthTec can be metered into the rice field as water is being applied. Do not exceed a copper concentration in water of 2.5 ppm of metallic copper concentration (9 gallons of EarthTec per acre with 8 inch depth of water or 13½ gallons of EarthTec per acre foot of water).

To Control Algae or Nonpublic Health Bacteria and Bacteria That Cause Odor Problems in Open Channel Irrigation Conveyance Systems and Chemigation Systems, Ditches, Canals and Similar Irrigation Conveyances: To prevent algae growth using a static application method, apply 1 gallon of EarthTec to 1,000,000 gallons of water to yield a rate of .06 ppm metallic copper in the water. If algae are present, apply 16.6 gallons of EarthTec to 1,000,000 gallons of water to yield 1.0 ppm metallic copper. To prevent algae growth using continuous flow systems, a metered flow rate of 1 milliliter per minute is added to a pumping flow of 267 gallons per minute to yield a rate of .06 ppm metallic copper. If algae are present, do not exceed the total dose of 1 gallon of EarthTec in 60,000 gallons of water (1.0 ppm metallic copper). See Example Calculation table below for continuous flow rates. To Control Algae or Nonpublic Health Bacteria and Bacteria That Cause Odor Problems in Sprinkler, Drip or Other Types of Irrigation Equipment: Agitation is not required. Do not mix with basic substances. EarthTec must be applied continuously for the duration of the water application. To prevent growth of algae, nonpublic health bacteria, and bacteria that cause odor problems, treat at a rate of 1 gallon EarthTec per 60,000 gallons of water to 1 gallon EarthTec per 1,000,000 gallons of water. This will yield a rate of 1.0 ppm to 0.06 ppm metallic copper (see Example Calculation table below). If algae are visible, start by cleaning the pipes or lines and then applying 1 gallon of EarthTec in 60,000 gallons of water (1.0 ppm metallic copper). See Example Calculation table below for continuous flow rates. Once the lines are cleaned, use the preventive dose described above.

EXAMPLE CALCULATION CHEMIGATION AND IRRIGATION FLOW RATES (0.06 ppm Cu)

Water Flow Rate gpm	Water Flow Rate cfm	Dosage Rate ppm Metallic Cu	EarthTec fl oz/min	Feeder Pump Setting EarthTec mL/min
3,000	400	0.06	0.4	11.3
6,000	800	0.06	0.8	22.6
9,000	1,200	0.06	1.1	34.0
12,000	1,600	0.06	1.5	45.3

CHEMIGATION AND IRRIGATION FLOW RATES (1.0 ppm Cu)

Water Flow Rate gpm	Water Flow Rate cfm	Dosage Rate ppm Metallic Cu	EarthTec fl oz/min	Feeder Pump Setting EarthTec mL/min
3,000	400	1.0	6.4	188.7
6,000	800	1.0	12.8	377.5
9,000	1,200	1.0	19.1	566.2
12,000	1,600	1.0	25.5	755.0

To Control Algae and Nonpublic Health Bacteria and Bacteria That Cause Odor Problems in Aquacultural Ponds: Apply at the rate of ¹/₄ to ¹/₂ gallon of EarthTec per acre foot (326,000 gallons) of water to yield concentrations ranging from 0.05 ppm to .09 ppm metallic copper, respectively. Metallic copper concentration is directly proportional to amount of EarthTec added per acre foot. A maintenance dose of 4 to 8 ounces per acre foot may be used every 14 days. The rate is dependent on water temperature, fish density and the degree of suppression targeted.

Amount	for Aquacultura of EarthTec Ap Foot (12 Inches	plied
Gallons EarthTec	Gallons Water	Copper ppm
0.25	326,000	0.05
0.5	326,000	0.09

To Control Algae and Nonpublic Health Bacteria, and Bacteria That Cause Odor Problems in Residential Swimming Pools, Spas and Hot Tubs:

For pools, apply at the rate of 2 fluid ounces of EarthTec per 1,000 gallons of water (20 fluid ounces per 10,000 gallons). Use a properly calibrated measurement device to determine the amount of EarthTec to be added.

For spas and hot tubs, apply at the rate of 6 mL of EarthTec per 100 gallons of water using supplied measuring cup.

Do not use a measurement cup or device that may also be used for food.

These dosages will yield a rate of 0.9 ppm metallic copper. Treated pool effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Application should be made before visible algae appear.

Every 14 days, test the copper level using a standard commercial swimming pool copper test kit. Add EarthTec to raise level back to 0.9 ppm (see tables). The amount of EarthTec to be added is proportional to the starting concentration and volume of water. Do not exceed 1.0 ppm metallic copper.

Pool Volume (gallons)	7,000	8,000	9,000	10,000	11,000	12,000	13,000	14,000
Measured Metallic Copper Level		ADDIT	IONAL	FLUID O	UNCES	OF EAR	ТНТЕС	
in Pool		A	DDED 7	TO MAIN	NTAIN C	ONTRO	L	in the
0.9 ppm	0	0	0	0	0	0	0	0
0.8 ppm	1	1	2	2	2	2	2	3
0.7 ppm	3	3	4	4	5	5	5	6
0.6 ppm	4	5	5	6	6	7	8	9
0.5 ppm	6	7	7	8	10	10	11	12
0.4 ppm	8	9	10	11	13	13	14	15
0.3 ppm	9	10	12	13	15	15	16	18

MAINTAINING METALLIC COPPER CONCENTRATION IN POOLS

0.2 ppm	11	12	14	15	17		18	20	21
0.1 ppm	12	14	15	17	19		20	22	24
MAINTAIN	ING MET	FALLIC	COPPI	ER CON	NCENT	RATIO	ON IN	SPAS	
Spa Volume (gallons)	100	200	300	400	500	700	800	900	1,000
Measured Metallic Copper Level in Spa			TIONA					THTEC DL	
0.9 ppm	0	0	0	0	0	0	0	0	0
0.8 ppm	1	1	2	3	3	4	5	6	6
0.7 ppm	1	3	4	5	6	9	10	11	13
0.6 ppm	2	4	6	8	9	13	15	17	19
0.5 ppm	3	5	8	10	13	18	20	23	25
0.4 ppm	3	6	9	13	16	22	25	28	32
0.3 ppm	4	8	11	15	19	27	30	34	38
0.2 ppm	4	9	13	18	22	31	35	40	44
0.1 ppm	5	10	15	20	25	35	40	45	51

An alternate method is to apply at the rate of 1 fluid ounce of EarthTec per 1,000 gallons of water. This will yield a rate of 0.45 ppm metallic copper. Repeat a maintenance dosage of 1 fluid ounce of EarthTec per 1,000 gallons of water once a month to maintain control. Application should be made before visible algae appear. Where visible algae are present apply at a rate of 2 fluid ounces of EarthTec per 1,000 gallons of water. This will yield a rate of 0.9 ppm metallic copper.

BACTERIAL ODOR CONTROL

To Control Bacterial Odor in Feedlot Run-Off Lagoons, Animal Waste or Confinement Pits, and Organic Sludge Pits: Apply by pouring product directly from the container into the pit or lagoon. Several application points speed up dispersal. Use 1 gallon of full strength EarthTec in 60,000 gallons (8,000 cubic feet) of sewage. This will yield a rate of 1.0 ppm metallic copper. Bacteria and odors should be noticeably reduced in 1 to 2 weeks. Repeat application when odor reoccurs. Minimum re-treatment interval is 14 days.

Feedlot Run-Off Lagoons: Add a portion of the required dosage of EarthTec at several locations around the lagoon to speed dispersal of the product. A minimum of 2 applications per year (spring and fall) is recommended. Additional applications may be required as needed when the lagoon is pumped.

Animal Waste or Confinement Pits: If pits are located under the confinement buildings, add EarthTec directly to these pits. If the pits are outside, add product to the transfer line to the pit.

APPLICATION AND HANDLING EQUIPMENT

Application, handling or storage equipment MUST consist of fiberglass, PVC, polypropylene, viton, corrosion resistant plastics or stainless steel. Never use mild steel, nylon, brass or copper around EarthTec. Always rinse and clean equipment thoroughly each night with plenty of fresh, clean water.

PESTICIDE STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a safe place away from pets and keep out of the reach of children. Store away from excessive heat. EarthTec will freeze. Always store EarthTec above 32 degrees F (Do Not Freeze). Freezing may cause product separation.



Always keep container closed. Store EarthTec in its original container only. Keep away from galvanized pipe, and any nylon storage or handling equipment.

DISPOSAL

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess EarthTec mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance. In the event of spill, neutralize with limestone or baking soda before disposal. May deteriorate concrete.

CONTAINER HANDLING:

Containers with capacities less than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times. Offer for recycling if available. If recycling is not available, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Containers with capacities greater than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¹/₄ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least 1 complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure 2 more times. Offer for recycling if available. If recycling is not available, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Containers too large to shake:

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure 2 more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



IMPORTANT READ BEFORE USING LIMITED WARRANTY AND LIMITATION OF REMEDIES

Read the entire Directions for Use, Limited Warranty and Limitation of Remedies (including limitations on liability) before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

The Directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product.

Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Earth Science Laboratories, Inc. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

To the extent consistent with applicable law, seller warrants that the product conforms to the chemical description and is reasonably fit for the purpose stated on the label for use under normal conditions, but makes no other warranties of FITNESS OR MERCHANTABILITY expressed or implied, or any other warranty if the product is used contrary to the label instructions, or under conditions not foreseeable to the seller. To the extent consistent with applicable law, the seller shall not be liable for more than the cost of this product to the buyer and will in no event be liable for any consequential, special or indirect damages connected with the use or handing of this product. This product is offered and the buyer or user accepts it subject to the foregoing terms which may not be varied. Seller makes no warranty for product which has been frozen.

Manufactured by:

Earth Science Laboratories, Inc. 113 SE 22nd Street, Suite 1 Bentonville, AR 72712 Phone: 800-257-9283