

PM 13

65168-1

172

ACCEPTED

MAY 2 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 65168-1

**MARCA
ABEJA
(BEE BRAND)
CHAMPU
LICE TREATMENT SHAMPOO**

**TRATAMIENTO PARA PIOJOS
PRECAUCION
MANTENGASE
FUERA DEL
ALCANCE
DE LOS NIÑOS**

**No to
Rasques
Natales
con
ABEJA**

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

READ WARNINGS BEFORE USING.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Kills head lice, pubic (crab) lice, and body lice. Apply to affected area until all the hair is thoroughly wet with product. Allow product to remain on area for 10 minutes but no longer. Add sufficient warm water to form a lather and shampoo as usual. Rinse thoroughly. A fine toothed comb or a special lice/nit removing comb may be used to help remove dead lice or their eggs (nits) from hair. A second treatment must be done in 7 to 10 days to kill any newly hatched lice.

ACTIVE INGREDIENTS

Technical Piperonyl Butoxide	4.0 %
Pyrethrum Extract	.33 %
INERT INGREDIENTS:	95.67 %

CAUTION: For external use only. Harmful if swallowed. Do not use near the eyes or permit contact with mucous membranes such as inside the nose, mouth, or vagina as irritation may occur. Keep out of eyes when rinsing hair. Adults and children: Close eyes tightly and do not open eyes until product is rinsed out. Also, protect children's eyes with washcloth, towel, or other suitable material, or by a similar method. If product gets into the eyes, immediately flush with water. If skin irritation or infection is present or develops, discontinue use and consult a doctor. Consult a doctor if infestation of eyebrows or eyelashes occurs. Avoid contamination of feed or foodstuffs. Use with caution on persons allergic to ragweed. **PHYSICAL OR CHEMICAL HAZARDS:** Do not use near heat or open flame.

(See back of label for additional warnings and directions).

12 FL. OZ. -- 354 ML.

**PACKED
Luchett
Bayamon, PR 00960
EPA Est --- 45168-PR-01
EPA Reg**

INC.

INDICATIONS: For the treatment of head, pubic (crab), and body lice and their eggs.

ADDITIONAL DIRECTIONS FOR USE: When Marca Abeja (BEE BRAND) Shampoo is applied, all clothing must be sterilized in order to prevent reinfestation with body lice.

Head Lice: Head lice live on the scalp and lay small white eggs (nits) on the hair shaft close to the scalp. The nits are most easily found on the nape of the neck or behind the ears. All personal headgear, scarfs, coats, and bed linen should be disinfected by machine washing in hot water and drying, using the hot cycle of a dryer for at least 20 minutes. Personal articles of clothing or bedding that cannot be washed may be drycleaned, sealed in a plastic bag for a period of about 2 weeks, or sprayed with a product specifically designed for this purpose. Personal combs and brushes may be disinfected by soaking in hot water (above 130° F) for 5 to 10 minutes. Thorough vacuuming of rooms inhabited by infected patients is recommended.

Pubic (crab) Lice: Pubic lice may be transmitted by sexual contact; therefore, sexual partners should be treated simultaneously to avoid reinfestation. The lice are very small and look almost like brown or gray dots on the skin. Pubic lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface. In hairy individuals, pubic lice may be present on the short hairs of the thighs and trunk, underarms, and occasionally on the beard and mustache. Underwear should be disinfected by machine washing in hot water; then drying, using the hot cycle for at least 20 minutes.

Body Lice: Body lice and their eggs are generally found in the seams of clothing, particularly in the waistline and armpit areas. They move to the skin to feed, then return to the seams of the clothing where they lay their eggs. Clothing worn and not laundered before treatment should be disinfected by the same procedure as described for head lice, except that sealing clothing in a plastic bag is not recommended for body lice because their nits (eggs) from these lice can remain dormant for a period of up to 30 days.

Statement of Practical Treatment

IF SWALLOWED: Call a doctor or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: If skin irritation or infection is present or develops, discontinue use and consult a physician.

STORAGE AND DISPOSAL. Store product in original container and keep in locked storage area. Do not reuse container. Rinse container thoroughly, securely wrap in several layers of newspaper, and discard in trash.

BEST AVAILABLE COPY