

CHLORO GLASS SANITIZER

Sanitizer for use on Bar, Restaurant Glassware, Utensils and Food Contact Surfaces. For Use on: Tabletops, Countertops, Sinks, Splashbacks, Trays, sealed Floors and Food Preparation Equipment. Surface composition may be: Stainless Steel, Glass, Fiberglass, Formica, glazed Porcelain, glazed Ceramic Tile or other Hard Nonporous Surfaces.

Active Ingredient: Trichloromelamine....18.7%
Inert Ingredients: .....81.3%
TOTAL.....100%

Provides 100 PPM Available Chlorine

KEEP OUT OF REACH OF CHILDREN
DANGER: See Precautions On Back Panel
NET WT. 1/4 OZ.

EPA REG. NO. 65169-1 EPA EST. NO. 65169-KY-1
© 2006 H&S Chemical Co. Inc., Covington, KY 41017
U.S.A. Distributor Made in the U.S.A.

ACCEPTED
APR 20 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 65169-1

Front of packet

PRECAUTIONARY STATEMENTS

Back of packet

HAZARD TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive to eyes. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Do not breathe dust. Wash hands thoroughly with soap and water after handling.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

In case of emergency call 800-255-3924 (Chem-Tel). Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LAST TANK 1.) Pre-Wash glassware. 2.) Rinse with clear, cool water. 3.) In sanitizing tank (last tank) mix contents of this package in 3 gallons of water. This provides 100ppm available chlorine for an 8 hour shift. Use test strips to assure required chlorine levels. Immerse glassware for at least two minutes or for contact time specified by governing sanitary code. 4.) Place sanitized glassware on a rack or drainboard to air dry. Do not towel dry.

FOOD CONTACT SURFACES: 1.) Remove heavy soil from all surfaces by a prescrape, preflush or presoak treatment. Then prewash all surfaces with a detergent solution followed by a rinse with potable water. 2.) Fill sanitizing container with 3 gallons of water then add entire contents of packet to water. 3.) Sanitize by wiping or swabbing surfaces. Allow solution to remain on surfaces for two minutes or for contact time specified by governing sanitary code. Then air dry. 4.) Store cloths, sponges or mops in sanitizing solution between uses during each 8-hour shift. Clean and rinse cloths, sponges or mops frequently in sanitizing solution. 5.) Always prepare a fresh sanitizing solution and use the same day prepared. Periodically determine chlorine level with test strip. Change solution when testing indicates a level below federal, state, or local regulations.

STORAGE AND DISPOSAL: Store this product in a cool dry area, away from direct sunlight and heat. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site. Dispose of packet in trash.