

65402-6

07/24/2012

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

JUL 24 2012

OFFICE OF
PREVENTION PESTICIDES
AND TOXIC SUBSTANCES

Ms LuAnn Maloney
Regulatory Manager for,
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Subject VigorOx[®] Citrus XA
EPA Registration Number 65402-6
Your Amendment Dated July 10, 2012
EPA Received Date July 10, 2012

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, FIFRA, as amended, to revise the Directions for Use for the "Surface Disinfections" section as per California Department of Pesticide Regulation's (CDPR) request of July 5, 2012, is acceptable

A stamped copy of the labeling is enclosed

If you have questions concerning this letter, please contact Karen M Leavy at (703)-308-6237

Sincerely,

A handwritten signature in black ink, which appears to read "M Swindell", is written over the typed name.

Marshall Swindell
Product Manager 33
Regulatory Management Branch I
Antimicrobials Division (7510P)

MASTER LABEL

VigorOx® Citrus XA

EPA Registration No 65402-6
EPA Est No 00279-NY 003

VigorOx® Citrus XA is for institutional/industrial sanitizing of previously cleaned non porous food contact surfaces in

- Dairies Wineries Breweries and Beverage Plants
- Meat and Poultry Processing/Packaging Plants
- Milk and Dairy Products Processing/Packaging Plants
- Seafood and Produce Processing/Packaging Plants
- Food Processing/Packing Plants
- Egg Processing/Packing Equipment Surfaces
- Eating Establishments

For Organic Production **VigorOx® Citrus XA** may be used in rinse or wash water on products labeled as organic in food processing facilities on commodities that will be further processed For use as a sanitizer on food contact surfaces in contact with products labeled as organic

VigorOx® Citrus XA is for institutional/industrial sanitizing of previously cleaned hard non porous food contact surfaces such as

- Eating Drinking and Food Preparation Utensils
- Countertops and Food Preparation Surfaces
- Tableware
- Plastic Glass and Metal Bottles (rinse)

VigorOx® Citrus XA is for sanitizing surfaces such as packinghouse conveyors harvesting equipment and containers It is effective against plant pathogens such as *Xanthomonas campestris* (axonopodis) pathovars citrumelo (citrus canker surrogate)

VigorOx® Citrus XA is for sanitizing of hard inanimate non food contact surfaces (general environmental surfaces)

VigorOx® Citrus XA is for use in the disinfection of hard non porous surfaces in general commercial and medical environments such as

- Hospitals Health Care Facilities Veterinary Hospital/Clinics Animal Life Science Labs Pharmaceutical Facilities and Equipment
- Schools Colleges Recreational Facilities Office Buildings
- Livestock Premises Poultry Premises Poultry Hatcheries Animal Housing Facilities
- Retail and Wholesale Establishments
- Bathroom Premises

VigorOx® Citrus XA is for use as a coarse spray for surfaces to be sanitized or disinfected

VigorOx® Citrus XA is for use in fogging applications as an adjunct to acceptable manual cleaning and disinfecting of room surfaces

VigorOx® Citrus XA can be used with the HRS™ and HRS™ II foaming agent For food contact applications the foaming agent must be used in compliance with applicable regulations under the Federal Food Drug and Cosmetic Act

VigorOx® Citrus XA can be used with the non foaming agent Peradigm™ as an antimicrobial container rinse and for hard non porous surface disinfection

VigorOx® Citrus XA is for use as an antimicrobial container rinse to control beverage spoilage microorganisms

VigorOx® Citrus XA is for use in the sanitization of ultra filtration and reverse osmosis (RO) membranes and their associated distribution systems

**KEEP OUT OF REACH OF CHILDREN
DANGER**

See side panel for additional precautionary statements

For Industrial Use Only

Active Ingredients	Peroxyacetic Acid	5 1%
	Hydrogen Peroxide	21 7%
Inert Ingredients		73 2%
Total		100 0%

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER – Corrosive Causes irreversible eye damage and skin burns Harmful if swallowed May be fatal if absorbed through skin Do not get in eyes on skin or on clothing Wear protective eyewear (goggles face shield or safety glasses) clothing and chemical resistant gloves Wash thoroughly with soap and water after handling and before eating drinking or using tobacco Remove contaminated clothing and wash before reuse

Physical or Chemical Hazards – Strong oxidizing agent Mix only with water VigorOx® Citrus XA is not combustible however at temperatures exceeding 156 F decomposition occurs releasing oxygen The oxygen released could initiate or promote combustion of other materials

Environmental Hazards – This pesticide is toxic to birds mammals fish and aquatic life Do not discharge effluent containing this product into lakes streams ponds estuaries oceans or other waters unless in accordance with requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority For guidance contact your State Water Board or Regional Office of the EPA Any solution released from the system should be diluted with water and tested for residuals to ensure that there is less than 3 ppm peroxygen remaining

ACCEPTED
with COMMENTS
in EPA Letter Dated:
JUL 24 2012

FMC Corporation

Peroxygens Division

1735 Market Street

Philadelphia Pennsylvania 19103

Under the Federal Insecticide
Fungicide and Rodenticide Act as
amended for the pesticide
registered under EPA Reg. No. 65402-6

Net Contents 55 Gallons

Pounds

MASTER LABEL

FMC and VigorOx® are trademarks of FMC Corporation



FIRST AID	
If in Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
If on Skin or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
If Inhaled	<ul style="list-style-type: none"> Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
If Swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor for further treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
Note to Physician Probably mucosal damage may contraindicate the use of gastric lavage	
This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body either into or in contact with the bloodstream or normally sterile areas of the body or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body This product may be used to pre-clean or decontaminate critical or semi critical medical devices prior to sterilization or high level disinfection	

ACCEPTED
with COMMENTS
in EPA Letter Dated.

JUL 24 2012

Under the Federal Insecticide
Fungicide and Rodenticide Act as
amended for the pesticide
registered under EPA Reg No

65402-16

Storage and Disposal

Storage

NEVER RETURN VigorOx® Citrus XA TO THE ORIGINAL CONTAINER AFTER IT HAS BEEN REMOVED Avoid all contaminants especially dirt caustic reducing agents and metals Contamination and impurities will reduce shelf life and can induce decomposition In case of a decomposition isolate container douse container with cool water and dilute with large volumes of water

Avoid damage to containers Keep container closed at all times when not in use Keep container out of direct sunlight To maintain product quality store at temperatures below 86 F Do not store on wooden pallets

Procedure for Leak or Spill

Stop leak if this can be done without risk Shut off ignition sources no flames smoking flares or spark producing tools Keep combustible and organic materials away Flush spilled material with large quantities of water Undiluted material should not enter confined spaces

Pesticide Disposal

If material has been spilled an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all local state and Federal environmental laws rules regulations standards and other requirements Because acceptable methods of disposal may vary by location regulatory agencies should be contacted prior to disposal

VigorOx® Citrus XA which is to be discarded should be disposed of as hazardous waste after contacting the appropriate local state or Federal agency to determine proper procedures

Container Disposal

Nonrefillable containers less than 5 gallons Nonrefillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

Nonrefillable containers greater than or equal to 5 gallons Nonrefillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container ¼ full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Empty drums are not returnable to FMC unless special arrangements have been made Triple rinse drums with water Dispose of drums in accordance with local state and Federal regulations

All Refillable containers Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Return to FMC for reuse

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For use in circulation cleaning and institutional/industrial sanitizing of previously cleaned hard non porous food contact surfaces and equipment such as food preparation surfaces pipelines tanks vats fillers evaporators pasteurizers and aseptic equipment in

- Dairies Wineries Breweries and Beverage Plants
- Meat and Poultry Processing/Packaging Plants
- Milk and Dairy Products Processing/Packing Plants
- Seafood and Produce Processing/Packing Plants
- Food Processing/Packing Plants
- Egg Processing/Packing Equipment Surfaces
- Eating Establishments
- Final Sanitizing Bottle Rinse

Foam Sanitization

VigorOx® Citrus XA can be applied as a foam for sanitization of previously cleaned hard non porous food contact surfaces and general environmental (non food contact) hard non porous surfaces such as floors walls ceilings drains and boots Foam applications can be used where penetration and retention of product for required times is difficult to achieve Examples include operating conveyor belts and vertical or uneven surfaces

- 1 Prepare a dilute VigorOx® Citrus XA solution by adding 1 to 17 fluid ounces per 4.5 gallons potable water Best results are achieved using water at 70 F or above
- 2 After preparing the VigorOx® Citrus XA diluted solution add 1 to 10 fluid ounces of HRS™ or add 2 to 20 fluid ounces of HRS™ II per the 4.5 gallons of diluted solution After the HRS™ or HRS™ II is added adjust volume to 5 gallons HRS™ and HRS™ II are only approved foam generating additives for use with VigorOx® Citrus XA
- 3 Apply the sanitizing solution as a foam using commercially available foam generating equipment Allow foam to contact surfaces at least one (1) minute For foot bath application allow foam to remain on the boot surface for one (1) minute upon exiting the bath
- 4 Drain items and/or surfaces thoroughly

Sanitizing Non Porous Food Contact Surfaces

VigorOx® Citrus XA is an effective sanitizer against *Staphylococcus aureus* *Escherichia coli* and *Salmonella typhimurium* Clean equipment immediately after use

- 1 Remove gross particulate matter with a warm water flush
- 2 Wash equipment with detergent or cleaning solution
- 3 Rinse equipment with potable water
- 4 Prepare VigorOx® Citrus XA solution by adding 1.0 to 1.7 fluid ounces to 5 gallons potable water This provides 87.7 to 149 ppm peroxyacetic acid and 373 to 635 ppm hydrogen peroxide
- 5 Fill closed systems with diluted sanitizer solution and allow a contact time of one (1) minute If sanitizing at temperatures of 5 C (40 F) or lower use at least 1.6 fluid ounces of product to 5 gallons of potable water
- 6 If sanitizing against *Listeria monocytogenes* use 1.2 to 1.7 fluid ounces of this product to 5 gallons of potable water This will provide 105 to 149 ppm of peroxyacetic acid and 448 to 635 ppm of hydrogen peroxide
- 7 For open or not completely closed systems use a coarse spray mop/wipe or flood technique to apply the solution to the surface for at least one (1) minute
- 8 Allow surfaces to drain thoroughly before resuming operation

Eating Establishment Sanitizing

VigorOx® Citrus XA is an effective sanitizer against *Staphylococcus aureus* *Escherichia coli* and *Salmonella typhimurium*

- 1 Scrape/prewash plates utensils cups glasses etc whenever possible
- 2 Wash all items with a detergent
- 3 Rinse thoroughly with potable water
- 4 Prepare VigorOx® Citrus XA solution as follows
Add 1.0 to 1.7 fluid ounces of the product to 5 gallons of potable water This will provide 87.7 to 149 ppm peroxyacetic acid and 373 to 635 ppm hydrogen peroxide

- 5 Immerse all items for at least two (2) minutes or for a contact time as specified by the local governing sanitizing code
- 6 If sanitizing against *Listeria monocytogenes* use at least 1.2 to 1.7 fluid ounces of this product to 5 gallons of potable water This will provide 105 to 149 ppm of peroxyacetic acid and 448 to 635 ppm of hydrogen peroxide
- 7 Place all sanitized items on a rack or drainboard to drain adequately air dry if items will not be re used immediately

Sanitizing Tableware

For sanitizing tableware in low to ambient temperature warewashing machines inject the diluted VigorOx® Citrus XA solution (1.0 to 1.7 fluid ounces of the product to 5 gallons of potable water) into the final rinse water Allow treated materials to air dry

Final Sanitizing Bottle Rinse

VigorOx® Citrus XA may be used as a final sanitizing rinse for plastic glass or metal returnable and non returnable bottles / cans

- 1 Wash bottles with detergent or cleaning solution and rinse with potable water
- 2 Rinse bottles/cans with a solution prepared by mixing 1.0 to 1.7 fluid ounces of VigorOx® Citrus XA to 5 gallons of potable water
- 3 Allow to drain adequately

Sanitization of Hatching Eggs

- 1 Prepare a dilute solution by adding 1.0 to 1.7 fluid ounces of VigorOx® Citrus XA to 5 gallons of potable water
- 2 Apply dilute solution as eggs are gathered or prior to setting as a coarse spray or flood so as to lightly wet all egg shell surfaces
- 3 Allow to drain dry

Sanitization of Conveyors Peelers Slicers and Saws for Meat Poultry Seafood Fruits and Vegetables

VigorOx® Citrus XA is an effective sanitizer against *Staphylococcus aureus* *Escherichia coli* and *Salmonella typhimurium*

For use in the static or continuous washing rinsing and sanitizing of conveyor equipment peelers collators slicers saws etc

- 1 Remove all products from equipment if during treatment the sanitizer will directly contact the items
- 2 Prepare VigorOx® Citrus XA solution by adding 1.0 to 1.7 fluid ounces to 5 gallons of potable water
- 3 Apply sanitizer solution to the return portion of the conveyor or to the equipment using a coarse spray or other means of wetting the surfaces Treat for at least one (1) minute Control the volume of solution so as to permit maximum drainage and to prevent puddles The conveyor may still be damp when food contact occurs
- 4 If sanitizing against *Listeria monocytogenes* use at least 1.2 to 1.7 fluid ounces of this product to 5 gallons of potable water
- 5 Allow equipment to drain adequately before reusing a dry surface is not required

General Environmental Surfaces Sanitization (Non Food Contact)

VigorOx® Citrus XA is an effective inanimate hard non food contact surface sanitizer against *Staphylococcus aureus* *Klebsiella pneumoniae* and *Saccharomyces cerevisiae*

Sanitization of surfaces such as floors walls tables chairs benches drains etc can be accomplished using the following procedures

- 1 Remove gross filth with a cleaner or other suitable detergent
- 2 Add 1 to 11 fluid ounces of VigorOx® Citrus XA to 16 gallons of potable water to prepare a solution containing 27 to 302 ppm of peroxyacetic acid and 117 to 1285 ppm of hydrogen peroxide
- 3 Soak items in/with diluted solution using mop/wipe coarse spray or flood techniques and allow contact for at least 5 minutes
- 4 Allow items and/or surfaces to drain adequately or air dry

Surfaces treated to control the spread of Citrus Canker

VigorOx® Citrus XA can be used to control the spread of citrus canker between inanimate surfaces and inanimate surfaces to plants This product is for use on inanimate surfaces such as packinghouse conveyors and harvesting equipment. This product is not for treatment of infected plants

Packinghouse Sanitization

VigorOx® Citrus XA is an effective sanitizer against microorganisms such as *Xanthomonas campestris* (axonopodis) pathovars citrumelo (citrus canker surrogate) and *Staphylococcus aureus* *Escherichia coli* and *Salmonella typhimurium*

- 1 Remove gross contamination with a cleaner or other suitable detergent and rinse with potable water
- 2 Use VigorOx® Citrus XA at a dilution of 9.7 to 15.5 fl oz per 50 gallons of water (85 to 135.8 ppm peroxyacetic acid and 360.5 to 577.7 ppm hydrogen peroxide) as a general sanitizing coarse spray to reduce bacterial and fungal contamination of walls floors conveyers and harvesting containers
- 3 Allow sanitizer to contact surface for at least one (1) minute
- 4 Allow to air dry do not rinse

Field Equipment Sanitization

VigorOx® Citrus XA may be used to sanitize harvest equipment such as pickers trailers trucks (including truck body parts and tires) bins packing crates ladders power tools hand tools gloves rubber boots pruning shears or other equipment that may transfer *Xanthomonas campestris* (axonopodis) pathovars citrumelo (citrus canker surrogate) This product can also be used to sanitize surfaces contaminated with *E. coli* *Salmonella typhimurium* and *S. aureus*

- 1 Remove gross contamination with a cleaner or other suitable detergent and rinse with water
- 2 Use VigorOx® Citrus XA at a dilution of 9.7 to 15.5 fl oz per 50 gallons of water (85 to 135.8 ppm peroxyacetic acid and 360.5 to 577.7 ppm hydrogen peroxide) as a general sanitizing coarse spray
- 3 Allow sanitizer to contact surface for at least one (1) minute
- 4 Allow to air dry Do not rinse

Fogging

VigorOx® Citrus XA is for sanitizing hard non porous room surfaces as an adjunct to acceptable manual cleaning and disinfecting of room surfaces

- 1 Prior to fogging remove or carefully protect all food products and packaging materials
- 2 Ensure room is properly ventilated Vacate all personnel from the room during fogging and for a minimum of 2 hours after fogging Ensure there is no strong odor characteristic of acetic acid before having personnel return to work area
- 3 Fog areas using one quart per 1000 cu ft of room area with a 0.30% (3.5 fl oz per 10 gallons of water) VigorOx® Citrus XA solution
- 4 Allow surfaces to drain thoroughly before operations are resumed

Surface Disinfection

VigorOx® Citrus XA is an effective disinfectant against vegetative forms of Gram positive and Gram negative bacteria and viruses This product is effective against *Staphylococcus aureus* *Salmonella choleraesuis* *Pseudomonas aeruginosa* Influenza A Virus (H1N1 H3N2 and H5N1 strains) Influenza B virus and Parainfluenza Virus Type 3 It may also be used to disinfect veterinary clinic surfaces and livestock equipment contaminated with Newcastle Disease virus Avian Reovirus Avian Infectious Bronchitis Infectious Bursal Disease and Infectious Bovine Rhinotracheitis and may be used in general commercial and medical and veterinary environments to clean disinfect and deodorize hard non porous inanimate surfaces including

- Floors walls and other non porous surfaces such as tables chairs counter tops garbage cans/bins bathroom fixtures sinks bed frames shelves racks carts refrigerators coolers glazed tile linoleum vinyl sealed asphalt glazed porcelain plastic (such as polypropylene and polyethylene) stainless steel or glass
- Hospitals surgical and obstetrical suites operating tables housekeeping services physical therapy departments nursing homes health care facilities autopsy facilities pharmaceutical and chemical processing facilities and equipment
- Schools colleges industrial facilities dietary areas office buildings recreational facilities retail and wholesale establishments
- Animal hospitals veterinary clinics animal life science laboratories kennels kennel runs cages feeding and watering equipment pet shops zoos pet animal quarters poultry premises trucks hatcheries and live stock quarters

To disinfect surfaces that may be contaminated with Gram positive or Gram negative bacteria including *S. aureus* *S. choleraesuis* or *P. aeruginosa*

- 1 Prepare VigorOx® Citrus XA disinfecting solution by adding 3.2 to 3.3 oz of the product to 5 gallons of potable water This will provide 280 to 2630 ppm peroxyacetic acid and 1195 to 11 200 ppm hydrogen peroxide
- 2 Remove gross filth from surfaces to be disinfected by cleaning with a detergent or suitable cleaning product Rinse with clean water
- 3 Apply VigorOx® Citrus XA solution by wiping mopping or as a coarse spray Allow to soak for at least 10 minutes then air dry drain treated surfaces (Applications on food contact surfaces require a sterile or potable water rinse following disinfection)

For surfaces contaminated with the viruses listed above

- 1 Prepare VigorOx® Citrus XA solution by adding 2 2/3 fluid ounce to 5 gallons of potable water This will provide 230 ppm peroxyacetic acid and 990 ppm hydrogen peroxide This product is effective against viruses in up to 500 ppm hard water and on surfaces with moderate organic soil
- 2 Remove gross filth from surfaces by cleaning with a detergent or suitable cleaning product Rinse with clean water
- 3 Apply VigorOx® Citrus XA solution by wiping mopping or as a coarse spray Allow to soak for at least 5 minutes then air dry

VigorOx® Citrus XA may be mixed with the non foaming agent Paradigm™ prior to disinfection

- 1 Prepare a diluted solution of VigorOx® Citrus XA by adding a minimum of 2.8 fl oz to 4.5 gallons of water
- 2 Add a maximum of 43 fl oz of Paradigm to the dilute solution and bring total volume to 5 gallons
- 3 Apply solution as previously described allowing a minimum contact time of 10 minutes

Antimicrobial Rinse of Pre Cleaned or New Returnable or Non Returnable Containers

To reduce the number of non pathogenic beverage spoilage organisms such as *Aspergillus versicolor* *Byssoschlamys fulva* *Pediococcus damnosus* *Lactobacillus buchnei* and *Saccharomyces cerevisiae*

- 1 Prepare VigorOx® Citrus XA by adding 7.0 to 30 fluid oz of product per 5 gallons of potable water This provides 614 to 2630 ppm peroxyacetic acid and 2614 to 11 200 ppm hydrogen peroxide
- 2 Apply solution allowing a minimum contact time of 5 seconds
- 3 Allow containers to drain thoroughly and then rinse with sterile or potable water

VigorOx® Citrus XA may be mixed with the non foaming agent Paradigm™ and applied at room temperature or at a minimum of 25 °C

- 1 Mix a minimum of 1.5 fl oz VigorOx® Citrus XA with Paradigm in 4.5 gallons of water
- 2 Add a maximum of 43 fl oz of Paradigm to the dilute solution and bring total volume to 5 gallons This provides 132 ppm peroxyacetic acid and 560 ppm hydrogen peroxide
- 3 Apply solution allowing a minimum contact time of 30 seconds

Disinfection and Deodorizing of Animal Housing Facilities Poultry Premises Coops, Trucks and Crates

- 1 Remove all animals / poultry from the facilities / items / areas to be disinfected
 - 2 Remove gross particulate litter droppings etc with a warm water flush or by sweeping
 - 3 Empty all troughs racks and other feeding and watering equipment
 - 4 Wash all items thoroughly with detergent or cleaning solution and rinse with potable water
 - 5 Prepare a disinfecting solution by adding 3.2 fluid ounces of VigorOx® Citrus XA to 5 gallons of potable water This provides 280 ppm of peroxyacetic acid and 1195 ppm of hydrogen peroxide which will disinfect surfaces contaminated with Gram positive and Gram negative bacteria as well as poultry and cattle viruses listed above Before instituting the treatment ensure that the work area is well ventilated
- For disinfection saturate surfaces with the diluted product for a period of at least 10 minutes

ACCEPTED
with COMMENTS
in EPA Letter to EPA
JUL 24 2012
Under the Federal Insecticide
Fungicide and Rodenticide Act as
amended for the pesticide
registered under EPA Reg No 65402-4

- 8 For surfaces contaminated with the viruses listed above under *Surface Disinfection* saturate surfaces with diluted product for a period of at least 5 minutes
- 9 Thoroughly scrub treated feed equipment (i.e. feed racks troughs fountains etc.) with a detergent and rinse with potable water
- 10 Do not return animals / poultry or use equipment until solution has been completely absorbed or air dried

Poultry Hatchery Disinfection

- 1 Remove remaining eggs and chicks and all gross particulate and other hatching related debris
- 2 Thoroughly wash all surfaces with a recommended detergent or cleaning solution and then rinse with potable water
- 3 Prepare the disinfecting solution by adding 3.2 fluid ounces of VigorOx® Citrus XA to 5 gallons of potable water
- 4 Before starting the treatment ensure that the work area / room and any closed spaces are well ventilated
- 5 Apply the disinfecting solution with a mop, cloth, brush or coarse spray, keeping surfaces wet for 10 minutes
- 6 Air dry before re-introducing eggs

Batch Sanitization of Ultra Filtration and Reverse Osmosis (RO) Membranes

VigorOx® Citrus XA is for use in the sanitization of ultra filtration medical and non medical institutional/industrial reverse osmosis (RO) membranes and their associated distribution systems. VigorOx® Citrus XA is not for use in kidney dialysis reprocessing equipment.

This product has been shown to be an effective disinfectant when tested by AOAC and EPA methods. This product may not totally eliminate all vegetative microorganisms in reverse osmosis membranes and their associated piping systems due to their construction and/or assembly but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. Check with equipment manufacturer for membrane compatibility with VigorOx® Citrus XA.

- 1 Remove biological or organic fouling from the membrane or other parts of the system with an appropriate cleaner
- 2 Flush the system with RO permeate or similar quality water
- 3 Remove mineral deposits with an acidic cleaner prior to sanitizing the membranes
- 4 Flush the system with RO permeate or similar quality water
- 5 Prepare an appropriate volume of 1% solution of the product (12 fluid ounces per 10 gallons of water). This will provide 526 ppm of peroxyacetic acid and 2338 ppm hydrogen peroxide
- 6 Fill the entire water circuit to be sanitized with the dilute solution and allow the solution to reach a minimum of 20°C (69°F)
- 7 Recirculate the dilute solution of VigorOx® Citrus XA for a minimum of 10 minutes
- 8 Allow membrane elements to soak in the solution for a minimum of 20 minutes
- 9 Rinse the RO system and test for residuals to ensure that there is less than 3 ppm per oxygen. Residuals can be reduced by diverting product water to drain.

Batch Sanitization of Piping Systems Associated with RO Membranes

- 1 Isolate incompatible equipment from piping system. This includes activated carbon filters and ion exchange equipment. Turn off power to ultraviolet light units.
- 2 Estimate total volume of water contained in the system (tanks, rinse stations and piping). Prepare an appropriate volume of 1.0 to 1.5% VigorOx® Citrus XA solution by adding 1.0 to 1.5 gallons of the product for every 100 gallons of solution prepared. Use RO permeate or similar quality water for dilution. This will provide 561 to 842 ppm peroxyacetic acid and 2389 to 3584 ppm hydrogen peroxide.
- 3 Recirculate the dilute VigorOx® Citrus XA solution through the system for a minimum of 4 hours. Process usage valves should be opened and closed to expose internals to the VigorOx® Citrus XA solution.
- 4 Completely drain the system of dilute VigorOx® Citrus XA solution. Thoroughly rinse the system by filling with RO permeate or similar quality water and recirculate before drainage. Repeat the process until test for residuals indicates there is less than 3 ppm peroxygen.

Continuous/Intermittent Addition to Minimize the Accumulation of Biological Matter Between Intermittent Sanitizing Episodes in Piping Systems Associated with RO Membranes

- 1 VigorOx® Citrus XA as received or diluted may be added continuously to the feed water stream between system sanitizing episodes to aid in minimizing the regrowth/accumulation of biological matter. The peroxygen residual in the system which will be effective will vary with the design and usage characteristics of the system. Adjust the addition rate of VigorOx® Citrus XA or the solution and periodically monitor residual peroxygen so that the desired effect is obtained.
- 2 For continuous addition do not exceed 20 ppm VigorOx® Citrus XA (0.1 fl. oz. per 40 gallons of water). This will give 1 ppm peroxyacetic acid and 4.5 ppm hydrogen peroxide. For intermittent feed do not exceed 2000 ppm VigorOx® Citrus XA (10 fl. oz. per 40 gallons of water). This will give 123 ppm peroxyacetic acid and 523 ppm hydrogen peroxide.

Note: Before using VigorOx® Citrus XA to sanitize metal surfaces, it is recommended that the diluted solution be tested on a small area to determine compatibility.

In all applications, always prepare a new sanitizing/disinfecting solution daily to ensure effectiveness. Do not reuse sanitizing/disinfecting solutions. Dispose of any unused sanitizing/disinfecting solution.

EMERGENCY TELEPHONE NUMBERS

(24 HOURS)

MEDICAL: COLLECT 303-595-9048

TRANSPORTATION: 800-424-9300

OTHER: COLLECT 716-879-0400

ESL112310

Rev 072612

ACCEPTED
with COMMENTS
in EPA Letter Dated.

JUL 24 2012
Under the Federal Insecticide
Fungicide and Rodenticide Act
amended for the pesticide
registered under EPA Reg No. 65402-4

4850 5757 2616 v 4