

74 19 65564-1
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
65564-1

Until Reregistration

JMS Stylet-Oil

JMS Flower Farms, Incorporated
1105 25th Avenue
Vero Beach, FL 32960

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) and FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Add the phrase "EPA Registration No. 65564-1" to your label before you release the product for shipment.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

BEST AVAILABLE COPY

63362:1:CR-30:Andreasen:C.Disk:KEVRIC:02/07/92/rj:dd:rj

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

U.S. GPO 1984-416-812

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting. Call a physician. Avoid breathing vapors and spray mists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

STORAGE AND DISPOSAL

STORAGE --- Keep drum tightly closed in storage to prevent entry of water. Do not freeze. Store drum on side. Roll drum before use if stored over-season.

DISPOSAL --- **Prohibitions** Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide Disposal Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

General Consult Federal, State or Local disposal authorities for approved alternative procedures.

Method of Application

Spray at no less than 400 psi pressure using ceramic spray nozzles (ALBUZ[®] ATR, lilac color or their equivalent). Do not apply this product through any type of irrigation system (chemigation).

APPLICATION RATE

Apply 2.0 to 3.0 gallons of diluted spray emulsion per acre depending on pest and crop. If additional coverage is necessary, a tractor speed of 3.5-4 mph is suggested. Vary the quantity of spray applied by changing the number of nozzles used; not by changing either tractor speed or spray pressure.

Timing of Applications

Sprays should be initiated as soon as plants emerge. Plants should be sprayed weekly or twice weekly depending on level of infestation. Continue sprays through harvest.

JMS
Flower Farms, Inc.

**JMS Stylet-Oil[®]
AS AN INSECTICIDE**

ACTIVE INGREDIENT:

BY WEIGHT

ALIPHATIC PETROLEUM DISTILLATE 97.1%

INERT INGREDIENTS 2.9%

TOTAL: 100 %

FEB 18 1992

Unsulfonated Residue of The Oil 99.1%

Flash Point 145°F

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 65564-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Do not induce vomiting. Call a physician.

IF INHALED - Avoid breathing of vapors and spray mist.

IF ON SKIN - Wash immediately with soap and water.

IF IN EYES - Wash immediately with copious amounts of water.

TERMS

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. All applicable directions, restrictions, and precautions on the label are to be followed. This labeling must be in the possession of the user at the time of the application of this material.

Manufactured by - Asgrow Florida Co., Plant City, FL 33566

EPA Establishment No. - 14775-FL-01

Manufactured for - 24(c) Registrant: J.N. Simons,

JMS FLOWER FARMS, INC., 1105 25th Avenue,

Vero Beach, FL 32960 (407) 567-9241

EPA Registration No. - SLN FL 770027

55 GALLONS NET • U.S. STANDARD

BEST AVAILABLE COPY

FIELD RE-ENTRY SAFETY

Do not enter treated areas without appropriate protective clothing.

PRE-HARVEST INTERVAL

No interval required.

ADDITIONAL COMMENTS

Do not spray on wet foliage or if temperature is below 50°F.

COMPATIBILITY INFO

CHEMICAL COMPATIBILITY - Add the following chemicals to the JMS Stylet-Oil[®] as the last ingredient. JMS Stylet-Oil[®] is compatible with: acephate (Orthene), benomyl (Benlate), (CCN), copper hydroxide (Kocide), mancozeb, maneb, methamidophos (M), and triadimefon (Bayleton).

DO NOT USE the following chemicals within two weeks have elapsed between the application of JMS Stylet-Oil[®] and the application of: chemical anilazine (Dyrene), chlorobenzilate (Dikar), dinocap (Karathane), Folpet, permethrin (Ambush, Pounce) and others.

DO NOT USE JMS Stylet-Oil[®] in combination with nutrient spray materials (Nutri-Leaf) or other materials can be sprayed separately. JMS Stylet-Oil[®] is compatible with urea (up to 5 lb per 100 gal) (MgSO₄) (3 lb per 100 gal).

For chemicals not listed above, try the basis. Chemicals which are phytotoxic when used with oil.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its label.

Controls - whiteflies, leafminers, spider mites.

Crop	Insects Controlled
Asparagus	Leafminers, mites.
Beans	Whiteflies, leafminers.
Beets	Leafminers, mites.
Cabbage	Whiteflies, leafminers.
Cauliflower	Whiteflies, leafminers.
Celery	Whiteflies, leafminers.
Cucumber	Whiteflies, leafminers.
Eggplant	Whiteflies, leafminers.
Grapes	Mealybugs.
Lettuce	Whiteflies, leafminers.
Melons	Whiteflies, leafminers.
Peppers	Whiteflies, leafminers.
Potatoes	Whiteflies, leafminers.
Pumpkins	Whiteflies, leafminers.
Squash	Whiteflies, leafminers.
Tobacco	Whiteflies, leafminers.
Tomatoes	Whiteflies, leafminers.