

**HAZARDS TO HUMANS  
& DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. If swallowed, do not induce vomiting. Call a physician. Avoid breathing vapors and spray mists.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

**STORAGE AND DISPOSAL**

**STORAGE** — Keep drum tightly closed in storage to prevent entry of water. Do not freeze. Store drum on side. Roll drum before use if stored over-season.

**DISPOSAL** — Prohibitions Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

**Pesticide Disposal** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**General Consult** Federal, State or Local disposal authorities for approved alternative procedures.

**Method of Application**

Spray at no less than 400 psi pressure using ceramic spray nozzles (ALBUZ<sup>®</sup> ATR, lilac color or their equivalent). Do not apply this product through any type of irrigation system (chemigation).

**APPLICATION RATE**

From 25-150 gallons of diluted spray emulsion per acre depending on plant size. Thorough coverage is necessary. A tractor speed of 3.5-4 mph (6-7 kph) is recommended. Vary the quantity of spray applied by changing the number of nozzles used; not by changing either tractor speed or spray pressure.

**Timing of Applications**

Sprays should be initiated as soon as plants emerge. Plants should be sprayed weekly or twice weekly depending on level of infestation. Continue sprays through harvest.

For use as a fungicide: spray weekly or once every two weeks depending on the level of disease pressure.

1119 65-64-1  
**JMS**  
**Flower Farms, Inc.**

**JMS Stylet-Oil<sup>®</sup>**  
**AS AN INSECTICIDE/FUNGICIDE**

ACTIVE INGREDIENT:	BY WEIGHT
ALIPHATIC PETROLEUM DISTILLATE.....	97.1%
INERT INGREDIENTS.....	2.9%
<b>TOTAL:</b> .....	<b>100%</b>
Unsulfonated Residue of The Oil.....	99.1%
Flash Point.....	45° F

ACCEPTED

DEC 14 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 65564-1

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**  
IF SWALLOWED - Do not induce vomiting. Call a physician.  
IF INHALED - Avoid breathing of vapors and spray mist.  
IF ON SKIN - Wash immediately with soap and water.  
IF IN EYES - Wash immediately with copious amounts of water.

**TERMS**

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. All applicable directions, restrictions, and precautions on the label are to be followed. This labeling must be in the possession of the user at the time of the application of this material.

Manufactured by - Asgrow Florida Co., Plant City, FL 33566  
EPA Establishment No. - 14775-FL-01

JMS FLOWER FARMS, INC., 1105 25th Avenue,  
Vero Beach, FL 32960 (407) 567-9241  
EPA Registration No. - 65564-1

**55 GALLONS NET • U.S. STANDARD**

Do not enter treated areas without appropriate clothing. Do not spray near draft.

**PRE-HARVEST INTERVAL**

No interval required.

**ADDITIONAL COMMENTS ON SPRAYING:**  
Do not spray on wet foliage or if temperature is below 60°F (16°C).

**COMPATIBILITY INFORMATION**

**CHEMICAL COMPATIBILITY** - Add the oil to tank mix formulations as the last ingredient. JMS Stylet-Oil<sup>®</sup> can be used as a tank mix with acephate (Orthene), benomyl (Benlate), copper ammonium carbonate (CCN), copper hydroxide (Kocide), fenvalerate (Avalon XL), mancozeb, maneb, methamidophos (Monitor), metalaxyl (Ridomil) and triadimefon (Bayleton).

**DO NOT USE** the following chemicals with JMS Stylet-Oil<sup>®</sup> unless at least two weeks have elapsed between the use of oil and the chemical: anilazine (Dyrene), chlorothalinalol (Bravo), dicloran (Botran), Dikar, dinocap (Kamthane), organic tin compds. (Duter, Folpet), permethrin (Ambush, Pounce) and sulfur.

**DO NOT USE** JMS Stylet-Oil<sup>®</sup> in combination with highly ionized nutrient spray materials (Nutm-Leaf (20-20-20), Bayfolan). These materials can be sprayed separately from oil. JMS Stylet-Oil<sup>®</sup> is compatible with urea (up to 5 lb per 100 gal) and epsom salt (MgSO<sub>4</sub>) (3 lb per 100 gal).

**TANK MIX** JMS Stylet-Oil<sup>®</sup> only with those chemicals registered for use on the specific crop to be treated.

For chemicals not listed above, try them with oil on a small scale basis. Chemicals which are phytotoxic when used alone will be more phytotoxic when used with oil.

**Directions For Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Insects Controlled	Quarts JMS Stylet-Oil <sup>®</sup> per 100 gal water
Asparagus	Leafminers, mites,	3 qts
Beans	Whiteflies, leafminers, mites	
Beets	Leafminers, mites	
Cabbage	Whiteflies, leafminers, mites	
Cauliflower	Whiteflies, leafminers, mites	
Celery	Whiteflies, leafminers, mites	
Cotton	Whiteflies, leafminers, mites	
Field Corn	Earworms, Fall Armyworm, Corn Rootworm	
Sweet Corn	Earworms, Fall Armyworm, Corn Rootworm	
Cucumber	Whiteflies, leafminers, mites	
Eggplant	Whiteflies, leafminers, mites	
Grapes	Mealybugs	
Lettuce	Whiteflies, leafminers, mites	
Melons	Whiteflies, leafminers, mites	
Peppers	Whiteflies, leafminers, mites	
Potatoes	Whiteflies, leafminers, mites	
Pumpkins	Whiteflies, leafminers, mites	
Sugarbeets	Mites, Leafminer, Leafhopper	
Squash	Whiteflies, leafminers, mites	
Tobacco	Whiteflies, leafminers, mites	
Tomatoes	Whiteflies, leafminers, mites	

Chrysanthemum Diplomela Philodendron Poinsettia Roses	Leafminers, Mealybugs, Scale insects, Mites, Whiteflies	↑ 1 OZ ↓
---	---	----------------

Crop	Diseases Controlled	Quarts JMS Stylet-Oil® per 100 gal water
Beans	Rust	↑ 3 QTS ↓
Cucurbits	Alternaria leafspot, Gummy stem blight, Powdery mildew	
Grapes	Powdery mildew	
Okra	Powdery mildew	
Roses	Black spot, Powdery mildew	
Sugar beets	Cercospora	

