

REGULATORY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting. Call a physician. Avoid breathing vapors and spray mists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

STORAGE AND DISPOSAL

STORAGE — Keep drum tightly closed in storage to prevent entry of water. Do not freeze. Store drum on side. Roll drum before use if stored over-season.

DISPOSAL — Prohibitions Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide Disposal Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

General Consult Federal, State or Local disposal authorities for approved alternative procedures.

Method of Application

For insects/plant virus: Spray at no less than 400 psi pressure using ceramic spray nozzles (ALBUZ® ATR, lilac color or their equivalent). Do not apply this product through any type of irrigation system (chemigation).

For fungicide use: Application system used for insects/plant virus is not required. However, maximum coverage and least potential for phytotoxicity results from use of this system. When using conventional application equipment, spray for coverage of upper leaf surface.

Timing of Applications

For use as an insecticide: Sprays should be initiated as soon as plants emerge. Plants should be sprayed weekly or twice weekly depending on level of infestation. Continue sprays through harvest.

For use as a fungicide: spray weekly or once every two weeks depending on the level of disease pressure.

For use in mitigating aphid transmission of plant viruses: Initiate sprays when winged aphids first appear. Spray weekly or twice weekly depending on level of disease/vector pressure. For cucurbits, begin spraying when 50% of plants germinate. Cucurbits should be sprayed twice weekly during rapid growth phase.

Reg # 65564-1


PM-19

192



JMS Stylet-Oil®

ACTIVE INGREDIENT:	BY WEIGHT
ALIPHATIC PETROLEUM DISTILLATE.....	97.1%
INERT INGREDIENTS.....	2.9%
TOTAL:	100%
Unsulfonated Residue of Total Oil.....	99.1%
Flash Point.....	345° F



KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Do not induce vomiting. Call a physician.
IF INHALED - Avoid breathing of vapors and spray mist.
IF ON SKIN - Wash immediately with soap and water.
IF IN EYES - Wash immediately with copious amounts of water.

TERMS

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. All applicable directions, restrictions, and precautions on the label are to be followed. This labeling must be in the possession of the user at the time of the application of this material.

Manufactured by - Asgrow Florida Co., Plant City, FL 33566
EPA Establishment No. - 14775-FL-01

JMS FLOWER FARMS, INC., 1105 25th Avenue,
Vero Beach, FL 32960 (407) 567-9241
EPA Registration No. 65564-1

55 GALLONS NET • U.S. STANDARD

APPLICATION RATE

From 25-150 gallons of diluted spray emulsion per acre depending on plant size. Thorough coverage is necessary. A tractor speed of 3.5-4 mph (6-7 kph) is recommended. Vary the quantity of spray applied by changing the number of nozzles used; not by changing either tractor speed or spray pressure.

FIELD RE-ENTRY SAFETY INTERVAL
Do not enter treated areas without appropriate clothing until sprays have dried.
PRE-HARVEST INTERVAL
No interval required.
ADDITIONAL COMMENTS ON SPRAYING
Do not spray on wet foliage or if temperature is below 60° F (16° C)

COMPATIBILITY INFORMATION

CHEMICAL COMPATIBILITY - Add the oil to tank mix formulations as the last ingredient. JMS Stylet-Oil® can be used as a tank mix with acephate (Orthene), benomyl (Banlate), copper ammonium carbonate (CCN), copper hydroxide (Kocide), fenvalerate (Asana XL), mancozeb, maneb, methamidophos (Monitor), metalaxyl (Ridomil) and triadimelon (Bayleton).

DO NOT USE the following chemicals with JMS Stylet-Oil® unless at least two weeks have elapsed between the use of oil and the chemical anilazine (Dyrene), chlorothalinalol (Bravo), dicloran (Botran), Dikar, dinocap (Kamthane), organic tin cmpds. (Duter, Folpet), permethrin (Ambush, Pounce) and sulfur.

DO NOT USE JMS Stylet-Oil® in combination with highly ionized nutrient spray materials (Nut-Leaf) (20-20-20), Bayolan. These materials can be sprayed separately from oil. JMS Stylet-Oil® is compatible with urea (up to 5 lb per 100 gal) and epsom salt (MgSO₄) (3 lb per 100 gal).

TANK MIX JMS Stylet-Oil® only with those chemicals registered for use on the specific crop to be treated.

For chemicals not listed above, try them with oil on a small scale basis. Chemicals which are phytotoxic when used alone will be more phytotoxic when used with oil.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Insects Controlled	Quarts JMS Stylet-Oil® per 100 gal water
------	--------------------	--

Asparagus	Leafminers, mites,	3 qts
Bears	Whiteflies, leafminers, mites	
Beets	Leafminers, mites	
Cabbage	Whiteflies, leafminers, mites	
Cauliflower	Whiteflies, leafminers, mites	
Celery	Whiteflies, leafminers, mites	
Cotton	Whiteflies, leafminers, mites	
Field Corn	Earworms, Fall Armyworm, Corn Rootworm	
Sweet Corn	Earworms, Fall Armyworm, Corn Rootworm	
Cucumber	Whiteflies, leafminers, mites	
Eggplant	Whiteflies, leafminers, mites	
Grapes	Mealybugs	
Lettuce	Whiteflies, leafminers, mites	
Melons	Whiteflies, leafminers, mites	
Peppers	Whiteflies, leafminers, mites	
Potatoes	Whiteflies, leafminers, mites	
Pumpkins	Whiteflies, leafminers, mites	
Sugarbeets	Mites, Leafminer, Leafhopper	
Squash	Whiteflies, leafminers, mites	
Tobacco	Whiteflies, leafminers, mites	
Tomatoes	Whiteflies, leafminers, mites	

BEST AVAILABLE COPY

Chrysanthemum Difffenbachia Philodendron Poinsettia Roses	Leafminers, Mealybugs, Scale Insects, Mites, Whiteflies	↑ 1 OZ ↓
---	---	----------------

Crop	Diseases Controlled	Quarts JMS Stylet-Oil® per 100 gal water
Beans Cucurbits	Rust Alternaria leafspot, Gummy stem blight, Powdery mildew	↑ 3 QTS ↓
Grapes	Powdery mildew	
Okra	Powdery mildew	
Roses	Black spot, Powdery mildew	
Sugar beets	Cercospora	

Crop	Plant Viruses Controlled	Quarts JMS Stylet-Oil® per 100 gal water
Pepper	Potato virus Y, tobacco etch virus, pepper mottle virus	↑ 3 QTS ↓
Tomato	Potato virus Y, tobacco etch virus	
Potato	Potato virus Y, potato leafroll virus	
Cucurbits	Papaya ringspot virus, watermelon mosaic virus 2, cucumber mosaic virus, zucchini yellow mosaic virus	

272