PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting. Call a physician. Avoid breathing vapors and spray mists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

STORAGE AND DISPOSAL

STORAGE – Keep drum tightly closed in storage to prevent entry of water. Do not freeze. Store drum on side. Roll drum before use if stored over-season.

DISPOSAL – Prohibitions do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide Disposal Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

General Consult Federal, State or Local disposal authorities for approved alternative procedures.

METHOD OF APPLICATION

For insects/plant virus: Spray at no less than 400 psi pressure using ceramic spray nozzles (ALBUZ® ATR, lilac color or their equivalent). Do not apply this product through any type of irrigation system (chemigation).

For fungicide use: Application system used for insects/plant virus is not required. However, maximum coverage and least potential for phytotoxicity results from use of this system. When using conventional application equipment, spray for coverage of upper leaf surface.

APPLICATION RATE

From 25-150 gallons of diluted spray emulsion per acre depending on plant size. Thorough coverage is necessary. A tractor speed of 3.5-4 mph (6-7 kph) is recommended. Vary the quantity of spray applied by changing the number of nozzles used; not by changing either tractor speed or spray pressure.

TIMING OF APPLICATIONS

For use as an insecticide: Sprays should be initiated as soon as plants emerge. Plants should be sprayed weekly or twice weekly depending on level of infestation. Continue sprays through harvest.

For use as a fungicide: Spray weekly or once every two weeks depending on the level of disease pressure.

For use in mitigating aphid transmission of plant viruses: Initiate sprays when winged aphids first appear. Spray weekly or twice weekly depending on level of disease/vector pressure. For cucurbits, begin spraying when 50% of plants germinate. Cucurbits should be sprayed twice weekly during rapid growth phase.

JMS Flower Farms, Inc.

ACCEPTED
MAY 21 1993
For Agricultural Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 65564-1

JMS Stylet-Oil®

Agricultural Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 65564-1

ACTIVE INGREDIENT:	BY WEIGHT
ALIPHATIC PETROLEUM DISTILLATE	97.1%
INERT INGREDIENTS	2.9%
TOTAL	100%

Unsulfonated Residue of The Oil	99.1%
Flash Point	345°F

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Do not induce vomiting. Call a physician.
IF INHALED - Avoid breathing of vapors and spray mist.
IF ON SKIN - Wash immediately with soap and water
IF IN EYES - Wash immediately with copious amounts of water.

TERMS

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. All applicable directions, restrictions, and precautions on the label are to be followed. This labeling must be in the possession of the user at the time of the application of this material.

Manufactured by - Asgrow Florida Co., Plant City, FL 33565
EPA Establishment No. - 14775-FL-01

JMS FLOWER FARMS, INC., 1105 25th Avenue,
Vero Beach, FL 32960
(407) 567-9241 • FAX (407) 567-9394
EPA Registration No. - 65564-1

55 GALLONS NET • U.S. STANDARD

F[illegible]EPA 3/93

FIELD RE-ENTRY SAFETY INTERVAL

Do not enter areas without appropriate clothing until sprays have dried

PRE HARVEST INTERVAL

No interval required.

ADDITIONAL COMMENTS ON SPRAYING

Do not spray on wet foliage or if temperature is below 60° F (16° C).

COMPATIBILITY INFORMATION

CHEMICAL COMPATIBILITY: Addition to tank mixformulations as the last ingredient. JMS Stylet-Oil® can be mixed with captafol (Orthene), benomyl (Benlate), copper ammonium carbonate (CCN), copper hydroxide (Kocide), fenvalerate (Asana XL), mancozeb, maneb, methamidophos (Monitor), metalaxyl (Ridomil), and triadimefon (Bayleton).

DO NOT USE the following chemicals with JMS Stylet-Oil® unless at least two weeks have elapsed between the use of oil and the chemical: aniazine (Dyrene), chlorothalonil (Bravo), diclobutrazol (Bolran), Dikar, dinocap (Karathane), organotin cmpds. (Duter, Folpel), permethrin (Ambuth, Pounce) and sulfur.

DO NOT USE JMS Stylet-Oil® in combination with highly ionized nutrient spray materials (Nutri-Leaf) (20-20-20), Bayfolan. These materials can be sprayed seperately from oil. JMS Stylet-Oil® is compatible with urea (up to 5 lb. per 100 gal.) and epsom salt ($MgSO_4$) (3 lb. per 100 g[illegible]

Tank mix JMS Stylet-Oil® only with those chemicals registered for use on the specific crop to be treated.,

For chemicals not listed above try them with oil on a small scale basis Chemicals which are phytotoxic when used alone will be more phytotoxic when used with oil.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR FUNGICIDE USE:

Crop	Plant Diseases Controlled	Rate JMS Stylet-Oil® per 100 gal water
Beans Cucurbits Okra Roses Sugar beets	Rust Alternaria leafspot, Gummy stem blight, Powdery mildew Powdery mildew Black spot, Powdery mildew Cercospora	3 QTS
Grapes	Powdery mildew	Use 1 to 2 Gal

FOR PLANT VIRUS USE:

Crop	Plant Viruses Controlled	Quarts JMS Stylet-Oil® per 100 gal water
Pepper Tomato Potato Cucurbits	Potato virus Y, tobacco etch virus, pepper mottle virus Potato virus Y, tobacco etch virus Potato virus Y, potato leafroll virus Papaya ringspot virus, watermelon mosaic virus 2, cucumber mosaic virus, zucchini yellow mosaic virus	3 QTS

FOR INSECTS:

Crop	Insects Controlled	Quarts JMS Stylet-Oil® per 100 gal water
Asparagus Beans Broccoli Cabbage Cucumber Celery Collards Cucumber Field Corn Sweet Corn Eggplant Lettuce Melons Peppers Potatoes Soybeans Spinach Squash Tomatoes	Leafminers, whitefly Whiteflies, leafminers, mites Leafminers, whitefly Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Earworms, Fall Armyworm, Corn Rootworm Earworms, Fall Armyworm, Corn Rootworm Whiteflies, leafminers, mites Mites Leafminers Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites Mites, Leafminers, Leafhopper Whiteflies, leafminers, mites Whiteflies, leafminers, mites Whiteflies, leafminers, mites	3 QTS

FOR ORNAMENTAL USE: Insects Controlled	RATE: JMS Stylet-Oil® per 1 gal water
Chrysanthemum, Gladiolus, Philodendron, Poinsettia, Roses — Leafminers, Mealybugs, Scale Insects, Mites, Whiteflies	1 OZ.