OCT 6 2003

Mr. Gary Gaumer President Preserve International 315 Wall Street, Suite 1 Chico, CA 95928

SUBJECT:

Synergize

EPA Registration Number: 66171-7 Your Amendment Dated: July 3, 2003 EPA Received Date: July 9, 2003

Dear Mr. Gary Gaumer:

The amendment referred to above, submitted in connection with registration under FIFRA as amended, is acceptable provided that you:

- 1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
- A. You have cited another product containing glutaraldehyde to support your addition of fogging directions to this label. Thus your directions must be in agreement with that label and meet Agency standards as to acceptable language for fogging.
- 1. Delete the entire section beginning: "Directions for Fogging: Synergize is recommended..." and ending "with potable water prior to reuse."
- 2. Add fogging directions to the section "Directions for Use in Poultry Houses" to read exactly as follows:

"Prior to fogging setters or hatchers, food products and packaging must be removed from the room or carefully protected. After cleaning, fog desired areas using one quart per 1000 cubic feet of room area with a product solution containing 0.25 to 1 ounce per gallon of water. Vacate all areas of personnel and poultry and feeds for a minimum of 2 hours after fogging. All food contact surfaces, including feed troughs, automatic feeders, and fountains must be disinfected followed by treatment with soap and water or detergent and rinsed with potable water before

| CONCURRENCES | | | | | | | | | |
|--------------|-----|---|---|---|--|--------|---------------|--|--|
| SYMBOL 75/00 | n | | | | | | | | |
| SURNAME) Jo | ايد | 1 | 1 |] | | | | | |
| DATE 10/6/0 | 3 | | | | | | ************* | | |
| | | | | | | OFFICE | AL EU E CORY | | |

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

*11 S. Gonzament Printing Office: 1982 vs. 620-866/40872

2/3

Note: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTION OF ROOM AND MACHINE SURFACES."

- 3. These are the only changes which have been accepted for this label at this time.
- 4. Submit two (2) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records. Should you have any questions concerning this letter, please contact Tracy Lantz at (703) 308-6415.

Sincerely,

Velma Noble

Product Manager (31)

Regulatory Management Branch I Antimicrobials Division (7510C)

Enclosure: Stamped Label

7510C·T Lantz·10/6/03·66171-7j

CONCURRENCES

SYMBOL | SURNAME | DATE

DESCRIPTION

SYNERGIZE is a multi-purpose disinfectant-cleaner. SYNERGIZE provides a broad spectrum, kill of gram negative and gram positive microorganisms. SYNERGIZE provides effective disinfecting and cleaning in one operation. SYNERGIZE is effective in warm, soft or hard water (as much as 1000 ppm CaCO₃) and performs especially well in areas containing 5% arms.

EYNERGIZE is for une no glazed ceramic and tile surfaces, stainless steet, aluminum, chrome, galvanized metal tin, brass zinc, glass, hard non-porous freated wood, polyethylene polypropytene. PVC (polyvinyl chloride), vinyl, tiberglass, viton, ethylene propylene, nitrile, acrylic and polyurethane.

DIRECTIONS FOR USE:

It is in violation of Federal Law to use this product in a manner inconsistent with its labeling. Registered for disinfecting and cleaning hard non-porous surfaces. Not for use on feed, food or litter.

SYNERGIZE cleaner-disintectant is specifically formulated for industrial use in the following Animal and Poultry Housing facilities: Turkey, Breeder, Grow Out, Fryer, Layer, Hatcheries and Swine Producing Facilities as well as the following equipment in these facilities: Waterers, Feeders, Hauling, Equipment, Dressing, Plants, Loading, Equipment, Farrowing Barns, Nursenes, Blocks, Creep Areas, and Chutes. This product is also for use in Farm Premise, Equine Facilities, Veal Barns, Ratile Facilities (Ostrich, Emu and Rhea Facilities), Call Pens and associated Farm Vehicles.

USE DIRECTIONS:

To disinfect and clean hard non-porous surfaces of walls, floors, tables, sinks, refrigerator exteriors, telephone booths and garbage cans in non-animal areas associated with the sites noted above. Apply a solution of 1/2 ounce SYNERGIZE per gallon water (dilute 1:256) to precleaned surfaces. Apply with cloth, sponge, mop or mechanical coarse spray device, making sure all surfaces are wetted thoroughly. Allow surfaces to remain wet for 10 minutes. All surfaces must be rinsed with clean water and allowed to dry before reuse.

BOOT/SHOE WASH: To prevent tracking harmful organisms into animal areas, shoe baths containing one inch of freshly made disinfecting solution should be placed at all entrances to building. Scrape waterproof shoes or boots and place in a 1/2 ounce SYNERGIZE per gallon water solution. Allow the solution to come in contact with the boot/shoe for 10 minutes. Change the disinfecting solution when solution becomes visibly dirty.

VEHICLES. To disinfect the non-porous, hard surfaces of vehicles use a solution of 1/2 ounce SYMERGIZE per gallon of water. Apply using a high pressure or course spray system. Leave all treated surfaces exposed to solution for 10 minutes, or more and allow to air dry. All treated surfaces that will contact feed or drinking water must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

DIRECTIONS FOR USE IN SWINE PRODUCING FACILITIES:

- 1) Remove all animals and feed from the premises, vehicles and enclosures.
- 2) Empty all troughs, racks, and other feeding and watering appliances.
- Remove all gross soil, bedding, manure, caked on dirt, spider webs, etc. from walls to ceilings of bams, pens, stalls, chutes, and other facilities occupied or traversed by hogs or other animals.
- 4) Thoroughly clean all surfaces with soap or detergent and rinse with water
- Thoroughty saturate all surfaces with a solution of 1/2 oz SYNERGIZE per gallon of water (1 256), using conventional spraying methods.
- Allow all treated areas to remain in contact with this solution for at least 10 minutes.
- 7)Immerse equipment used in handling and restraining animals, forks, shovels, and scrapers used for removing manure in SYNERGIZE at a dilution of 1:256. Allow to remain in contact for at least 10 minutes.
- Ventitate buildings, and other closed spaces. Do not house swine or other livestock until treatment has been absorbed, set, or dried
- 9)Thoroughly scrub treated feed racks, troughs, automatic leeders, fountains, waterers and equipment which may come in contact with food with soap or detergent, and rinse with potable water before reuse.

DIRECTIONS FOR FOGGING:

Synergize is recommended for use in fogging or atomizing (wet mist) operations as an adjunct to acceptable manual cleaning and disinfecting of room surfaces. Before fogging, all food and food packaging items must be removed or carefully protected. Use at a dilution of 1/2 - 3 ounces per gallon. Check with a Preserve International representative to determine an effective fogging schedule. Vacate the area of all personnel during fogging. Treated food contact surfaces must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.



SYNERGIZE

CLEANER-DISINFECTANT

Synergize cleaner-disinfectant is specifically formulated for industrial use in the following Animal and Poultry Housing facilities: Turkey, Breeder, Grow Out, Fryer, Layer, Hatcheries and Swine Producing Facilities as well as the following equipment in these facilities: Waterers, Feeders, Hauling Equipment, Loading Equipment, Farrowing Barns, Nurseries, Blocks, Creep Areas, and Chutes. This product is also for use in Farm Premise, Equine Facilities, Veal Barns, Ratite Facilities (Ostrich, Emu and Rhea Facilities), Calf Pens and associated Farm Vehicles.

Staphylococcus aureus A.T.C.C. 6538 Salmonella choleraesuis A.T.C.C. 10708 Pseudomonas aeruginosa A T C C 15442 Aspergillus fumigatus A T C C 36607

ACTIVE INGREDIENTS

| Alkyl (Cn 67%, Cn 25%, Cn 7%, Cn 1%) dimethyl benzyl ammonium chloride 26.00% | |
|---|--|
| Glutaraldehyde | |
| INERT INGREDIENTS | |
| TOTAL | |

KEEP OUT OF REACH OF CHILDREN DANGER

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
See First Aid statement and additional precautions on side panel

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed, by storage and disposal.

PESTICIDE STORAGE:

Store away from heat. Do not transfer to another container. Do not reuse empty container.

PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Registration No.: 66171-7

EPA Establishment No.: 66171-TN-ORCCEPTED

66171-CA ANICOMMENTS 1 922 in EPA Letter Dated:

Patent Number: 5,891,922 in EPA I

NFPA HAZARD CLASSIFICATION HEALTH FIRE REACTIVITY SPECIFIC

Under Precipic Pederal Insecticide, and Rodenticide Act as

Zephyr Cove, NV 89448 (209) 664-1607 • www.preserveinternational.com for the pesticide, registered under EPA Reg. No. registered under EPA Reg. No.

DIRECTIONS FOR USE IN FARM PREMISES, EQUINE FACILITIES, YEAL BARNS, RATITES FACILITIES, CALF PENS AND VEHICLES:

BARNS, RATITES FACILITIES, CALF PENS AND VEH

1) Do not use in milking stalls, milking parlors, or milk houses

2) Hemove all animals and feed from the premises, vehicles, and enclosures

 Remove all gross soil, bedding, manure, caked on dirt, spider webs, etc. from floors, surlaces, and from walls to ceiting of barns, pens, stalls, chutes, and other facilities occupied or traversed by hogs or other animals.

4) Empty all troughs, racks, and other feeding aixt watering appliances

5) Thoroughly clean all surfaces with కుండి or detergent and rinse with viater * * ? ?

6) Thoroughly saturate all surfaces with a solution of 1/2 oz SYNEGGE per gallon of water (1.256), using conventional engaging method. Allow to remain inhontact with this solution for at least 10 minutes.

7) Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing manure in a solution of 1/2 oz. SYNERGIZE per gallon of water (1:256). Allow to remain in contact for 10 minutes.

Source to the minutes

Source to the minutes of the

9) Thoroughly scrub treated feed racks, troughs, automatic deducts, fountains, waterers and equipment which may come in contact with ford with roap or detergent, and rinse with potable water before reuse.

DIRECTIONS FOR USE IN POULTRY HOUSE:

1) Remove all poultry and feeds from premises, trucks, coops, and crates.

 Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.

3) Empty all troughs, racks, and other feeding and watering appliances.

4) Thoroughly clean all surfaces with soap or detergent and rinse with water.

 Saturate surfaces with 1/2 ounce SYNERGIZE per gallon water (dilute1 256) for a period of 10 minutes.

6) Ventilate building, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.

7) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, waterers and equipment which may come in contact with food with soap or detergent, and rinse with potable water before reuse.

PRECAUTIONARY STATEMENTS Hazards to humans and domestic animals

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. May be latal if inhaled. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Wear goggles or face should be wear a dust/mist/litering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N, P, R, or HE prefilter. Wear protective clothing and rubber gloves. Prolonged or frequent repeated skin contact may cause allergic reaction in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

FIRST AID

Net Contents.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes, remove contact lenses, if present, after the first 5 minutes, then continue mising eye; call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing; rinse skin immediately with plenty of water for 15-20 minutes; call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible, call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice; have person sip a glass of water if able to swallow; do not induce vomiting unless told to do so by the poison control center or doctor; do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1 (800) 255-3924 for emergency medical treatment information. For information on this pesticide (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1 (800) 858-7378

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage Measures against circulatory shock, respiratory depression and convulsion may be needed

ENVIRONMENTAL HAZARDS: This product is toxic to lish. Do not apply directly to water Do not contaminate water by cleaning of equipment or disposal of wastes. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other water unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

U.S. Gallons

Batch No.